

Interface regularity for semilinear one-phase problems

Original

Interface regularity for semilinear one-phase problems / Audrito, Alessandro; Serra, Joaquim. - In: ADVANCES IN MATHEMATICS. - ISSN 0001-8708. - 403:(2022). [10.1016/j.aim.2022.108380]

Availability:

This version is available at: 11583/2985054 since: 2024-01-14T14:01:13Z

Publisher:

Elsevier

Published

DOI:10.1016/j.aim.2022.108380

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)




ELSEVIER

Contents lists available at ScienceDirect

Advances in Mathematics

www.elsevier.com/locate/aim



Interface regularity for semilinear one-phase problems [☆]

Alessandro Audrito ^{*}, Joaquim Serra

Department of Mathematics, ETH Zurich, Rämistrasse 101, 8092 Zurich, Switzerland

ARTICLE INFO

Article history:

Received 18 October 2021

Received in revised form 8 February 2022

Accepted 24 March 2022

Available online 14 April 2022

Communicated by C. Fefferman

MSC:

35J61

35J75

35J20

35R35

Keywords:

Blow-down

Improvement of flatness

One-phase free boundary problem

ABSTRACT

We study critical points of a one-parameter family of functionals arising in combustion models. The problems we consider converge, for infinitesimal values of the parameter, to Bernoulli's free boundary problem, also known as one-phase problem. We prove a $C^{1,\alpha}$ estimates for the “interfaces” (level sets separating the *burnt* and *unburnt* regions). As a byproduct, we obtain the one-dimensional symmetry of minimizers in the whole \mathbb{R}^N , for $N \leq 4$, answering positively a conjecture of Fernández-Real and Ros-Oton. Our results are to Bernoulli's free boundary problem what Savin's results for the Allen-Cahn equation are to minimal surfaces.

© 2022 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

[☆] This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement 892017 (LNLFB-Problems). JS has been supported by the European Research Council (ERC) under the grant agreement 948029.

^{*} Corresponding author.

E-mail addresses: alessandro.audrito@math.ethz.ch (A. Audrito), joaquim.serra@math.ethz.ch (J. Serra).

1. Introduction

In this paper we study critical points of the following (non-convex) energy functional

$$\mathcal{E}_\varepsilon(u, \Omega) := \int_{\Omega} |\nabla u|^2 + \Phi_\varepsilon(u) \, dx, \quad (1.1)$$

where $\varepsilon \in (0, 1]$ is a parameter, $\Omega \subset \mathbb{R}^N$ some open domain, and

$$\Phi_\varepsilon(t) := \Phi(t/\varepsilon) \quad (1.2)$$

$$\Phi(t) := \begin{cases} \int_0^t \beta(\tau) \, d\tau & \text{for } t \geq 0 \\ 0 & \text{for } t < 0, \end{cases} \quad (1.3)$$

for some given function $\beta \in C_c^\infty([0, +\infty))$ satisfying

$$\beta \geq 0, \quad \beta(0) = 0, \quad \beta'(0) > 0, \quad \int_0^\infty \beta = 1. \quad (1.4)$$

When $\varepsilon = 1$, $\mathcal{E}_1(u, \Omega)$ will be sometimes denoted by $\mathcal{E}(u, \Omega)$. The assumption $\beta'(0) > 0$ is made for simplicity, but in all our main results it could be actually replaced by $\liminf_{\tau \downarrow 0} \beta(\tau) \tau^{-p} > 0$ for some $p \in (1, \infty)$ —see Remark 2.3.

These type of functional arises in combustion models (e.g. flame propagation) [12,4,13,27,21], and were studied in detail in the book of Caffarelli and Salsa [11].

Connection to the one-phase problem

Due to the assumptions on Φ , as $\varepsilon \downarrow 0$, the energy \mathcal{E}_ε formally converges towards

$$\mathcal{E}_0(u, \Omega) := \int_{\Omega} |\nabla u|^2 + \chi_{\{u>0\}} \, dx. \quad (1.5)$$

Critical points of \mathcal{E}_0 are solutions to Bernoulli's (or one-phase) free boundary problem:

$$u \geq 0, \quad \Delta u = 0 \quad \text{in } \{u > 0\}, \quad \partial_n u = 1 \quad \text{on } \partial\{u > 0\}, \quad (1.6)$$

where n is the inwards unit normal to $\partial\{u > 0\}$. The regularity of solutions and free boundaries for minimizers of \mathcal{E}_0 has been extensively studied in [1,2,6–8,10,15,16,19,20,24,25] (see also the treatment given in [23]). The convergence of \mathcal{E}_ε towards \mathcal{E}_0 as $\varepsilon \downarrow 0$ is not merely formal: as proven in [11, Theorem 1.15], sequences of minimizers u_{ε_k} of $\mathcal{E}_{\varepsilon_k}$ converge as $\varepsilon_k \downarrow 0$ (and up to subsequences) towards minimizers of the functional \mathcal{E}_0 .

A conjecture “à la De Giorgi”

By the results in [10,19], it is known that every minimizer $u_0 : \mathbb{R}^N \rightarrow [0, \infty)$ of \mathcal{E}_0 in \mathbb{R}^N must have one-dimensional symmetry in dimensions $N \leq 4$, while this fails for $N \geq 7$ (see [16]). On the other hand if $u : \mathbb{R}^N \rightarrow \mathbb{R}_+$ is a minimizer of $\mathcal{E} = \mathcal{E}_1$ in \mathbb{R}^N , then the *blow-down sequence* $u_{\varepsilon_k}(x) := \varepsilon_k u(x/\varepsilon_k)$, $\varepsilon_k \downarrow 0$, are minimizers of $\mathcal{E}_{\varepsilon_k}$ in \mathbb{R}^N . Since by [11, Theorem 1.15] u_{ε_k} converges (up to subsequence and uniformly in every compact subset of \mathbb{R}^N) to some entire minimizer of \mathcal{E}_0 , every blow-down of u must be one-dimensional if $N \leq 4$. By analogy with De Giorgi’s conjecture for the Allen-Cahn equation (see for instance [14,22]), Fernández-Real and Ros-Oton raised the following

Conjecture 1.1 ([18]). *Let $N \leq 4$ and $u : \mathbb{R}^N \rightarrow \mathbb{R}_+$ be a minimizer of \mathcal{E} in \mathbb{R}^N (see Definition 1.2 below). Then, u must be of the form*

$$u(x) = v(\nu \cdot x - l) \quad \text{where } v(t) = \psi^{-1}(t) \quad \text{for } \psi(z) := \int_1^z \frac{d\zeta}{\sqrt{\Phi(\zeta)}}, \quad (1.7)$$

for some $\nu \in \mathbb{S}^{N-1}$ and $l \in \mathbb{R}$.

The results in this paper answer positively this conjecture.

Minimizers and critical points

We define next minimizer and critical point of \mathcal{E}_ε .

Definition 1.2. Let $\Omega \subseteq \mathbb{R}^N$ be some open domain and let $\varepsilon > 0$. We say that $u_\varepsilon \in H_{loc}^1(\Omega)$ is a *minimizer* of (1.1) in Ω if for every $V \subset\subset \Omega$ and for every $\xi \in H_0^1(V)$ we have

$$\mathcal{E}_\varepsilon(u_\varepsilon, V) \leq \mathcal{E}_\varepsilon(u_\varepsilon + \xi, V).$$

Definition 1.3. Let Ω , N and $\varepsilon > 0$, as in Definition 1.2. We say that $u_\varepsilon \in H_{loc}^1(\Omega)$ is a *critical point* of (1.1) in Ω if for every $V \subset\subset \Omega$ and for every $\xi \in H_0^1(V)$ we have

$$\left. \frac{d}{dt} \right|_{t=0} \mathcal{E}_\varepsilon(u_\varepsilon + t\xi, V) = 0 \quad \Leftrightarrow \quad \int_V 2\nabla u_\varepsilon \cdot \nabla \xi + \Phi'_\varepsilon(u_\varepsilon)\xi \, dx = 0.$$

Notice that (after integration by parts) any critical point u_ε of satisfies

$$\Delta u_\varepsilon = \frac{1}{2}\Phi'_\varepsilon(u_\varepsilon), \quad (1.8)$$

in the weak sense. Since Φ' is smooth and bounded, by elliptic regularity and the standard “bootstrap argument” for semilinear equations, any critical point is locally smooth (with estimates which degenerate in principle as $\varepsilon \downarrow 0$) and hence satisfies (1.8) in the classical sense.

New results

We describe next the main results of the paper. Our main contribution is the following rigidity results for critical points of \mathcal{E} in \mathbb{R}^N which are “asymptotic” to $(\nu \cdot x)_+$ at very large scales. In its statement (and in the rest of the paper) we use the following convenient notation for inclusion of sets: we write “ $X \subset Y$ in Z ” when $X \cap Z \subset Y \cap Z$.

Theorem 1.4. *Let Φ be as in (1.3)-(1.4). There exist constants ϑ_1 and ϑ_2 depending only on Φ such that the following holds. Let $u : \mathbb{R}^N \rightarrow \mathbb{R}_+$ be a critical point of \mathcal{E} in \mathbb{R}^N .*

Assume there exist $\nu \in \mathbb{S}^{N-1}$ and sequences $R_k \uparrow \infty$ and $\delta_k \downarrow 0$ such that

$$|u - (\nu \cdot x)_+| \leq \delta_k R_k \quad \text{in } B_{R_k}, \quad (1.9)$$

and

$$\{\nu \cdot x \leq -\delta_k R_k\} \subset \{u \leq \vartheta_1\} \subset \{u \leq \vartheta_2\} \subset \{\nu \cdot x \leq \delta_k R_k\} \quad \text{in } B_{R_k}. \quad (1.10)$$

Then u is of the form (1.7).

On the other hand, building on the results of [11, Chapter 1] (and introducing new ideas) we establish the following

Proposition 1.5. *Let Φ be as in (1.3)-(1.4) and let ϑ_1 and ϑ_2 be the constants from Theorem 1.4. Let $u : \mathbb{R}^N \rightarrow \mathbb{R}_+$ be a minimizer of \mathcal{E} in \mathbb{R}^N which is not identically 0. Then, for every sequence $R_k \uparrow \infty$ there exists a subsequence R_{k_ℓ} , a 1-homogeneous minimizer u_0 of \mathcal{E}_0 in \mathbb{R}^N — also not identically zero — and a sequence $\delta_\ell \downarrow 0$ such that*

$$|u - u_0| \leq \delta_\ell R_{k_\ell} \quad \text{in } B_{R_{k_\ell}}, \quad (1.11)$$

and

$$\begin{aligned} \{x : \text{dist}(x, \{u_0 > 0\}) \geq \delta_\ell R_{k_\ell}\} &\subset \{u \leq \vartheta_1\} \subset \{u \leq \vartheta_2\} \\ &\subset \{x : \text{dist}(x, \{u_0 = 0\}) \leq \delta_\ell R_{k_\ell}\} \quad \text{in } B_{R_{k_\ell}}. \end{aligned} \quad (1.12)$$

Combining Theorem 1.4, Proposition 1.5, and using the classification results for 1-homogeneous minimizers of \mathcal{E}_0 of [10,19] we obtain

Corollary 1.6. *Conjecture 1.1 holds true.*

2. Overview of the proofs and organization of the paper

The proof of Theorem 1.4 is split into several intermediate steps, some of them having independent interest. The main step (and our main contribution) is establishing an

“improvement of flatness” result for critical points of \mathcal{E} that we state below. Before that, we need to introduce two positive constants ϑ_1 and ϑ_2 , with $\vartheta_1 < \vartheta_2$ and depending only on Φ , that will appear throughout the paper. Under our assumptions on Φ —see (1.2)-(1.4)— we can choose positive constants ϑ_1 , ϑ_2 , and c_1 , such that the following holds:

$$\begin{cases} \Phi = 0 & \text{in } (-\infty, 0], \quad \Phi = 1 & \text{in } [\vartheta_2, \infty), \\ \frac{1}{c_1}u \leq \frac{1}{2}\Phi'(u) \leq c_1u, & \forall u \in [0, \vartheta_1]. \end{cases} \quad (2.1)$$

We can now give the statement of our “improvement of flatness” result.

Theorem 2.1. *Let Φ be as in (1.3)-(1.4) and let ϑ_1 and ϑ_2 as in (2.1). Fix $\gamma \in (0, 1)$. There exist constants $\delta_0 > 0$ and $\varrho_0 \in (0, 1/4)$ depending only on N and Φ , such that the following holds. For every $R > 0$, every $\delta \in (0, \delta_0]$, every $\varepsilon/R \in (0, \delta^2)$, and every critical point u_ε of (1.1) in $B_R \subset \mathbb{R}^N$ satisfying*

$$u_\varepsilon(0) \in [\vartheta_1\varepsilon, \vartheta_2\varepsilon] \quad (2.2)$$

and

$$\begin{array}{ll} u_\varepsilon(x) - x_N \leq \delta R & \text{in } B_R \cap \{u_\varepsilon \geq \vartheta_1\varepsilon\} \\ -\delta R \leq u_\varepsilon(x) - x_N & \text{in } B_R, \end{array} \quad (2.3)$$

there exists $\nu \in \mathbb{S}^{N-1}$ such that

$$\begin{array}{ll} u_\varepsilon(x) - \nu \cdot x \leq \delta \varrho_0^{1+\gamma} R & \text{in } B_{\varrho_0 R} \cap \{u_\varepsilon \geq \vartheta_1\varepsilon\} \\ -\delta \varrho_0^{1+\gamma} R \leq u_\varepsilon(x) - \nu \cdot x & \text{in } B_{\varrho_0 R} \end{array} \quad (2.4)$$

with

$$|\nu - e_N| \leq \sqrt{2N}\delta. \quad (2.5)$$

Let us discuss some key aspects in the statement of Theorem 2.1:

Assumption (2.2) must be thought as the analogue of asking 0 to be a free boundary point in the one-phase setting ($\varepsilon = 0^+$). Indeed, on the one hand it follows from the definition of ϑ_2 that u_ε is harmonic in $\{u_\varepsilon > \vartheta_2\varepsilon\}$. On the other hand, using the definition of ϑ_1 we will show (cf. Lemma 3.6) that u_ε has “exponentially small size in ε ” inside $\{u_\varepsilon < \vartheta_1\varepsilon\}$. Consequently, the “fat hypersurface” $\{\vartheta_1\varepsilon < u_\varepsilon < \vartheta_2\varepsilon\}$ is really analogous the free boundary in the one-phase setting.

Assumption (2.3) and conclusion (2.4) must be thought, respectively, as a δ -flatness property of u_ε at scale $R > 0$ and a $(\varrho_0^\gamma \delta)$ -flatness property at scale $\varrho_0 R$. In our framework this turns out to be the appropriate notion of δ -flatness. As it is customary, the flatness is a dimensionless parameter: Roughly speaking, it measures the ratio between

$\min_{e \in \mathbb{S}^{N-1}} \text{dist}(\{\vartheta_1 \varepsilon < u_\varepsilon < \vartheta_2 \varepsilon\} \cap B_R, \{e \cdot x = 0\} \cap B_R)$ and R . With respect to [15], we remark that in (2.3)-(2.4) inequality from above is not required to hold in $\{u_\varepsilon > 0\}$, but only in $\{u_\varepsilon \geq \vartheta_1 \varepsilon\}$ (otherwise the result would be empty since non-zero solutions to our semilinear PDE are everywhere positive!).

The conclusion of the theorem can be phrased as an “improvement of flatness”: if u_ε is δ -flat at scale R (for small values of ε and δ), it is $(\varrho_0^\gamma \delta)$ -flat at scale $\varrho_0 R$.

We now say a few words about the proof of Theorem 2.1. In some sense, this proof is an “interpolation” of the proofs of De Silva in [15] and Savin in [22] (although an additional “sliding method” step in the spirit of Berestycki, Caffarelli, Nirenberg [3] is also needed, by similar reasons as in [17]). Indeed, our goal is to generalize the proof of De Silva [15] for the one-phase free boundary problem to the setting of critical points of $\mathcal{E}_\varepsilon(\cdot, \mathbb{R}^N)$. But since we need to go from a scaling invariant problem to a non-scaling invariant semilinear problem, there is an obvious analogy with what Savin did in his celebrated paper [22]. In this work Savin proved a version of the De Giorgi’s improvement of flatness for area-minimizing hypersurfaces (a scaling invariant problem), in the framework of energy minimizers of the Allen-Cahn equation (a semilinear PDE).

Both Savin’s and De Silva’s proofs follow a “small perturbations” approach (linearization around flat solutions). In both cases — although for different reasons — the *deviation* between an almost-flat solution and the flat one which best approximates it, is found to be an “almost-harmonic” function. Further, in both proofs, the quadratic decay of harmonic function towards their linear Taylor expansion is somehow transferred to the almost-flat solutions in order to obtain the improvement of flatness property. To accomplish this, both proofs use a delicate compactness argument, where deviations converge in C^0 towards some limit function which is proved to be harmonic in the viscosity sense. This type of argument requires some C^α estimate, or *improvement of oscillation estimate*, which guarantees the compactness in C^0 (via Arzelà-Ascoli) of the sequences the deviations.

In our proof we also need such improvement of oscillation estimate, and finding an appropriate statement we could use in our setting turned out to be not easy at all! Indeed, in a first “naive approximation”, one could try to extend De Silva’s improvement of oscillation ([15, Theorem 3.1]) to the semilinear setting as follows:

Lemma 2.2. *Let v_ε be the solution of (1.8) in \mathbb{R} satisfying $v_\varepsilon(0) = \vartheta_1 \varepsilon$ (see Lemma 3.1, part (i)). There exist $\delta_0, c_0 \in (0, 1)$ and $\theta_0 \in (0, 1)$ depending only on N, Φ such that the following holds. For every $R > 0$, every $\delta \in (0, \delta_0)$, every $a \in \mathbb{R}$ and $b \leq 0$ such that $a + |b| = \delta R$, every $\varepsilon/R \in (0, c_0 \delta)$ and every critical point u_ε of (1.1) in B_R satisfying*

$$v_\varepsilon(x_N - a) \leq u_\varepsilon(x) \leq v_\varepsilon(x_N - b) \quad \text{in } B_R, \quad (2.6)$$

there exist $a' \in \mathbb{R}$, $b' \leq 0$ such that

$$\begin{aligned}
v_\varepsilon(x_N - a') &\leq u_\varepsilon(x) \leq v_\varepsilon(x_N - b') \quad \text{in } B_{R/4}, \\
b &\leq b' \leq a' \leq a, \\
a' + |b'| &\leq \theta_0(a + |b|).
\end{aligned}$$

Lemma 2.2 is true.¹ Unfortunately, it seems useless: the reason is that we cannot exclude the existence of minimizers \mathcal{E}_ε in B_{2R} which are $\frac{\delta}{100}$ -close to $(x_N)_+$ —with $\varepsilon > 0$ and ε/δ arbitrarily small— but failing to satisfy (2.6).

Lemma 6.3, where the δ -shifts of v_ε are replaced by δ -shifts of two suitable $1D$ super- and subsolutions, is the right replacement to the previous naive statement. We construct these useful super and subsolutions in Lemma 3.1 part (ii) and (iii). Since they play a very important role in the paper, we devote the entire Section 3 to the classification of $1D$ (super- and sub-) solutions and the study of their properties. We do not give yet the statement of Lemma 6.3 because such preliminaries are needed.

Let us remark that this notion of δ -flatness consisting in “being trapped” between δ -shifts of $1D$ super and subsolutions is essentially equivalent to the notion (2.3) when $\varepsilon \in (0, \delta^2)$ —this is actually the reason behind this nonlinear relation between ε and δ in the statement of Theorem 2.1. Definition 6.1 and Lemma 6.2 establish this essential equivalence, when $\varepsilon \in (0, \delta^2)$, of these two notions of flatness which are used throughout the paper.

Last, but not least, in order to prove Theorem 1.4 we need to be able to apply our new improvement of flatness result (Theorem (2.1)) to $u_\varepsilon := \varepsilon u(\cdot/\varepsilon)$ where u is a minimizer of \mathcal{E}_1 in \mathbb{R}^N , $N \leq 4$. To do so, first we need to show that the assumption (2.3) will be satisfied—for some $\delta = \delta_0$ and $R = 1$ —when ε is taken sufficiently small. This part essentially combines previous results in [10,19] and [11] (although some improvements are needed) and it is contained in Section 4. However there is an important difference with respect to [22] that is related to our assumption $\varepsilon/R < \delta^2$ in Theorem 2.1. Indeed, in contrast with the Allen-Cahn setting (where ε and δ are comparable and the analogue of Theorem 1.4 is a corollary of the improvement of flatness), in our setting Theorem 1.4 does not follow as a direct consequence of Theorem 2.1. The reason is the following: suppose you want to apply Theorem 2.1 iteratively (in balls of radius $R\varrho_0^{-i}$) to an entire minimizer u of \mathcal{E}_1 , starting from a huge ball B_R (for which u is δ_0 -flat). Then, at a mesoscale $1 \ll R' \ll R$ the flatness will have improved to $\delta = (R'/R)^\gamma \delta_0$. So, if we want to continue applying Theorem 2.1 to u in $B_{R'}$, we must check that $1/R' < (R'/R)^{2\gamma} \delta_0^2$ (since $\varepsilon = 1$) and hence, we will always reach a critical mesoscale $R' = CR^{\frac{2\gamma}{1+2\gamma}}$ for which we cannot continue iterating. To solve this, we need an additional “sliding method” step in the spirit of Berestycki, Caffarelli, Nirenberg [3]. This last step follows the ideas of [17] and is done in Section 7.

¹ By a small modification of the proof of Lemma 6.3.

Remark 2.3. We assume $\beta'(0) > 0$ for simplicity, although this assumption is not really necessary. Indeed, our same proofs gives almost identical results if the assumption $\beta'(0) > 0$ is relaxed to $\liminf_{t \downarrow 0} \beta(t)t^{-p} > 0$, for some $p > 1$.

More precisely, Theorem 2.1 can be proved under this more general condition, up to assuming $\varepsilon/R < \delta^q$ (instead of $\varepsilon/R < \delta^2$), for some suitable $q = q(p) > 2$. The reason for this change is the following: while $\beta'(0) > 0$ implies the exponential decay increasing 1D solutions at $-\infty$, $\beta(t) \geq t^p$ gives a slower power-like decay. Accordingly, the properties of 1D solutions like (3.2) and (3.4) change to similar ones where powers replace logarithms. Up to this changes, all of our statements and proofs are still valid—with minor modifications—in this more general framework. The most important modifications are localized in Section 3 and only propagate to rest of the paper through Lemma 6.2, where the size of the error is not $\sqrt{\varepsilon/R}$ but $(\varepsilon/R)^{1/q}$ (for some $q > 2$). This is the reason why we need to assume $\varepsilon/R < \delta^q$ instead of $\varepsilon/R < \delta^2$ in Theorem 2.1. By the rest, all the proofs remain essentially the same.

3. ODEs analysis and barriers

In this section we consider the family of second order ODEs

$$\ddot{u}_\varepsilon = \frac{1}{2}\Phi'_\varepsilon(u_\varepsilon) \quad \text{in } \mathbb{R}, \quad (3.1)$$

and we provide a classification of its solutions, for every $\varepsilon \in (0, 1)$ fixed. With respect to [18, Section 2.3], our ODEs analysis shows finer properties of global solutions such as (3.2), (3.3) and (3.4), which will be needed later in the proofs of our main theorems.

Lemma 3.1. (1D global solutions) Fix $\varepsilon \in (0, 1)$ and let Φ be as in (2.1). Then:

(i) Equation (3.1) has a unique solution v_ε with

$$v_\varepsilon(0) = \vartheta_1\varepsilon, \quad \lim_{x \rightarrow +\infty} v_\varepsilon(x) = 1,$$

which is implicitly given by

$$\int_{\vartheta_1\varepsilon}^{v_\varepsilon(x)} \frac{dw}{\sqrt{\Phi_\varepsilon(w)}} = x.$$

This solution v_ε is smooth, positive, increasing, convex, and satisfies $v_\varepsilon(x) \rightarrow 0$ as $x \rightarrow -\infty$.

(ii) For every $t > 0$, equation (3.1) has a unique solution v_ε^t with

$$v_\varepsilon^t(0) = \vartheta_1\varepsilon, \quad \lim_{x \rightarrow +\infty} v_\varepsilon^t(x) = 1 + t.$$

Moreover, v_ε^t is of class C^2 , increasing, convex, and satisfies $v_\varepsilon^t(x) \rightarrow -\infty$, $\dot{v}_\varepsilon^t(x) \rightarrow \sqrt{2t+t^2}$ as $x \rightarrow -\infty$. Also, if x_ε^t denotes the unique root of v_ε^t —i.e. the point where $v_\varepsilon^t(x_\varepsilon^t) = 0$ —, then

$$x_\varepsilon^t \geq -\varepsilon\sqrt{2c_1} \log\left(1 + \frac{\vartheta_1}{t}\right), \quad (3.2)$$

where $c_1 > 0$ is the constant in (2.1).

(iii) For any $\tau \in (-1, 0)$, equation (3.1) has a unique solution v_ε^τ with

$$v_\varepsilon^\tau(0) = \vartheta_1\varepsilon, \quad \lim_{x \rightarrow +\infty} \dot{v}_\varepsilon^\tau(x) = 1 - |\tau|.$$

Moreover, v_ε^τ is smooth, positive, and satisfies $v_\varepsilon^\tau(x) \rightarrow +\infty$, $\dot{v}_\varepsilon^\tau(x) \rightarrow -1 + |\tau|$ as $x \rightarrow -\infty$. Also, v_ε^τ has a unique point of minimum y_ε^τ satisfying

$$\sqrt{\frac{|\tau|}{c_1}}\varepsilon \leq v_\varepsilon^\tau(y_\varepsilon^\tau) \leq \sqrt{2c_1|\tau|}\varepsilon, \quad (3.3)$$

and

$$y_\varepsilon^\tau \geq -\varepsilon\sqrt{2c_1} \left(2 + \log \frac{\vartheta_1}{\sqrt{2|\tau|/c_1}}\right), \quad (3.4)$$

where $c_1 > 0$ is the constant in (2.1).

Proof. After scaling, let us assume $\varepsilon = 1$ and set $u = u_\varepsilon$, $v = v_\varepsilon$, $v^t = v_\varepsilon^t$ and $v^\tau = v_\varepsilon^\tau$.

Since Φ' is bounded, nonnegative and continuous, a local C^2 solution $u = u(x)$ to (3.1) with $(u(0), \dot{u}(0)) = (\vartheta_1, \dot{u}_0)$ exists and it is convex on its maximal interval of definition I . Using the assumptions on Φ' , it is not difficult to see that $I = \mathbb{R}$. Further, since (3.1) is invariant under even reflections ($x \rightarrow -x$), we assume $\dot{u}_0 > 0$.

Step 1. Since \dot{u} is nondecreasing the limits $\lim_{x \rightarrow \pm\infty} \dot{u}$ exist. Since $\dot{u}_0 > 0$ we see that $u(x) \rightarrow +\infty$ as $x \rightarrow +\infty$. Let us define

$$\lim_{x \rightarrow +\infty} \dot{u}(x) =: A \in (0, +\infty).$$

Hence, using that the Hamiltonian $x \rightarrow \dot{u}(x)^2 - \Phi(u(x))$ must be constant (and $\Phi(u) = 1$ for $u > 0$ large enough) we obtain

$$\dot{u}(x)^2 - \Phi(u(x)) \equiv A^2 - 1, \quad x \in \mathbb{R}. \quad (3.5)$$

Step 2. Let us classify first monotone solutions: assume $\lim_{x \rightarrow -\infty} \dot{u} \geq 0$ and hence $\dot{u} > 0$ in \mathbb{R} . In this case (since $\Phi = \Phi'(u) = 0$ for $u < 0$) we obtain that either

$$\lim_{x \rightarrow -\infty} u(x) = 0 \quad \text{and} \quad \lim_{x \rightarrow -\infty} \dot{u}(x) = 0$$

or

$$\lim_{x \rightarrow -\infty} u(x) = -\infty \quad \text{and} \quad \lim_{x \rightarrow -\infty} \dot{u}(x) =: B \in (0, A).$$

From (3.5), we obtain that in the first case $A = 1$, while in the second one we have

$$A^2 - B^2 = 1,$$

and hence $A > 1$.

Now in the first case integrating (3.5) —with $A = 1$ — we get

$$\int_{v(y)}^{v(x)} \frac{dw}{\sqrt{\Phi(w)}} = x - y, \quad (3.6)$$

for every $y \leq x$ and so (i) follows. The solution in (ii), is obtained in the case $A = 1 + t$, so $B^2 = A^2 - 1 = 2t + t^2$. To complete (ii) we are left to show (3.2). Integrating (3.5) between $x^t \leq 0$ (the root of v^t) and 0 (recall $v^t(0) = \vartheta_1$) and using (2.1) we obtain

$$\begin{aligned} 0 - x^t &= \int_0^{\vartheta_1} \frac{dw}{\sqrt{\Phi(w) + 2t + t^2}} \leq \int_0^{\vartheta_1} \frac{dw}{\sqrt{\frac{1}{2c_1}w^2 + 2t + t^2}} \\ &\leq \sqrt{2c_1} \int_0^{\vartheta_1} \frac{dw}{w + t} = \sqrt{2c_1} \log \left(1 + \frac{\vartheta_1}{t} \right). \end{aligned}$$

Step 3. Let us consider now the case where \dot{u} changes sign. If so, there is $x_0 \in \mathbb{R}$ such that $\dot{u}(x) \leq 0$ for $x \leq x_0$ and $\dot{u}(x) \geq 0$ for $x \geq x_0$ (by convexity of u). Since the equation is invariant under the reflection $x \mapsto 2x_0 - x$, it follows that $u(x) = u(2x_0 - x)$ and thus $\lim_{x \rightarrow -\infty} \dot{u} = -A$. Note that the solutions $u = v^\tau$ described in (iii) corresponds to the setting $A = 1 - |\tau|$, with $\tau \in (-1, 0)$.

To show (3.3), we notice that if y^τ is the minimum point of v^τ , then $\dot{v}^\tau(y^\tau) = 0$. Thus, by (3.5), it follows

$$\Phi(v^\tau(y^\tau)) = 2|\tau| - \tau^2. \quad (3.7)$$

Using again (2.1) —note that $v^\tau(y^\tau) < v^\tau(0) = \vartheta_1$ — we obtain

$$\frac{|\tau|}{c_1} \leq \frac{2|\tau| - \tau^2}{c_1} \leq \frac{1}{2}(v^\tau(y^\tau))^2 \leq c_1(2|\tau| - \tau^2) \leq 2c_1|\tau|$$

and (3.3) follows.

We are left to prove (3.4). We use now (2.1) to obtain that, for all $w \in (v^\tau(y^\tau), \vartheta_1)$,

$$\begin{aligned} \Phi(w) - 2|\tau| + \tau^2 &= \Phi(w) - \Phi(v^\tau(y^\tau)) = \int_{v^\tau(y^\tau)}^w \Phi'(t) dt \geq \frac{1}{c_1} [t^2]_{v^\tau(y^\tau)}^w \\ &= \frac{1}{c_1} (w^2 - (v^\tau(y^\tau))^2) \geq \frac{w}{2c_1} (w - v^\tau(y^\tau)). \end{aligned} \quad (3.8)$$

Hence, integrating (3.5) between y^τ and 0 (recall $v^\tau(0) = \vartheta_1$) we obtain

$$\begin{aligned} 0 - y^\tau &= \int_{v^\tau(y^\tau)}^{\vartheta_1} \frac{dw}{\sqrt{\Phi(w) - 2|\tau| + \tau^2}} \leq \sqrt{2c_1} \int_{v^\tau(y^\tau)}^{\vartheta_1} \frac{dw}{\sqrt{w} \sqrt{w - v^\tau(y^\tau)}} \\ &= \sqrt{2c_1} \int_1^{\vartheta_1/v^\tau(y^\tau)} \frac{d\omega}{\sqrt{\omega} \sqrt{\omega - 1}} \\ &\leq \sqrt{2c_1} \left(\int_1^2 \frac{d\omega}{\sqrt{\omega} \sqrt{\omega - 1}} + \int_2^{\vartheta_1/v^\tau(y^\tau)} \frac{d\omega}{\omega - 1} \right) \\ &= \sqrt{2c_1} \left(\log(3 + 2\sqrt{2}) + \log \left(\frac{\vartheta_1}{v^\tau(y^\tau)} \right) \right) \leq \sqrt{2c_1} \left(2 + \log \frac{\vartheta_1}{\sqrt{2|\tau|/c_1}} \right). \quad \square \end{aligned}$$

In the following remark we introduce important one-dimensional super- and sub-solutions which will be used in the sequel.

Remark 3.2. Lemma 3.1 gives a classification of solutions to (3.1) in one dimension. The properties of such solutions are determined by their slopes at infinity, 1 , $1+t$, or $1-|\tau|$, where $t > 0$ and $\tau \in (-1, 0)$ are parameters. As done in Lemma 3.1 it is convenient to “center” these solutions so that their value at $x = 0$ is $\vartheta_1\varepsilon$.

In what follows, we will always take

$$t = \varepsilon, \quad \tau = -\varepsilon.$$

Within this setting, we define

$$w_\varepsilon^\varepsilon(x) := \begin{cases} 0 & \text{if } x \leq x_\varepsilon^\varepsilon \\ v_\varepsilon^\varepsilon(x) & \text{if } x > x_\varepsilon^\varepsilon, \end{cases} \quad w_\varepsilon^{-\varepsilon}(x) := \begin{cases} v_\varepsilon^{-\varepsilon}(x_\varepsilon^{-\varepsilon}) & \text{if } x \leq y_\varepsilon^{-\varepsilon} \\ v_\varepsilon^{-\varepsilon}(x) & \text{if } x > y_\varepsilon^{-\varepsilon}, \end{cases}$$

where $x_\varepsilon^\varepsilon$ and $y_\varepsilon^{-\varepsilon}$ are, respectively, the (unique) root of $v_\varepsilon^\varepsilon$ and the point of minimum of $v_\varepsilon^{-\varepsilon}$.

It is immediate to see that $w_\varepsilon^\varepsilon$ and $w_\varepsilon^{-\varepsilon}$ are, respectively, a sub- and a super- solution of (3.1), both in the viscosity sense or in the weak sense.

The next two lemmata are auxiliary results, which will be crucial in the proofs of our main theorems (see Section 6). In the statement of the next lemma we use the following standard notation $\text{diam}(X) := \sup X - \inf X$ for subsets $X \subset \mathbb{R}$.

Lemma 3.3. *There exists $c > 1$ depending only on ϑ_1, ϑ_2 and $c_1 > 0$ as in (2.1) such that*

$$\text{diam}(\{\vartheta_1\varepsilon \leq w_\varepsilon^\varepsilon \leq \vartheta_2\varepsilon\}) \leq c\varepsilon, \quad \forall \varepsilon > 0,$$

and

$$\text{diam}(\{\vartheta_1\varepsilon \leq w_\varepsilon^{-\varepsilon} \leq \vartheta_2\varepsilon\}) \leq c\varepsilon, \quad \forall \varepsilon \in \left(0, \frac{\vartheta_1^2}{8c_1}\right).$$

Proof. By scaling, we need to prove that $w^\varepsilon := w_1^\varepsilon$ and $w^{-\varepsilon} := w_1^{-\varepsilon}$ satisfy

- (i) $\text{diam}(\{\vartheta_1 \leq w^\varepsilon \leq \vartheta_2\}) \leq c$;
- (ii) $\text{diam}(\{\vartheta_1 \leq w^{-\varepsilon} \leq \vartheta_2\}) \leq c$.

To prove (i) we notice that (3.5) reads as $(\dot{w}^\varepsilon)^2 = \Phi(w^\varepsilon) + 2\varepsilon + \varepsilon^2$ in $\{w^\varepsilon > 0\}$ and so, by (2.1), we find

$$\frac{\vartheta_1^2}{c_1} \leq \Phi(\vartheta_1) \leq (\dot{w}^\varepsilon)^2 \quad \text{in } \{\vartheta_1 \leq w^\varepsilon \leq \vartheta_2\}.$$

Integrating between y and x , it follows

$$w^\varepsilon(x) - w^\varepsilon(y) \geq \frac{\vartheta_1}{\sqrt{c_1}}(x - y).$$

So, choosing x such that $w^\varepsilon(x) = \vartheta_2$, $y = 0$ and recalling that $w^\varepsilon(0) = \vartheta_1$, we find $\frac{\vartheta_1}{\sqrt{c_1}}x \leq \vartheta_2 - w^\varepsilon(0) = \vartheta_2 - \vartheta_1$, and (i) is proved.

To prove (ii) we use again (3.5): $(\dot{w}^{-\varepsilon})^2 - \Phi(w^{-\varepsilon}) = -2\varepsilon + \varepsilon^2$. Hence, for $\varepsilon \in \left(0, \frac{\vartheta_1^2}{8c_1}\right)$, we find

$$(\dot{w}^{-\varepsilon})^2 \geq \Phi(w^{-\varepsilon}) - 2\varepsilon \geq \frac{\vartheta_1^2}{2c_1} - 2\varepsilon \geq \frac{\vartheta_1^2}{4c_1} > 0, \quad \text{in } \{\vartheta_1 \leq w^{-\varepsilon} \leq \vartheta_2\},$$

which allows us to conclude similarly as for (i). \square

Lemma 3.4. *For every $\sigma \in (0, 1)$, there exists $\varepsilon_0 \in (0, 1)$ depending only on $\vartheta_1, \vartheta_2, c_1 > 0$ in (2.1) and σ , such that for every $\delta \in [0, 1)$ and every $\varepsilon \in (0, \varepsilon_0)$, if $w_\varepsilon^\varepsilon$ and $w_\varepsilon^{-\varepsilon}$ are as in Remark 3.2, then:*

(i) *If $x_\varepsilon^\varepsilon$ is such that $v_\varepsilon^\varepsilon(x_\varepsilon^\varepsilon) = 0$, then*

$$\begin{aligned} w_\varepsilon^\varepsilon(x - \delta - \varepsilon^\sigma) + \delta + \frac{1}{2}\varepsilon^\sigma &\leq x, & x &\in (x_\varepsilon^\varepsilon + \delta + \varepsilon^\sigma, 1) \\ w_\varepsilon^\varepsilon(x + \delta + \varepsilon^\sigma) - \delta - \frac{1}{2}\varepsilon^\sigma &\geq x, & x &\in (-1, 1). \end{aligned} \tag{3.9}$$

(ii) If $y_\varepsilon^{-\varepsilon}$ is the minimum point of $v_\varepsilon^{-\varepsilon}$, then

$$\begin{aligned} w_\varepsilon^{-\varepsilon}(x - \delta - \varepsilon^\sigma) + \delta + \frac{1}{2}\varepsilon^\sigma &\leq x, & x \in (y_\varepsilon^{-\varepsilon} + \delta + \varepsilon^\sigma, 1) \\ w_\varepsilon^{-\varepsilon}(x + \delta + \varepsilon^\sigma) - \delta - \frac{1}{2}\varepsilon^\sigma &\geq x, & x \in (-1, 1). \end{aligned} \quad (3.10)$$

Proof. Let us prove part (i). To simplify the notations, we set $w^\varepsilon := w_\varepsilon^\varepsilon$ and $x_\varepsilon = x_\varepsilon^\varepsilon$. Let $\tilde{x}_\varepsilon > 0 > x_\varepsilon$ such that $w^\varepsilon(\tilde{x}_\varepsilon) = \vartheta_2\varepsilon$, hence w^ε is linear for $x \geq \tilde{x}_\varepsilon$. Then if $x \in (\tilde{x}_\varepsilon + \delta + \varepsilon^\sigma, 1)$, we have

$$\begin{aligned} w^\varepsilon(x - \delta - \varepsilon^\sigma) - x &= \vartheta_2\varepsilon + (1 + \varepsilon)(x - \delta - \varepsilon^\sigma - \tilde{x}_\varepsilon) - x \\ &= (\vartheta_2 + x)\varepsilon - (1 + \varepsilon)(\delta + \varepsilon^\sigma) - (1 + \varepsilon)\tilde{x}_\varepsilon \\ &\leq (\vartheta_2 + 1)\varepsilon - \varepsilon^\sigma - \delta \leq -\delta - \frac{1}{2}\varepsilon^\sigma, \end{aligned}$$

for every $\varepsilon \leq \varepsilon_0 \leq [2(\vartheta_2 + 1)]^{\frac{1}{\sigma-1}}$, while if $x \in (x_\varepsilon + \delta + \varepsilon^\sigma, \tilde{x}_\varepsilon + \delta + \varepsilon^\sigma)$, we obtain by (3.2) (with $t = \varepsilon$)

$$\begin{aligned} w^\varepsilon(x - \delta - \varepsilon^\sigma) - x &\leq \vartheta_2\varepsilon - (x_\varepsilon + \delta + \varepsilon^\sigma) \leq \vartheta_2\varepsilon + C\varepsilon|\log \varepsilon| - \delta - \varepsilon^\sigma \\ &\leq -\delta - \frac{1}{2}\varepsilon^\sigma, \end{aligned}$$

taking eventually ε_0 smaller. Notice that the constant $C > 0$ depends only on ϑ_1 , and c_1 (cf. (3.2)).

To show the second inequality in (3.9), we assume first $x + \delta + \varepsilon^\sigma \geq \tilde{x}_\varepsilon$ and we notice that, since $\tilde{x}_\varepsilon \in (0, c\varepsilon)$ (where $c > 0$ is as in Lemma 3.3), we have

$$\begin{aligned} w^\varepsilon(x + \delta + \varepsilon^\sigma) - x &= \vartheta_2\varepsilon + (1 + \varepsilon)(x + \delta + \varepsilon^\sigma - \tilde{x}_\varepsilon) - x \\ &\geq \delta + \varepsilon^\sigma - \tilde{x}_\varepsilon + \varepsilon(x + \delta + \varepsilon^\sigma - \tilde{x}_\varepsilon) \geq \delta + \varepsilon^\sigma - c\varepsilon \geq \delta + \frac{1}{2}\varepsilon^\sigma, \end{aligned}$$

provided that ε_0 is small enough. Further, since $\tilde{x}_\varepsilon \leq c\varepsilon$, when $x \leq \tilde{x}_\varepsilon - \delta - \varepsilon^\sigma$ we have $x \leq 0$, and the second inequality in (3.9) follows.

To show (ii), we set $w^{-\varepsilon} = w_\varepsilon^{-\varepsilon}$, $y_\varepsilon = y_\varepsilon^{-\varepsilon}$, and we take \tilde{y}_ε such that $w^{-\varepsilon}(\tilde{y}_\varepsilon) = \vartheta_2\varepsilon$. The proof of the first inequality works exactly as before, using (3.4) instead of (3.2). To show the second, we assume first $x \in (\tilde{y}_\varepsilon - \delta - \varepsilon^\sigma, 1)$ and, recalling that $\tilde{y}_\varepsilon \leq c\varepsilon$, we write

$$\begin{aligned} w^{-\varepsilon}(x + \delta + \varepsilon^\sigma) - x &= \vartheta_2\varepsilon + (1 - \varepsilon)(x + \delta + \varepsilon^\sigma - \tilde{y}_\varepsilon) - x \\ &= (\vartheta_2 - x)\varepsilon + (1 - \varepsilon)(\delta + \varepsilon^\sigma) - (1 - \varepsilon)\tilde{y}_\varepsilon \\ &\geq \delta + (1 - \varepsilon)\varepsilon^\sigma - (1 + c)\varepsilon - \varepsilon\delta \geq \delta + \frac{1}{2}\varepsilon^\sigma, \end{aligned}$$

taking eventually ε_0 smaller. As above, if $x \leq \tilde{y}_\varepsilon - \delta - \varepsilon^\sigma$, then x is negative and the inequality is automatically satisfied. \square

We end this section by proving that solutions u_ε to (1.8) decay exponentially fast inside $\{u_\varepsilon \leq \vartheta_1 \varepsilon\}$ as $\varepsilon \rightarrow 0$. This is a main fact we will use later in Section 6 (see for instance Lemma 6.2). This decay is obtained in Lemma 3.6 using a sliding type argument based on the continuous family of super-solutions constructed in the following lemma.

Lemma 3.5. Fix $c_1 > 0$ as in (2.1) and $c^2 := \frac{1}{c_1}$. For every $\varepsilon \in (0, 1)$, $\varrho > 0$ and $R \geq \varrho$, let

$$\varphi(r) = \varphi_{\varepsilon, \varrho, R}(r) := e^{-\frac{\mu_+}{\varepsilon}(R-r)} \frac{1 - \frac{\mu_+}{\mu_-} e^{-\frac{\mu_+ - \mu_-}{\varepsilon}(r-\varrho)}}{1 - \frac{\mu_+}{\mu_-} e^{-\frac{\mu_+ - \mu_-}{\varepsilon}(R-\varrho)}}, \quad r \in [\varrho, R], \quad (3.11)$$

where μ_\pm are defined by

$$\mu_\pm = -\frac{N-1}{2\varrho} \varepsilon \pm \sqrt{\left(\frac{N-1}{2\varrho}\right)^2 \varepsilon^2 + c^2}. \quad (3.12)$$

Then, for every $x_0 \in \mathbb{R}^N$ and $\varrho > 0$, the function

$$\psi(x) := \psi_{\varepsilon, \varrho, R, x_0}(x) := \begin{cases} \varphi(\varrho) & \text{in } B_\varrho(x_0) \\ \varphi(|x - x_0|) & \text{in } B_R(x_0) \setminus B_\varrho(x_0) \end{cases} \quad (3.13)$$

satisfies

$$\begin{cases} -\Delta \psi + \frac{1}{c_1 \varepsilon^2} \psi \geq 0 & \text{in } B_R(x_0) \\ \psi = 1 & \text{in } \partial B_R(x_0) \\ \partial_r \psi \geq 0 & \text{in } B_R(x_0), \end{cases} \quad (3.14)$$

in the weak sense.

Proof. Up to translations and scaling, we may assume $x_0 = 0$, $\varepsilon = 1$ and set $\varphi = \varphi_1$, $\psi = \psi_1$. Notice that if $\varrho = R$, we have $\psi \equiv 1$ in B_R (i.e. $\varphi = 1$ in $(0, R)$) and (3.14) is trivial.

If $0 < \varrho < R$, since $\varphi(\varrho) > 0$ and $\varphi(R) = 1$, it suffices to verify that the differential inequality in (3.14) is satisfied in $B_R \setminus B_\varrho$ with $\varphi'(\varrho) = 0$ and $\varphi' \geq 0$ in (ϱ, R) .

To see this, we notice that if $r \in (\varrho, R)$ and $\varphi' \geq 0$, then

$$-\Delta \varphi + c^2 \varphi = -\varphi'' - \frac{N-1}{r} \varphi' + c^2 \varphi \geq -\varphi'' - \frac{N-1}{\varrho} \varphi' + c^2 \varphi,$$

and so, it is enough to check that

$$\begin{cases} -\varphi'' - \frac{N-1}{\varrho} \varphi' + c^2 \varphi = 0 & \text{in } (\varrho, R) \\ \varphi' \geq 0 & \text{in } (\varrho, R) \\ \varphi'(\varrho) = 0. \end{cases}$$

Integrating the equation above, we easily see that

$$\varphi(r) = Ae^{\mu_+ r} + Be^{\mu_- r} \quad r \in (R/2, R),$$

for some suitable constants $A, B \in \mathbb{R}$, and μ_{\pm} as in (3.12). Imposing that $\varphi'(\varrho) = 0$ and $\varphi(R) = 1$, we deduce

$$A = \frac{1}{e^{\mu_+ R} \left(1 - \frac{\mu_+}{\mu_-} e^{-(\mu_+ - \mu_-)(R - \varrho)}\right)}, \quad B = -\frac{\mu_+}{\mu_-} e^{(\mu_+ - \mu_-)\varrho} A$$

and, substituting into the expression of φ , (3.11) follows. Checking that $\varphi' \geq 0$ in (ϱ, R) is a straightforward computation. \square

Lemma 3.6. *There exists $\varepsilon_0 \in (0, 1)$ depending only on N and c_1 such that for every $\varepsilon \in (0, \varepsilon_0)$, every solution u_{ε} to (1.8), every $x_0 \in \{u_{\varepsilon} \leq \vartheta_1 \varepsilon\}$ and every ball $B_{\varepsilon^{3/4}}(x_0) \subset \{u \leq \vartheta_1 \varepsilon\}$, then*

$$u_{\varepsilon} \leq 3\vartheta_1 \varepsilon e^{-\frac{\varepsilon^{-1/4}}{4c_1^{1/2}}} \quad \text{in } B_{\varepsilon^{3/4}}(x_0). \quad (3.15)$$

Proof. Fix $R > 0$ and $x_0 \in \{u \leq \vartheta_1 \varepsilon\}$ such that $B_R(x_0) \subset \{u \leq \vartheta_1 \varepsilon\}$. Let $\psi_{\varrho} := \psi_{\varepsilon, \varrho, R, x_0}$ be defined as in (3.13), satisfying (3.14), and let $\tilde{\psi}_{\varrho} := \vartheta_1 \varepsilon \psi_{\varrho}$.

If $\varrho = R$, then $\tilde{\psi}_R = \vartheta_1 \varepsilon$ satisfies (3.14), with $\tilde{\psi}_R \geq u_{\varepsilon}$ in $B_R(x_0)$. Setting $v := \tilde{\psi}_R - u_{\varepsilon}$ and recalling that $B_R(x_0) \subset \{u \leq \vartheta_1 \varepsilon\}$, we obtain

$$-\Delta v + \frac{1}{c_1 \varepsilon^2} v = -\Delta \tilde{\psi}_R + \frac{1}{c_1 \varepsilon^2} \tilde{\psi}_R + \Delta u - \frac{1}{c_1 \varepsilon^2} u \geq \Delta u - \frac{1}{2} \Phi'_{\varepsilon}(u) = 0,$$

and thus

$$\begin{cases} -\Delta v + \frac{1}{c_1 \varepsilon^2} v \geq 0 & \text{in } B_R(x_0) \\ v \geq 0 & \text{in } B_R(x_0). \end{cases}$$

By the strong maximum principle, $v > 0$ in $B_R(x_0)$ (it cannot be $v = 0$ since ψ_R is a strict super-solution), that is $\psi_R > u_{\varepsilon}$ in $B_R(x_0)$. Now, let

$$\varrho_* := \inf\{\varrho \in (0, R] : \psi_{\varrho} > u_{\varepsilon} \text{ in } B_R(x_0)\}.$$

We have $\varrho_* = 0$. If by contradiction, $\varrho_* > 0$, we may repeat the above argument setting $v := \tilde{\psi}_{\varrho_*} - u_{\varepsilon}$ and noticing that $v \geq 0$ in $B_R(x_0)$ with $v(x_*) = 0$, for some $x_* \in \overline{B}_R(x_0)$. Since by construction $B_R(x_0) \subset \subset \{u \leq \vartheta_1 \varepsilon\}$, $\tilde{\psi}_{\varrho_*} = \vartheta_1 \varepsilon$ on $\partial B_R(x_0)$, and ψ_{ϱ_*} is radially increasing near the boundary of the ball, it must be $x_* \in B_R(x_0)$. Thus using the linear equation for v and the strong maximum principle either $v \equiv 0$ or $v > 0$ in $B_R(x_0)$. Since both scenarios are impossible, our contradiction follows.

In particular, we have $\varrho_* < \frac{R}{2}$ and so, $u_\varepsilon \leq \psi_{R/2}$ in $B_R(x_0)$. Now, choosing $R = \varepsilon^{3/4}$, taking $\varrho = \frac{R}{2}$ in (3.11) and using (3.13), we obtain

$$u_\varepsilon \leq \varepsilon \vartheta_1 \varphi_{\varepsilon^{3/4}/2}(\varepsilon^{3/4}/2) \leq \varepsilon \vartheta_1 \left(1 - \frac{\mu_+}{\mu_-}\right) e^{-\frac{\mu_+}{2\sqrt[4]{\varepsilon}}} \quad \text{in } B_{\varepsilon^{3/4}/2}(x_0),$$

where μ_\pm are defined in (3.12) (with $R = \varepsilon^{3/4}$). Since $\mu_\pm \rightarrow \pm \frac{1}{\sqrt{c_1}}$ as $\varepsilon \rightarrow 0^+$, there is $\varepsilon_0 \in (0, 1)$ (depending only on N and c_1) such that $\mu_+ \geq 1/(2\sqrt{c_1})$ and $-\mu_+/\mu_- \leq 2$ for every $\varepsilon \in (0, \varepsilon_0)$ and thus (3.15) follows. \square

4. Lipschitz and non-degeneracy estimates

We recall now a useful Lipschitz estimate from [11].

Proposition 4.1 (Uniform Lipschitz estimate; see [11, Theorem 1.2]). *For any $V \subset\subset B_1$, there exists $\overline{C} > 0$ depending only on N , L , ϑ_2 and V such that for every $\varepsilon \in (0, 1)$ and for every critical point u_ε of (1.1) in B_1 with $u_\varepsilon(0) \leq \vartheta_2 \varepsilon$ we have*

$$\sup_V |\nabla u_\varepsilon| \leq \overline{C}. \quad (4.1)$$

We also need a non-degeneracy estimate related to [11, Theorem 1.8]. Our estimate is stronger since balls $B_r(z)$ do not need to be centered at some point in $\{u \geq C\varepsilon\}$, with C large, and can be centered at any point in $\{u_\varepsilon \geq \vartheta_1 \varepsilon\}$

Lemma 4.2 (Uniform non-degeneracy). *There exists $\varepsilon_0 \in (0, 1)$ depending only on ϑ_1 and c_1 such that for every $\kappa > 0$, there exists $c_\kappa > 0$ depending only on N , L , ϑ_2 and κ such that for every $\varepsilon \in (0, \varepsilon_0)$, every local minimizer u_ε of (1.1) in B_1 , every $z \in \{u_\varepsilon \geq \vartheta_1 \varepsilon\}$ and every $r \geq \kappa \varepsilon$ such that $B_r(z) \subset\subset B_1$, then*

$$\sup_{B_r(z)} u_\varepsilon \geq c_\kappa r. \quad (4.2)$$

Proof. Let us fix $\kappa > 0$, and assume that $\varepsilon \in (0, \varepsilon_0)$, $u = u_\varepsilon$, $z \in \{u_\varepsilon \geq \vartheta_1 \varepsilon\}$ and $r \geq \kappa \varepsilon$. Define

$$\omega(r) := \frac{1}{r} \sup_{B_r(z)} u. \quad (4.3)$$

Our goal is to prove a lower bound for ω , which holds if ε_0 is small enough. Up to translate and scaling, we may assume $r = 1$ and $z = 0$. Let $\sigma := \frac{1}{3}$ where $c > 0$ is the constant appearing in (4.7) depending only on N and c_1 .

Step 1: Estimates. Let $\varphi \in C_0^\infty(B_1)$, $0 \leq \varphi \leq 1$, with $\varphi = 1$ in $B_{7/8}$. Assume also

$$|\nabla \varphi| \leq c_N, \quad |\Delta \varphi| \leq c_N, \quad (4.4)$$

for some $c_N > 0$. Testing the equation of u with $\eta = u\varphi^2$, it is not difficult to find

$$\int_{B_1} [|\nabla u|^2 + \tfrac{1}{2}\Phi'_\varepsilon(u)u] \varphi^2 \, dx = \tfrac{1}{2} \int_{B_1} u^2 \Delta(\varphi^2) \, dx,$$

which, since $\Phi'_\varepsilon(u)u \geq 0$ implies

$$\int_{B_{7/8}} |\nabla u|^2 \, dx \leq c_N \int_{B_1} u^2 \, dx, \quad (4.5)$$

for some new $c_N > 0$.

Now, let $\phi \in C^\infty(\mathbb{R}^N)$, $0 \leq \phi \leq 1$ with $\phi = 0$ in $B_{3/4}$ and $\phi = 1$ in $\mathbb{R}^N \setminus B_{7/8}$, satisfying (4.4). Taking $v = \phi u$ as a competitor for u , we deduce

$$\begin{aligned} \int_{B_1} \Phi_\varepsilon(u) - \Phi_\varepsilon(\phi u) \, dx &\leq \int_{B_1} |\nabla(u\phi)|^2 - |\nabla u|^2 \, dx \\ &\leq \int_{B_1} (\phi^2 - 1) |\nabla u|^2 \, dx + 2 \int_{B_{7/8}} u^2 |\nabla \phi|^2 \, dx + \int_{B_{7/8}} |\nabla u|^2 \phi^2 \, dx \\ &\leq c_N \int_{B_{7/8}} |\nabla u|^2 + u^2 \, dx, \end{aligned}$$

for some new $c_N > 0$ and so, recalling that $\phi \leq 1$, $\Phi'_\varepsilon \geq 0$ and using (4.5), it follows

$$\int_{B_{3/4}} \Phi_\varepsilon(u) \, dx \leq \int_{B_1} \Phi_\varepsilon(u) - \Phi_\varepsilon(\phi u) \, dx \leq c_N \int_{B_1} u^2 \, dx.$$

In particular, by the definition of ω , we conclude

$$\int_{B_{3/4}} \Phi_\varepsilon(u) \, dx \leq c_N \omega(1)^2, \quad (4.6)$$

for some new $c_N > 0$.

Step 2: Decay of ω . Note that for all $y \in B_{1/2}$, since u is subharmonic, we have

$$u(y) \leq \fint_{B_{1/4}(y)} u \, dx \leq c_N \int_{B_{3/4}} u \, dx = c_N \left(\int_{B_{3/4} \cap \{u \geq t\}} u \, dx + \int_{B_{3/4} \cap \{u \leq t\}} u \, dx \right),$$

for every $t > 0$. Recalling that Φ is nondecreasing, there holds $\{u \geq t\} \subseteq \{\Phi_\varepsilon(u) \geq \Phi_\varepsilon(t)\}$ and, using that $\Phi_\varepsilon(t) \geq \frac{1}{2c_1}(t/\varepsilon)^2$ for $t \in (0, \vartheta_1\varepsilon]$ combined with (4.3), it follows

$$\begin{aligned} \int_{B_{3/4} \cap \{u \geq t\}} u \, dx &\leq \omega(1) \int_{B_{3/4} \cap \{u \geq t\}} dx \leq \omega(1) \int_{B_{3/4} \cap \{\Phi_\varepsilon(u) \geq \Phi_\varepsilon(t)\}} dx \\ &\leq c_1 \omega(1) \left(\frac{\varepsilon}{t}\right)^2 \int_{B_{3/4}} \Phi_\varepsilon(u) \, dx \leq c_1 c_N \left(\frac{\varepsilon}{t}\right)^2 \omega^3(1), \end{aligned}$$

where the last inequality is a direct application of (4.6). Substituting into the inequality above, we deduce

$$u(y) \leq c_N \left[c_1 c_N \left(\frac{\varepsilon}{t}\right)^2 \omega^3(1) + t \right] \leq c \left[\left(\frac{\varepsilon}{t}\right)^2 \omega^3(1) + t \right],$$

for some $c > 0$ depending only on N and c_1 , and so, by the arbitrariness of $y \in B_{1/2}$,

$$\omega\left(\frac{1}{2}\right) \leq c \left[\left(\frac{\varepsilon}{t}\right)^2 \omega^3(1) + t \right]. \quad (4.7)$$

Setting $t := \min\{\max\{\varepsilon, \omega(1)\}^{1+2\sigma}, \vartheta_1 \varepsilon\}$, we have that $t \leq \vartheta_1 \varepsilon$ thanks to the definition of ε_0 . So, using that $\sigma = \frac{1}{3}$, we may re-write (4.7) as

$$\omega\left(\frac{1}{2}\right) \leq c \max\{\varepsilon, \omega(1)\}^{1+2\sigma}. \quad (4.8)$$

Let us now assume by contradiction that we have $\varepsilon \leq \omega_0$ and $\omega(1) \leq \omega_0$, for $\omega_0 \in (0, 1/4)$ sufficiently small so that (4.8) implies

$$\omega\left(\frac{1}{2}\right) \leq \max\{\varepsilon, \omega(1)\}^{1+\sigma}.$$

After scaling (applying the above inequality to $u_\varepsilon(rx)/r$), we obtain provided $\varepsilon/r \in (0, \omega_0)$,

$$\omega\left(\frac{r}{2}\right) \leq \max\{\varepsilon/r, \omega(r)\}^{1+\sigma}.$$

Iterating the above inequality, we obtain that whenever $2^k \varepsilon \leq \omega_0$, we have either

$$(i) \quad \omega(2^{-k}) \leq (2^k \varepsilon)^{1+\sigma} \quad \text{or} \quad (ii) \quad \omega(2^{-k}) \leq \omega(1)^{(1+\sigma)^k},$$

for all $k \in \mathbb{N}$. Finally, choosing

$$k := \lceil \log_2(\varepsilon^{-1/2}) \rceil,$$

we have $2^{-k} \leq \varepsilon^{1/2} \leq 2^{-k+1}$ and hence $2^k \varepsilon \leq 2\varepsilon^{1/2} \in (0, \omega_0)$, provided $\varepsilon \in (0, \varepsilon_0)$ with $\varepsilon_0 > 0$ sufficiently small.

Hence, recalling $\omega(1) \leq \omega_0 \leq \frac{1}{4}$ and that by assumption $0 \in \{u_\varepsilon \geq \vartheta_1 \varepsilon\}$, we have

$$\max\{(2^k \varepsilon)^{1+\sigma}, (1/4)^{(1+\sigma)^k}\} \geq \omega(2^{-k}) := 2^k \sup_{B_{2^{-k}}} u \geq \vartheta_1 \varepsilon^{1/2}, \quad (4.9)$$

which clearly gives a contradiction if $\varepsilon \in (0, \varepsilon_0)$ with ε_0 chosen sufficiently small (since $(2^k \varepsilon)^{1+\sigma} \leq (2\varepsilon^{1/2})^{1+\sigma} \ll \varepsilon^{1/2}$ and $(1/4)^{(1+\sigma)^k} \ll 4^{-k} \ll \varepsilon^{1/2}$ as $\varepsilon \downarrow 0$). \square

5. Proof of Proposition 1.5

This section is devoted to the proof of Proposition 1.5. It will be obtained as a corollary of the following result, which is its equivalent version in terms of blow-down families.

Proposition 5.1. *Let Φ be as in (1.3)–(1.4) and let ϑ_1 and ϑ_2 as in (2.1). Let $u : \mathbb{R}^N \rightarrow \mathbb{R}_+$ be a minimizer of \mathcal{E} in \mathbb{R}^N not identically 0, with $0 \in \{\vartheta_1 \leq u \leq \vartheta_2\}$. Let $\{\varepsilon_j\}_{j \in \mathbb{N}}$ be a sequence satisfying $\varepsilon_j \rightarrow 0$ as $j \rightarrow +\infty$ and let u_{ε_j} be the corresponding blow-down family.*

Then for every $\alpha \in (0, 1)$, there exist sequences $\varepsilon_{j_\ell}, \delta_\ell \rightarrow 0$ and a 1-homogeneous entire local minimizer of (1.5) $u_0 \in W_{loc}^{1,\infty}(\mathbb{R}^N)$ — also not identically 0 — such that

$$|u_{\varepsilon_{j_\ell}} - u_0| \leq \delta_\ell \quad \text{in } B_1, \quad (5.1)$$

and

$$\begin{aligned} \{x : \text{dist}(x, \{u_0 > 0\}) \geq \delta_\ell\} &\subset \{u_{\varepsilon_{j_\ell}} \leq \vartheta_1 \varepsilon_{j_\ell}\} \subset \{u_{\varepsilon_{j_\ell}} \leq \vartheta_2 \varepsilon_{j_\ell}\} \\ &\subset \{x : \text{dist}(x, \{u_0 = 0\}) \leq \delta_\ell\} \quad \text{in } B_1, \end{aligned} \quad (5.2)$$

for every $\ell \in \mathbb{N}$.

The above statement will follow as a byproduct of several auxiliary results, having independent interest: in Lemma 5.2 we prove that families of minimizers of (1.1) converge (in a suitable sense, up to subsequences) to a minimizer of (1.5), while in Lemma 5.3 and Corollary 5.4 we deal with the convergence of the level sets of u_ε . Proposition 5.1 is a consequence of these facts and a Weiss type monotonicity formula (Lemma 5.5).

Lemma 5.2. *Let $R > 0$ and $\{u_{\varepsilon_j}\}$, $\varepsilon_j \downarrow 0$, be a sequence of minimizers of (1.1) in B_R , with $\varepsilon = \varepsilon_j$. Assume $u_{\varepsilon_j}(0) \leq \vartheta_2 \varepsilon_j$. Then, up to subsequence, we have*

$$u_{\varepsilon_j} \rightarrow u_0 \quad \text{in } H_{loc}^1(B_R) \cap C_{loc}^\alpha(B_R), \quad \text{for all } \alpha \in (0, 1), \quad (5.3)$$

as $j \rightarrow +\infty$, where $u_0 \in W_{loc}^{1,\infty}(B_R)$ is a minimizer of (1.5) in B_R .

Proof. By scaling we may assume $R = 1$. By Proposition 4.1, the family $\{u_\varepsilon\}_{\varepsilon \in (0,1)}$ is uniformly bounded in $W_{loc}^{1,\infty}(B_1)$. So, by the Ascoli-Arzelà theorem, for every $\alpha \in (0, 1)$, there exists $u_0 \in W_{loc}^{1,\infty}(B_1)$ and $\varepsilon_j \rightarrow 0$ as $j \rightarrow +\infty$ such that $u_{\varepsilon_j} \rightarrow u_0$ in $C_{loc}^\alpha(B_1)$. Furthermore, since in addition each u_ε is subharmonic and $\{u_\varepsilon\}_{\varepsilon \in (0,1)}$ is uniformly bounded in $L_{loc}^2(B_1)$, we deduce $u_{\varepsilon_j} \rightarrow u_0$ in $W_{loc}^{1,1}(B_1)$, up to subsequence (see for

instance [5, Lemma A.1]). Consequently, since $u_{\varepsilon_j}, u_0 \in W_{loc}^{1,\infty}(B_1)$ we deduce $u_{\varepsilon_j} \rightarrow u_0$ in $H_{loc}^1(B_1)$ by interpolation and (5.3) is proved.

Now, let us set for simplicity $u := u_0$ and $u_j := u_{\varepsilon_j}$. Let us fix $V \subset\subset B_1$ and show that

$$\mathcal{E}_0(u, V) \leq \liminf_{j \rightarrow +\infty} \mathcal{E}_{\varepsilon_j}(u_j, V). \quad (5.4)$$

Indeed, by H_{loc}^1 convergence, it is enough to check that

$$\int_V \chi_{\{u>0\}} dx \leq \liminf_{j \rightarrow +\infty} \int_V \Phi_{\varepsilon_j}(u_j) dx. \quad (5.5)$$

To show (5.5), we first notice that $\Phi_{\varepsilon_j}(u_j) \rightarrow 1$ in $\{u > 0\}$. Indeed, if $x \in \{u > 0\}$, that is $u(x) \geq \epsilon_x$ for some $\epsilon_x > 0$, then $u_j(x) \geq \epsilon_x/2 > 0$ for all j large enough. Now, by the monotonicity of Φ , $\Phi_{\varepsilon_j}(\epsilon_x/2) \leq \Phi_{\varepsilon_j}(u_j(x))$ for j large enough and thus, by definition of Φ_ε ,

$$1 = \lim_{j \rightarrow +\infty} \Phi_{\varepsilon_j}(\epsilon_x/2) \leq \limsup_{j \rightarrow +\infty} \Phi_{\varepsilon_j}(u_j(x)) \leq 1.$$

Consequently, by Fatou's lemma

$$\int_V \chi_{\{u>0\}} dx = \int_{V \cap \{u>0\}} dx \leq \liminf_{j \rightarrow +\infty} \int_{V \cap \{u>0\}} \Phi_{\varepsilon_j}(u_j) dx \leq \liminf_{j \rightarrow +\infty} \int_V \Phi_{\varepsilon_j}(u_j) dx,$$

and (5.5) follows.

Once (5.4) is established, let us fix $V := B_r$, $r < 1$, $\xi \in C_0^\infty(V)$, and $\varphi \in C^\infty(\overline{B_r})$ vanishing on ∂B_r with $\varphi > 0$ in B_r . Since u_j is a local minimizer, we have

$$\mathcal{E}_{\varepsilon_j}(u_j, V) \leq \mathcal{E}_{\varepsilon_j}(u_j + \xi - \delta\varphi, V), \quad (5.6)$$

for all $j \in \mathbb{N}$ and $\delta > 0$. Since $u_j \rightarrow u$ in $H^1(V)$, we immediately see that

$$\int_V |\nabla(u_j + \xi) - \delta\nabla\varphi|^2 dx \rightarrow \int_V |\nabla(u + \xi) - \delta\nabla\varphi|^2 dx \quad (5.7)$$

as $j \rightarrow +\infty$. Now, if $x \in \{u + \xi - \delta\varphi > 0\} \cap V$, there is $\epsilon_x > 0$ such that $u(x) + \xi(x) - \delta\varphi(x) \geq \epsilon_x$ and, since $u_j \rightarrow u$ locally uniformly, it must be $u_j(x) + \xi(x) - \delta\varphi(x) \geq \epsilon_x/2$ for every j large enough. Consequently, by monotonicity,

$$1 = \lim_{j \rightarrow +\infty} \Phi_{\varepsilon_j}(\epsilon_x/2) \leq \limsup_{j \rightarrow +\infty} \Phi_{\varepsilon_j}(u_j(x) + \xi(x) - \delta\varphi(x)) \leq 1.$$

Similar, whenever $x \in \{u + \xi - \delta\varphi < 0\} \cap V$, then $u(x) + \xi(x) - \delta\varphi(x) \leq -\epsilon_x$ for some $\epsilon_x > 0$ and so $u_j(x) + \xi(x) - \delta\varphi(x) \leq -\epsilon_x/2$ for every j large enough, which implies

$$0 \leq \Phi_{\varepsilon_j}(u_j(x) + \xi(x) - \delta\varphi(x)) \leq \Phi_{\varepsilon_j}(-\epsilon_x/2) = 0,$$

when j is large enough. On the other hand, for and every $m \in \mathbb{N}$, we have²

$$|\{u + \xi - \delta\varphi = 0\} \cap B_{r-1/m}| = 0 \quad \text{for all } \delta \in E_m \subset (0, 1), \text{ where } |(0, 1) \setminus E_m| = 0. \quad (5.8)$$

Consequently, since $|\cup_m ((0, 1) \setminus E_m)| = 0$,

$$|\{u + \xi - \delta\varphi = 0\} \cap B_r| = 0 \quad \text{for a.e. } \delta \in (0, 1), \quad (5.9)$$

and we deduce that for a.e. $\delta > 0$, $\Phi_{\varepsilon_j}(u_j + \xi - \delta\varphi) \rightarrow \chi_{\{u + \xi - \delta\varphi > 0\}}$ a.e. in B_r , as $j \rightarrow +\infty$.

So, putting together (5.4), (5.6), (5.7), noticing that $\{u + \xi - \delta\varphi > 0\} \subseteq \{u + \xi > 0\}$ and passing to the limit as $j \rightarrow +\infty$ by means of the dominated convergence theorem, we find

$$\begin{aligned} \mathcal{E}_0(u, V) &\leq \int_V |\nabla(u + \xi) - \delta\nabla\varphi|^2 + \chi_{\{u + \xi - \delta\varphi > 0\}} \, dx \\ &\leq \mathcal{E}_0(u + \xi, V) + 2\delta \|\nabla(u + \xi)\|_{L^2(V)} \|\nabla\varphi\|_{L^2(V)} + \delta^2 \|\nabla\varphi\|_{L^2(V)}^2, \end{aligned}$$

for a.e. $\delta > 0$. Finally, passing to the limit along a sequence $\delta = \delta_k \rightarrow 0$ for which (5.9) is satisfied for every $k \in \mathbb{N}$, we find $\mathcal{E}_0(u, V) \leq \mathcal{E}_0(u + \xi, V)$ and the thesis follows by the arbitrariness of $B_r \subset\subset B_1$ and $\xi \in C_0^\infty(B_r)$. \square

Lemma 5.3. *Let $R > 0$, $\{u_\varepsilon\}_{\varepsilon \in (0,1)}$ and u_0 as in Lemma 5.2. Then, for every $\vartheta \geq \vartheta_1$, there exists a sequence $\varepsilon_j \rightarrow 0$ such that*

$$\{u_{\varepsilon_j} \geq \vartheta_{\varepsilon_j}\} \rightarrow \overline{\{u_0 > 0\}} \quad \text{locally Hausdorff in } B_R, \quad (5.10)$$

as $j \rightarrow +\infty$.

Proof. By scaling, we may assume $R = 1$. Fix $\varrho \in (0, 1)$ and $\vartheta \geq \vartheta_1$. Set $u = u_0$, $u_j = u_{\varepsilon_j}$, $U_j := \{u_{\varepsilon_j} > \vartheta_{\varepsilon_j}\} \cap B_\varrho$, $\Omega := \{u > 0\} \cap B_\varrho$, and notice that by assumption $0 \in \Omega^c$. We first show that for every $z \in \overline{\Omega}$ and every $r > 0$ such that $B_r(z) \subset\subset B_1$, then

$$\sup_{B_r(z)} u \geq \frac{\varepsilon}{2}r, \quad (5.11)$$

² To see this, it is enough to apply the Coarea formula to the function $\frac{u+\xi}{\varphi}$, which is Lipschitz in $B_{r-1/m}$.

where $c > 0$ is the constant appearing in Lemma 4.2 for $\kappa = 1/2$. Given such $z \in \overline{\Omega}$ and $r > 0$, we take $y \in B_{r/2}(z)$ such that $u(y) > 0$. So, by uniform convergence, $y \in U$ for j large enough (and thus $u_j(y) > \vartheta_1 \varepsilon_j$). So by (4.2) (with $\kappa = 1$), there is $x_j \in B_{r/2}(y)$ such that $u_j(x_j) \geq \frac{c}{2}r$. Now, up to passing to a subsequence, $x_j \rightarrow x \in \overline{B_{r/2}(y)}$ as $j \rightarrow +\infty$ and thus, by C_{loc}^α convergence, $u(x) \geq \frac{c}{2}r$ and (5.11) follows since $x \in B_r(z)$.

Now fix $\sigma > 0$, and define

$$\Omega_\sigma := \{x : \text{dist}(x, \Omega) \leq \sigma\}, \quad U_{j,\sigma} := \{x : \text{dist}(x, U_j) \leq \sigma\}.$$

Let us show that $U_j \subset \Omega_\sigma$ for every $j \geq j_\sigma$, for some j_σ large enough. Indeed, assume by contradiction there is a sequence z_j such that $u_j(z_j) \geq \vartheta \varepsilon_j \geq \vartheta_1 \varepsilon_j$, but $z_j \notin \Omega_\sigma$. Then, by (4.2), there is j_σ such that

$$u_j(x_j) := \sup_{B_{\sigma/2}(z_j)} u_j \geq \frac{c}{2} \sigma,$$

for every $j \geq j_\sigma$ and some $x_j \in \overline{B_{\sigma/2}(z_j)}$. In addition, up to passing to a subsequence, $z_j \rightarrow z$, $x_j \rightarrow x \in \overline{B_{\sigma/2}(z)} \subset \subset \Omega^c$, and $u_j(x_j) \rightarrow u(x)$ as $j \rightarrow +\infty$, by C_{loc}^α convergence. Since $u(x) = 0$ by construction, we obtain a contradiction.

We also have $\overline{\Omega} \subset U_{j,\sigma}$ for every $j \geq j_\sigma$. Indeed, assume by contradiction there is $z_j \in \overline{\Omega}$ such that $z_j \notin U_{j,\sigma}$. Then, by (5.11), there is $x_j \in \overline{B_{\sigma/2}(z_j)}$ such that $u(x_j) \geq \frac{c}{4}\sigma$ while, by construction, $u_j < \vartheta \varepsilon_j$ in $\overline{B_{\sigma/2}(z_j)}$. So, since $z_j \rightarrow z$, $x_j \rightarrow x \in \overline{B_{\sigma/2}(z)}$ (up to a subsequence), we have $\frac{c}{4}\sigma \leq u(x) \leq 0$, a contradiction. The limit (5.10) follows from the arbitrariness of $\sigma > 0$. \square

Corollary 5.4. *Let $R > 0$, $\{u_\varepsilon\}_{\varepsilon \in (0,1)}$ and u_0 as in Lemma 5.2. Then, for every $\vartheta \geq \vartheta_1$, there exists a sequence $\varepsilon_j \rightarrow 0$ such that*

$$\{u_{\varepsilon_j} \leq \vartheta \varepsilon_j\} \rightarrow \{u_0 = 0\} \quad \text{locally Hausdorff in } B_R, \quad (5.12)$$

as $j \rightarrow +\infty$.

Proof. It is enough to apply Lemma 5.3 and noticing that $\{u_{\varepsilon_j} \leq \vartheta \varepsilon_j\} = \{u_{\varepsilon_j} \geq \vartheta \varepsilon_j\}^c$ and $\{u_0 = 0\} = \{u_0 > 0\}^c$. \square

Lemma 5.5. *Let u be a nonnegative entire local minimizer of (1.1) with $\varepsilon = 1$.*

Then, for every $x_0 \in \mathbb{R}^N$, the function

$$r \rightarrow \mathcal{W}(u, x_0, r) := r^{-N} \int_{B_r(x_0)} |\nabla u|^2 + \Phi(u) \, dx - r^{-1-N} \int_{\partial B_r(x_0)} u^2 \, d\sigma \quad (5.13)$$

is well-defined in $(0, \infty)$ and satisfies

$$\frac{d}{dr} \mathcal{W}(u, x_0, r) = 2r^{-N} \int_{\partial B_r(x_0)} \left(\partial_n u - \frac{u}{r} \right)^2 d\sigma + r^{-1-N} \int_{B_r(x_0)} u \Phi'(u) d\sigma, \quad (5.14)$$

where $\partial_n u := \nabla u \cdot n$ and n is the outward unit normal to $\partial B_r(x_0)$. In particular, the function $r \rightarrow \mathcal{W}(u, x_0, r)$ is non-decreasing.

Proof. We follow [26, Theorem 2]. Note first that under our assumptions u is a critical point of $\int |\nabla u| + \Phi(u)$ with Φ of class $C^{1,1}$. Hence u satisfies a semilinear equation of the type $\Delta u = f(u)$ with f Lipschitz. Hence, by standard elliptic regularity and “semilinear bootstrap” we have $u \in C_{\text{loc}}^{2,\alpha}(\mathbb{R}^n)$. This qualitative regularity is enough in order to justify the computations below.

Fix $x_0 \in \mathbb{R}^N$ and let $u_r(x) := \frac{u(x_0 + rx)}{r}$. Then

$$\mathcal{W}(u, x_0, r) = \int_{B_1} |\nabla u_r|^2 dx + \int_{B_1} \Phi(r u_r) dx - \int_{\partial B_1} u_r^2 d\sigma.$$

Noticing that $r \frac{d}{dr} u_r = \nabla u_r \cdot x - u_r$ and using the equation of u_r , we obtain

$$\begin{aligned} \frac{d}{dr} \int_{B_1} |\nabla u_r|^2 dx &= \frac{2}{r} \int_{B_1} \nabla u_r \cdot \nabla (\nabla u_r \cdot x - u_r) dx \\ &= -\frac{2}{r} \int_{B_1} \Delta u_r (\nabla u_r \cdot x - u_r) dx + \frac{2}{r} \int_{\partial B_1} (\nabla u_r \cdot x) (\nabla u_r \cdot x - u_r) d\sigma \\ &= -\int_{B_1} \Phi'(r u_r) (\nabla u_r \cdot x - u_r) dx + \frac{2}{r} \int_{\partial B_1} (\nabla u_r \cdot x) (\nabla u_r \cdot x - u_r) d\sigma. \end{aligned}$$

Similar,

$$\begin{aligned} \frac{d}{dr} \left(\int_{B_1} \Phi(r u_r) dx \right) &= \int_{B_1} \Phi'(r u_r) (\nabla u_r \cdot x) dx, \\ -\frac{d}{dr} \int_{\partial B_1} u_r^2 d\sigma &= -\frac{2}{r} \int_{\partial B_1} u_r (\nabla u_r \cdot x - u_r) d\sigma. \end{aligned}$$

Summing and rearranging terms, we find

$$\frac{d}{dr} \mathcal{W}(u, x_0, r) = \frac{2}{r} \int_{\partial B_1} (\nabla u_r \cdot x - u_r)^2 d\sigma + \frac{1}{r} \int_{B_1} u_r \Phi'_{1/r}(u_r).$$

Changing variables $x \rightarrow \frac{x-x_0}{r}$, (5.14) follows. \square

Proof of Proposition 5.1. By scaling, $\{u_{\varepsilon_j}\}_{j \in \mathbb{N}}$ is a family of minimizers of (1.1) in \mathbb{R}^N and thus, by Lemma 5.2, Lemma 5.3, Corollary 5.4 and using a standard diagonal argument, we deduce the existence of sequences $\varepsilon_\ell = \varepsilon_{j_\ell}$, $\delta_\ell \rightarrow 0$ and a minimizer u_0 of (1.5) in \mathbb{R}^N with $0 \in \partial\{u_0 > 0\}$ such that (5.1) and (5.2) are satisfied. The fact that u_0 is nontrivial follows by uniform non-degeneracy (Lemma 4.2).

We are left to show that u_0 is 1-homogeneous. To see this, we use Weiss' monotonicity formula. For every $\varepsilon \in (0, 1)$, we consider the function

$$r \rightarrow \mathcal{W}_\varepsilon(u_\varepsilon, r) := r^{-N} \int_{B_r} |\nabla u_\varepsilon|^2 + \Phi_\varepsilon(u_\varepsilon) \, dx - r^{-1-N} \int_{\partial B_r} u_\varepsilon^2 \, d\sigma.$$

Noticing that $\mathcal{W}_\varepsilon(u_\varepsilon, r) = \mathcal{W}(u, r/\varepsilon)$, we easily compute

$$\frac{d}{dr} \mathcal{W}_\varepsilon(u_\varepsilon, r) = \frac{1}{\varepsilon} \frac{d}{dr} \mathcal{W}(u, r/\varepsilon) = 2r^{-N} \int_{\partial B_r} \left(\partial_n u_\varepsilon - \frac{u_\varepsilon}{r} \right)^2 \, d\sigma + r^{-1-N} \int_{B_r} u_\varepsilon \Phi'_\varepsilon(u_\varepsilon) \, d\sigma,$$

and thus, integrating and neglecting the second term in the r.h.s., we deduce

$$\mathcal{W}(u, R/\varepsilon) - \mathcal{W}(u, \varrho/\varepsilon) \geq 2 \int_{\varrho}^R r^{-N} \int_{\partial B_r} \left(\partial_n u_\varepsilon - \frac{u_\varepsilon}{r} \right)^2 \, d\sigma \, dr, \quad (5.15)$$

for every $0 < \varrho < R$ fixed. On the other hand, since u is globally Lipschitz and $\Phi \leq 1$, we have

$$\mathcal{W}(u, r) \leq r^{-N} \int_{B_r} |\nabla u_\varepsilon|^2 + \Phi_\varepsilon(u_\varepsilon) \, dx \leq c_N (1 + \|\nabla u\|_{L^\infty(\mathbb{R}^N)}) < +\infty, \quad \forall r > 0.$$

This, together with the monotonicity $r \rightarrow \mathcal{W}(u, r)$, yields $\mathcal{W}(u, r) \rightarrow l$ as $r \rightarrow +\infty$, for some $l < +\infty$ (depending on u). Consequently, taking $\varepsilon = \varepsilon_\ell$ and passing to the limit as $\ell \rightarrow +\infty$ in (5.15), we obtain by H^1_{loc} and C^α_{loc} convergence

$$\int_{\varrho}^R r^{-N} \int_{\partial B_r} \left(\partial_n u_0 - \frac{u_0}{r} \right)^2 \, d\sigma \, dr = 0.$$

By the arbitrariness of ϱ and R , it follows $\partial_n u_0 = \frac{u_0}{r}$ in ∂B_r , for every $r > 0$, that is, u_0 is 1-homogeneous. \square

Proof of Proposition 1.5. Let $\{R_j\}_{j \in \mathbb{N}}$ be any sequence satisfying $R_j \rightarrow +\infty$ as $j \rightarrow +\infty$, and let $\varepsilon_j := \frac{1}{R_j}$. Let ε_{j_ℓ} , δ_ℓ , $u_{\varepsilon_{j_\ell}}$ and u_0 as in Proposition 5.1 and $R_{j_\ell} := \frac{1}{\varepsilon_{j_\ell}}$. Then, since u_0 is 1 homogeneous, (1.11) and (1.12) follow by scaling back to u into (5.1) and (5.2). \square

6. Improvement of flatness

This section is devoted to the proof of Theorem 2.1. As mentioned in the introduction, its proof can be regarded as a suitable “interpolation” of the methods by De Silva [15] and Savin [22], and requires some auxiliary results: a uniform Hölder type estimate given in Lemma 6.3 and Lemma 6.4, and a compactness result provided by Lemma 6.5. Further, we will crucially use the 1D solutions studied in Lemma 3.1 and their truncations (cf. Remark 3.2).

Definition 6.1. Let u_ε be a critical point of (1.1) in $B_R \subset \mathbb{R}^N$.

- We say that u_ε satisfies $\text{Flat}_1(\nu, \delta, R)$ if

$$\begin{aligned} u_\varepsilon(x) - \nu \cdot x &\leq \delta R && \text{in } B_R \cap \{u_\varepsilon \geq \vartheta_1 \varepsilon\} \\ -\delta R &\leq u_\varepsilon(x) - \nu \cdot x && \text{in } B_R. \end{aligned} \quad (6.1)$$

- We say that u_ε satisfies $\text{Flat}_2(\nu, \delta, R)$ if

$$w_\varepsilon^\varepsilon(\nu \cdot x - \delta R) < u_\varepsilon(x) < w_\varepsilon^{-\varepsilon}(\nu \cdot x + \delta R) \quad \text{in } B_R. \quad (6.2)$$

Lemma 6.2. There exist $\varepsilon_0, \delta_0 \in (0, 1)$ depending only on ϑ_1, ϑ_2 and $c_1 > 0$ as (2.1), such that for every $R > 0$, every $\nu \in \mathbb{S}^{N-1}$, every $\varepsilon/R \in (0, \varepsilon_0)$, $\delta \in [0, \delta_0]$ and every critical point u_ε of (1.1) in B_R , we have

$$u_\varepsilon \text{ satisfies } \text{Flat}_1(\nu, \delta, R) \Rightarrow u_\varepsilon \text{ satisfies } \text{Flat}_2(\nu, \delta + \sqrt{\varepsilon/R}, (1 - \sqrt{\varepsilon/R})R), \quad (6.3)$$

$$u_\varepsilon \text{ satisfies } \text{Flat}_2(\nu, \delta, R) \Rightarrow u_\varepsilon \text{ satisfies } \text{Flat}_1(\nu, \delta + \sqrt{\varepsilon/R}, (1 - \sqrt{\varepsilon/R})R). \quad (6.4)$$

Proof. Let $\varepsilon_0 \in (0, 1)$ as in Lemma 3.4, $\varepsilon \in (0, \varepsilon_0)$, and set $U_\varepsilon := \{u_\varepsilon \geq \vartheta_1 \varepsilon\}$. By scaling, we may assume $R = 1$ while, up to a rotation of the coordinate system, we can set $\nu = e_N$.

Step 1. Let us prove first (6.3). Assume that u_ε satisfies $\text{Flat}_1(\nu, \delta, 1)$, as defined in (6.1). On the one hand we have $u_\varepsilon(x) \geq x_N - \delta$ in B_1 . Then, by the first inequality in (3.9) with $\sigma = 1/2$, we have

$$u_\varepsilon(x) \geq x_N - \delta \geq w_\varepsilon^\varepsilon(x_N - \delta - \sqrt{\varepsilon}) \quad \text{in } B_1 \cap \{w_\varepsilon^\varepsilon(x_N - \delta - \sqrt{\varepsilon}) > 0\}. \quad (6.5)$$

Further, since $u_\varepsilon \geq 0$, the same inequality holds true in $B_1 \cap \{w_\varepsilon^\varepsilon(x_N - \delta - \sqrt{\varepsilon}) = 0\}$ and the first inequality in (6.3) follows.

To show the second inequality, we use that, on the other hand, $u_\varepsilon(x) \leq x_N + \delta$ in $B_1 \cap U_\varepsilon$. Then, by the second inequality in (3.10), we have

$$u_\varepsilon(x) \leq x_N + \delta \leq w_\varepsilon^{-\varepsilon}(x_N + \delta + \sqrt{\varepsilon}) - \frac{\sqrt{\varepsilon}}{2} \quad \text{in } B_1 \cap U_\varepsilon. \quad (6.6)$$

Now notice that by Lemma 3.6 (cf. (3.15)), we have

$$u_\varepsilon \leq 3\vartheta_1\varepsilon e^{-\frac{\varepsilon^{-1/4}}{4c_1^{1/2}}} \quad \text{in } B_{1-\sqrt{\varepsilon}} \setminus V_\varepsilon, \quad V_\varepsilon := B_1 \cap \{x : d(x, U_\varepsilon) \leq \varepsilon^{3/4}\}. \quad (6.7)$$

Thanks to (3.3) (with $|\tau| = \varepsilon$), we also know that $w_\varepsilon^{-\varepsilon}(x_N + \delta + \sqrt{\varepsilon}) \geq \frac{1}{\sqrt{c_1}}\varepsilon^{3/2}$ and thus by (6.7)

$$u_\varepsilon < w_\varepsilon^{-\varepsilon}(x_N + \delta + \sqrt{\varepsilon}) \quad \text{in } B_{1-\sqrt{\varepsilon}} \setminus V_\varepsilon,$$

for every $\varepsilon \in (0, \varepsilon_0)$, taking eventually ε_0 smaller. We are left to check that $u_\varepsilon(x) < w_\varepsilon^{-\varepsilon}(x_N + \delta + \sqrt{\varepsilon})$ in $B_{1-\sqrt{\varepsilon}} \cap (V_\varepsilon \setminus U_\varepsilon)$. Let $x \in B_{1-\sqrt{\varepsilon}} \cap (V_\varepsilon \setminus U_\varepsilon)$. Let $\bar{x} \in B_1 \cap U_\varepsilon$ such that $|x - \bar{x}| \leq \varepsilon^{3/4}$. From (6.6) (using $u_\varepsilon \geq 0$) we know that $w_\varepsilon^{-\varepsilon}(\bar{x}_N + \delta + \sqrt{\varepsilon}) \geq \sqrt{\varepsilon}/2$. Hence (using that $w_\varepsilon^{-\varepsilon}$ is 1-Lipschitz),

$$u_\varepsilon(x) \leq \vartheta_1\varepsilon \leq \sqrt{\varepsilon}/2 - \varepsilon^{3/4} \leq w_\varepsilon^{-\varepsilon}(\bar{x}_N + \delta + \sqrt{\varepsilon}) - \varepsilon^{3/4} \leq w_\varepsilon^{-\varepsilon}(x_N + \delta + \sqrt{\varepsilon}).$$

This completes the proof of (6.3).

Step 2. Now we show (6.4). Assume $u_\varepsilon(x) > w_\varepsilon^\varepsilon(x_N - \delta)$ in B_1 . Then, by the second inequality in (3.9) (with $\sigma = 1/2$), we obtain

$$u_\varepsilon(x) > w_\varepsilon^\varepsilon(x_N - \delta) > x_N - \delta - \frac{\sqrt{\varepsilon}}{2} > x_N - \delta - \sqrt{\varepsilon} \quad \text{in } B_1,$$

and the first inequality in (6.4) follows. On the other hand, if $u_\varepsilon(x) < w_\varepsilon^{-\varepsilon}(x_N + \delta)$ in B_1 , the first inequality in (3.10) yields

$$u_\varepsilon(x) < w_\varepsilon^{-\varepsilon}(x_N + \delta) < x_N + \delta + \frac{\sqrt{\varepsilon}}{2} < x_N + \delta + \sqrt{\varepsilon} \quad \text{in } B_1 \cap \{x_N \geq y_\varepsilon^{-\varepsilon} - \delta\},$$

where $y_\varepsilon^{-\varepsilon}$ is as in Lemma 3.4. Finally, since $u_\varepsilon(x) \geq \vartheta_1\varepsilon$ and the assumption imply $w_\varepsilon^{-\varepsilon}(x_N + \delta) \geq \vartheta_1\varepsilon$, we deduce, by monotonicity, that $x_N + \delta \geq 0 \geq y_\varepsilon^{-\varepsilon}$ in $U_\varepsilon = \{u_\varepsilon \geq \vartheta_1\varepsilon\}$. Thus $B_1 \cap U_\varepsilon \subset B_1 \cap \{x_N \geq y_\varepsilon^{-\varepsilon} - \delta\}$ and the second inequality in (6.4) follows too. \square

Lemma 6.3. *There exist $\delta_0, c_0 \in (0, 1)$ and $\theta_0 \in (\frac{1}{2}, 1)$ depending only on $N, \vartheta_1, \vartheta_2$ and c_1 as in (2.1) such that for every $R > 0$, every $\delta \in (0, \delta_0)$, every $a \in \mathbb{R}$ and $b \leq 0$ such that $a + |b| = \delta R$, every $\varepsilon/R \in (0, c_0\delta)$ and every critical point u_ε of (1.1) in B_R satisfying*

$$\begin{aligned} w_\varepsilon^\varepsilon(x_N - a) &\leq u_\varepsilon(x) \leq w_\varepsilon^{-\varepsilon}(x_N - b) \quad \text{in } B_R, \\ u_\varepsilon(0) &\in [\vartheta_1\varepsilon, \vartheta_2\varepsilon], \end{aligned} \quad (6.8)$$

where $w_\varepsilon^\varepsilon$ and $w_\varepsilon^{-\varepsilon}$ are as in Remark 3.2 with $w_\varepsilon^\varepsilon(0) = w_\varepsilon^{-\varepsilon}(0) = \vartheta_1\varepsilon$, then there exist $a' \in \mathbb{R}, b' \leq 0$ such that

$$\begin{aligned}
w_\varepsilon^\varepsilon(x_N - a') &\leq u_\varepsilon(x) \leq w_\varepsilon^{-\varepsilon}(x_N - b') \quad \text{in } B_{R/4}, \\
b &\leq b' \leq a' \leq a, \\
a' + |b'| &\leq \theta_0(a + |b|).
\end{aligned} \tag{6.9}$$

Proof. By scaling, we may assume $R = 1$. Set $u = u_\varepsilon$, $w^\varepsilon = w_\varepsilon^\varepsilon$, $w^{-\varepsilon} = w_\varepsilon^{-\varepsilon}$, and define

$$w^{\varepsilon,a}(x_N) := w^\varepsilon(x_N - a), \quad w^{-\varepsilon,b}(x_N) := w^{-\varepsilon}(x_N - b).$$

Notice that, up to replace δ with $\delta + 1/j$ and then taking the limit as $j \rightarrow +\infty$, we may assume

$$w^{\varepsilon,a} < u < w^{-\varepsilon,b} \quad \text{in } B_1,$$

and, since $0 \in \{\vartheta_1\varepsilon \leq u_\varepsilon \leq \vartheta_2\varepsilon\}$, we also have

$$a \geq -c\varepsilon, \quad |b| \leq \delta + c\varepsilon, \tag{6.10}$$

where $c > 1$ is as in Lemma 3.3. This can be easily verified since $w^\varepsilon(0) = \vartheta_1\varepsilon$ and $\{\vartheta_1\varepsilon \leq w^\varepsilon \leq \vartheta_2\varepsilon\} \subset \{|x_N| \leq c\varepsilon\}$ by Lemma 3.3.

We define

$$\delta_0 := \frac{1}{32}, \quad c_0 := \frac{1}{16c}, \quad \theta_0 := 1 - c_N, \tag{6.11}$$

where $c > 1$ is as in Lemma 3.3 respectively (depending only on ϑ_1 , ϑ_2 and $c_1 > 0$ in (2.1)), and $c_N \in (0, 1)$ is the dimensional constant appearing in (6.14) (notice that we may assume $\theta_0 > \frac{1}{2}$ taking eventually c_N smaller). In particular, since $\delta \in (0, \delta_0)$, we have

$$\{\varepsilon\vartheta_1 \leq u \leq \varepsilon\vartheta_2\} \subset \{|x_N| < \tfrac{1}{32}\}. \tag{6.12}$$

Fix $y = (y', y_N) = (0, \frac{1}{8})$. We consider the following alternative. Either:

$$(a) \quad w^{-\varepsilon,b}(y) - u(y) \leq u(y) - w^{\varepsilon,a}(y)$$

or

$$(b) \quad w^{-\varepsilon,b}(y) - u(y) \geq u(y) - w^{\varepsilon,a}(y)$$

First case. Assume (a) holds. We first prove that

$$u > w^{\varepsilon,a-c_N\delta} \quad \text{in } B_{15/16} \cap \{|x_N| \geq \tfrac{1}{16}\}, \tag{6.13}$$

for some $c_N \in (0, 1)$. Let $v := u - w^{\varepsilon,a}$. In view of (6.12), v is harmonic and positive in $B_1 \cap \{|x_N| > \frac{1}{32}\}$ and so, by the Harnack inequality, it follows

$$\inf_{B_{15/16} \cap \{|x_N| \geq 1/16\}} v \geq 4c_N v(y) \geq 2c_N [w^{-\varepsilon, b}(y) - w^{\varepsilon, a}(y)] \geq c_N \delta,$$

for some $c_N > 0$. To justify the last inequality we proceed as follows. If $\tilde{a} > a$ and $\tilde{b} > b$ are such that $w^{\varepsilon, \delta}(\tilde{a}) = w^{-\varepsilon, -\delta}(\tilde{b}) = \vartheta_2 \varepsilon$, then $|\tilde{a} - a| \leq c\varepsilon$, $|\tilde{b} - b| \leq c\varepsilon$ where $c > 0$ is the constant appearing in the statement of Lemma 3.3. Consequently, since $w^{\varepsilon, a}(y) = \vartheta_2 \varepsilon + (1 + \varepsilon)(y_N - \tilde{a})$, $w^{-\varepsilon, b}(y) = \vartheta_2 \varepsilon + (1 - \varepsilon)(y_N - \tilde{b})$ and $c > 1$, we find

$$\begin{aligned} w^{-\varepsilon, b}(y) - w^{\varepsilon, a}(y) &= (1 - \varepsilon)(y_N - \tilde{b}) - (1 + \varepsilon)(y_N - \tilde{a}) = \tilde{a} - \tilde{b} - \frac{\varepsilon}{4} + \varepsilon(\tilde{a} + \tilde{b}) \\ &\geq \delta - 2c\varepsilon - \frac{\varepsilon}{4} + \varepsilon(a + b - 2c\varepsilon) \geq \delta - 6c\varepsilon - \varepsilon - \varepsilon\delta, \end{aligned}$$

thanks to (6.10). Further, recalling that $\varepsilon < c_0 \delta$ by assumption, it follows

$$w^{-\varepsilon, b}(y) - w^{\varepsilon, a}(y) \geq (1 - 8cc_0)\delta > \frac{1}{2}\delta,$$

in view of the definition of c_0 in (6.11). As a consequence, $u \geq w^{\varepsilon, a} + c_N \delta$ in $B_{15/16} \cap \{|x_N| \geq \frac{1}{16}\}$ and thus, using that $w^{\varepsilon, a}$ is a line with slope $1 + \varepsilon$ in $\{w^{\varepsilon, a} > \vartheta_2 \varepsilon\}$ and $\varepsilon < 1$, we deduce (6.13).

The second step is to show

$$u \geq w^{\varepsilon, a - c_N \delta} \quad \text{in } B_{1/4}, \quad (6.14)$$

for some new $c_N \in (0, 1)$. If (6.14) holds true, then (6.9) follows by setting $a' = a - c_N \delta$, $b' = b$, in view of the definition of θ_0 .

To prove (6.14) we use a sliding argument: given any smooth, nonnegative and bounded h , we define the family of functions

$$v_\lambda(x) := w^{\varepsilon, a}(x_N + \lambda h(x)), \quad x \in B_1, \quad \lambda \in [0, c_N \delta].$$

Notice that $v_0 = w^{\varepsilon, a}$. Using the equation of w^ε , it is not difficult to check that

$$\Delta v_\lambda = \frac{1}{2} \Phi'_\varepsilon(v_\lambda) (1 + 2\lambda \partial_N h + \lambda^2 |\nabla h|^2) + \lambda \dot{w}^\varepsilon \Delta h, \quad (6.15)$$

where $\partial_N := \partial_{x_N}$. We choose $h(x) := \tilde{h}(x - y)$, where \tilde{h} is the unique radially decreasing harmonic function in $B_{1/2} \setminus B_{1/32}$ satisfying $\tilde{h} = 1$ in $\overline{B}_{1/32}$ and $\tilde{h} = 0$ in $\mathbb{R}^N \setminus B_{1/2}$. Consequently,

$$\Delta v_\lambda > \frac{1}{2} \Phi'_\varepsilon(v_\lambda) \quad \text{in } D := B_{1/2}(y) \cap \{x_N < \frac{1}{16}\}, \quad (6.16)$$

for every $\lambda \in (0, c_N \delta]$. This follows neglecting the nonnegative terms in (6.15) and noticing that $\partial_N h > 0$ in D by construction. On the other hand,

$$v_\lambda < u \quad \text{in } \partial D, \quad (6.17)$$

for every $\lambda \in [0, c_N \delta]$. Indeed, recalling that $h = 0$ in $\partial B_{1/2}(y)$ it follows $v_\lambda = w^{\varepsilon, a} < u$ in $\partial D \cap \{x_N < \frac{1}{16}\}$ while, since $h \leq 1$ and $\lambda \leq c_N \delta$, we have $v_\lambda \leq w^{\varepsilon, a - c_N \delta}$ and so $v_\lambda < u$ in $\partial D \cap \{x_N = \frac{1}{16}\}$ in view of (6.13). Now, we define

$$\lambda_* := \max\{\lambda \in [0, c_N \delta] : v_\lambda \leq u \text{ in } D\},$$

and show that $\lambda_* = c_N \delta$. If this is not true, there must be $\lambda \in [0, c_N \delta)$ and $x_\lambda \in \overline{D}$ such that $v_\lambda \leq u$ in D , with $v_\lambda(x_\lambda) = u(x_\lambda)$. Recalling that $v_0 = w^{\varepsilon, a}$ and that $w^{\varepsilon, a} < u$ by assumption, we immediately see that $\lambda > 0$ and, by (6.17), it must be $x_\lambda \in D$. Thus, using the equation of u (or equivalently u) and (6.16), we obtain that the function $\tilde{v}_\lambda := u - v_\lambda$ satisfies

$$\begin{cases} \tilde{v}_\lambda \geq 0 & \text{in } D \\ \tilde{v}_\lambda(x_\lambda) = 0, \Delta \tilde{v}_\lambda(x_\lambda) < 0, \end{cases}$$

which leads to a contradiction since $x_\lambda \in D$ is a minimum point for \tilde{v}_λ . Combining (6.13) with $\lambda_* = c_N \delta$, and noticing that $B_{1/4} \subset B_{1/2}(y)$, we deduce

$$u(x) \geq w^\varepsilon(x_N - a + c_N \delta h(x)) \quad \text{in } B_{1/4},$$

and thus, since $h \geq c_N$ in $B_{1/4}$ for some new constant $c_N > 0$ by construction, the monotonicity of w^ε yields (6.14).

Second case. Assume now that (b) holds. In this case, following the proof of (6.13), we find

$$u < w^{-\varepsilon, b + c_N \delta} \quad \text{in } B_{15/16} \cap \{|x_N| \geq \frac{1}{16}\},$$

where $c_N > 0$ can be taken as in (6.13). So, following the ideas of Step 1, we must prove

$$u \leq w^{-\varepsilon, b + c_N \delta} \quad \text{in } B_{1/4}, \quad (6.18)$$

where $c_N \in (0, 1)$ is as in (6.14). As above, (6.18) implies (6.9) taking $a' = a$ and $b' = b + c_N \delta$.

To do so, we consider

$$v_\lambda(x) := w^{-\varepsilon, a}(x_N + \lambda h(x)), \quad x \in B_1, \quad \lambda \in [-c_N \delta, 0],$$

where h is as in Step 1 (note however that now $\lambda < 0$). Using (6.15), we deduce $\Delta v_\lambda < \frac{1}{2} \Phi'_\varepsilon(v_\lambda)$ in D , for every $\lambda \in [-c_N \delta, 0)$. To see this, it is enough to notice that

$$2\partial_N h + \lambda |\nabla h|^2 \geq \bar{c}_N + \lambda |\nabla h|^2 > 0 \quad \text{in } D,$$

for some small $\bar{c}_N > 0$, if $|\lambda|$ is small enough and so, choosing eventually δ_0 smaller (depending only on N), the above inequality is satisfied for $\lambda \in [-c_N\delta, 0)$. Proceeding exactly as above, we find

$$\lambda_* := \min\{\lambda \in [-c_N\delta, 0] : v_\lambda \geq u \text{ in } D\} = -c_N\delta,$$

and we are led to

$$u(x) \leq w^{-\varepsilon}(x_N - b - c_N\delta) \quad \text{in } B_{1/4},$$

for some new $c_N > 0$, which is (6.18). \square

Lemma 6.4. *There exist $\alpha, \tilde{\delta}_0 \in (0, 1)$ and $C > 0$ depending only on $N, \vartheta_1, \vartheta_2$ and c_1 as in (2.1) such that for every $\delta \in (0, \tilde{\delta}_0)$, every $\varepsilon \in (0, \delta^2)$ and every critical point u_ε of (1.1) in B_1 satisfying*

$$\begin{aligned} u_\varepsilon(x) - x_N &\leq \delta && \text{in } B_1 \cap \{u_\varepsilon \geq \vartheta_1\varepsilon\} \\ -\delta &\leq u_\varepsilon(x) - x_N && \text{in } B_1, \end{aligned} \quad (6.19)$$

with $u_\varepsilon(0) \in [\vartheta_1\varepsilon, \vartheta_2\varepsilon]$, then the function

$$v_{\varepsilon, \delta}(x) := \frac{u_\varepsilon(x) - x_N}{\delta}$$

satisfies

$$\begin{aligned} v_{\varepsilon, \delta}(x) - v_{\varepsilon, \delta}(z) &\leq \omega_\delta(x - z) && \text{in } B_{1/2} \cap \{u_\varepsilon \geq \vartheta_1\varepsilon\} \\ -\omega_\delta(x - z) &\leq v_{\varepsilon, \delta}(x) - v_{\varepsilon, \delta}(z) && \text{in } B_{1/2}, \end{aligned} \quad (6.20)$$

for every $z \in B_{1/2} \cap \{u_\varepsilon \geq \vartheta_1\varepsilon\}$, where

$$\omega_\delta(y) := C(\delta + |y|)^\alpha.$$

Proof. Let $\delta_0, \theta_0 \in (0, 1)$ and $c_0 \in (0, 1)$ as in Lemma 6.3, and $\varepsilon_0 \in (0, 1)$ as in Lemma 6.2. We set

$$\tilde{\delta}_0 := \min\{\delta_0/4, \sqrt{\varepsilon_0/4}, c_0/4\},$$

and take $\delta \in (0, \tilde{\delta}_0)$, $\varepsilon \in (0, \delta^2)$. Notice that the definition of $\tilde{\delta}_0$ guarantees $4\delta < \delta_0$ and $4\varepsilon < \varepsilon_0$. For simplicity we also set $u = u_\varepsilon$, $w^\varepsilon = w_\varepsilon^\varepsilon$ and $w^{-\varepsilon} = w_\varepsilon^{-\varepsilon}$, and define

$$\kappa := \frac{1}{\tilde{\delta}_0}, \quad 0 < \alpha < |\log_4(\theta_0)|, \quad C \geq 4^{1+2\alpha}. \quad (6.21)$$

Step 1. We first prove that (6.20) holds true for every $z \in \{\vartheta_1\varepsilon \leq u_\varepsilon \leq \vartheta_2\varepsilon\} \cap B_{1/2}$. Let us set

$$\tilde{\delta} := 2(\delta + \sqrt{\varepsilon/2}).$$

Notice that $\varepsilon < \delta^2$ implies $\tilde{\delta} \leq 4\delta$ and thus, since $\tilde{\delta} \geq 2\delta$ by definition, it is equivalent to work with $\tilde{\delta}$ instead of δ , which is what we will do from now on.

So, we fix $j \in \mathbb{N}$ such that

$$4^{-j-2} \leq \frac{\tilde{\delta}}{4\tilde{\delta}_0} = \frac{\kappa}{4}\tilde{\delta} < 4^{-j-1}, \quad (6.22)$$

and we use the definition of $\tilde{\delta}$ to combine (6.19) and (6.3), which yield

$$w^\varepsilon(x_N - \tilde{\delta}) < u(x) < w^{-\varepsilon}(x_N + \tilde{\delta}) \quad \text{in } B_{3/4}.$$

Now, in view of (6.22), we have $\tilde{\delta} \leq 4^{-j}\tilde{\delta}_0$ and, since $\varepsilon < \delta^2$, we also have $\varepsilon \leq \tilde{\delta}^2 \leq c_0\tilde{\delta}$ and so we may apply Lemma 6.3 (rescaled and translated from B_1 to $B_{1/4}(z)$, i.e. applied to the function $u(z + 4 \cdot)$) iteratively on $B_{4^{-k}}(z)$ for $1 \leq k \leq j$, deducing the existence of a_k and b_k (with $a_0 = -b_0 = \tilde{\delta}$) for which

$$\begin{aligned} w^\varepsilon(x_N - z_N - a_k) &\leq u(x) \leq w^{-\varepsilon}(x_N - z_N - b_k) \quad \text{in } B_{4^{-k}}(z), \\ 0 < a_k + |b_k| &\leq 4\theta_0^k \tilde{\delta}. \end{aligned} \quad (6.23)$$

Then, applying (6.4) to (6.23) (choosing $R = 4^{-k}$ and $\delta = (a_k + |b_k|)4^k$) and recalling that $\theta_0 \in (\frac{1}{2}, 1)$, it follows

$$\begin{aligned} u(x) - (x_N - z_N) &\leq 2\theta_0^k \tilde{\delta} \quad \text{in } B_{4^{-k}/2}(z) \cap \{u_\varepsilon \geq \vartheta_1 \varepsilon\} \\ -2\theta_0^k \tilde{\delta} &\leq u(x) - (x_N - z_N) \quad \text{in } B_{4^{-k}/2}(z), \end{aligned} \quad (6.24)$$

for all $1 \leq k \leq j$ (notice that since $\varepsilon < \delta^2 < \tilde{\delta}^2$ and $\tilde{\delta}_0 < \sqrt{\varepsilon_0}$ we automatically have $\varepsilon < \varepsilon_0 4^{-k}$, for every $k \leq j$).

Now, assume that $|x - z| \geq \kappa \tilde{\delta}$. Then $4^{-j+n-2} \leq |z - x| < 4^{-j+n-1}$, for some $0 \leq n \leq j$ ($n \in \mathbb{N}$) by the definition of j . Applying (6.24) with $k = j - n$, we find

$$u(x) - x_N - (u(z) - z_N) = u(x) - (x_N - z_N) - u(z) \quad \begin{cases} \leq 4\theta_0^{j-n} \tilde{\delta} & \text{if } x \in \{u_\varepsilon \geq \vartheta_1 \varepsilon\} \\ \geq -4\theta_0^{j-n} \tilde{\delta}, \end{cases}$$

and thus

$$\begin{aligned} v(x) - v(z) &\leq 4\theta_0^{j-n} \quad \text{if } x \in \{u_\varepsilon \geq \vartheta_1 \varepsilon\} \\ -4\theta_0^{j-n} &\leq v(x) - v(z), \end{aligned} \quad (6.25)$$

where we have set $v := v_{\varepsilon, \delta}$ for simplicity. Using the definitions of α and C in (6.21) and that $|x - z| \geq 4^{-j+n-2}$, we have $4\theta_0^{j-n} \leq C|x - z|^\alpha$ and (6.20) follows.

If $|x - z| \leq \kappa\tilde{\delta}$, then, proceeding as above, we find that (6.25) holds true with $n = 0$ and so, since $4\theta_0^j \leq C(4^{-\alpha})^{j+2}$ for α and C as in (6.21), and $\kappa\tilde{\delta} \geq 4^{-j-2}$, we deduce

$$\begin{aligned} v(x) - v(z) &\leq C(\kappa\tilde{\delta})^\alpha \quad \text{if } x \in \{u_\varepsilon \geq \vartheta_1\varepsilon\} \\ -C(\kappa\tilde{\delta})^\alpha &\leq v(x) - v(z), \end{aligned}$$

and (6.20) follows.

Step 2. Now we consider the case $x, z \in \{u \geq \vartheta_2\varepsilon\} \cap B_{1/4}$. We fix $x_0 \in \partial\{u > \varepsilon\vartheta_2\} \cap B_{1/4}$ such that $|x - x_0| = \text{dist}(x, \partial\{u > \varepsilon\vartheta_2\}) := d(x)$.

Set $d := d(x)$ and assume first $\kappa\delta \vee |x - z| < d/4$. In this case, using *Step 1*, we easily obtain

$$|v(\xi) - v(x_0)| \leq C(\kappa\delta \vee |\xi - x_0|)^\alpha \leq 2^\alpha C d^\alpha, \quad \forall \xi \in B_{d/2}(x).$$

So, since v is harmonic in $B_d(x)$, we have

$$\sup_{B_{d/4}(x)} |\nabla v| \leq c_N \frac{\text{osc}_{B_{d/2}(x)} v}{d} \leq c_N C_\alpha d^{\alpha-1},$$

for some $C_\alpha > 0$ and thus $|v(x) - v(z)| \leq c_N C_\alpha d^{\alpha-1} |x - z| \leq C_\alpha (\kappa\delta \vee |x - z|)^\alpha$ for some new $C_\alpha > 0$.

On the other hand, if $\kappa\delta \vee |x - z| \geq d/4$, we may apply the estimate of *Step 1* twice to obtain

$$\begin{aligned} |v(x) - v(z)| &\leq |v(x) - v(x_0)| + |v(x_0) - v(z)| \\ &\leq C[(\kappa\delta \vee |x - x_0|)^\alpha + (\kappa\delta \vee |z - x_0|)^\alpha] \\ &\leq C\{[\kappa\delta \vee d(x)]^\alpha + [\kappa\delta \vee (|x - z| + d(x))]^\alpha\} \leq C_\alpha (\kappa\delta \vee |x - z|)^\alpha, \end{aligned}$$

for some $C, C_\alpha > 0$ and our statement follows.

Step 3. If $x \in \{u \leq \vartheta_1\varepsilon\}$ and $z \in \{u > \vartheta_2\varepsilon\}$ then there exists $\bar{z} \in \{\vartheta_1\varepsilon \leq u_\varepsilon \leq \vartheta_2\varepsilon\}$ which belongs to the segment xz . Hence, using the previous steps

$$\begin{aligned} v(x) - v(z) &\geq v(x) - v(\bar{z}) - |v(\bar{z}) - v(z)| \\ &\geq -C(\kappa\delta \vee |x - \bar{z}|)^\alpha - C(\kappa\delta \vee |x - \bar{z}|)^\alpha \geq -C(\kappa\delta \vee |x - z|)^\alpha, \end{aligned}$$

and the proof of (6.20) is complete. \square

Lemma 6.5. *There exists a Hölder continuous function $v : \{x_N \geq 0\} \cap \overline{B_{1/4}} \rightarrow \mathbb{R}$, harmonic in $\{x_N > 0\} \cap B_{1/4}$ and with $\|v\|_{L^\infty} = 1$ such that for every sequence $\delta_j \rightarrow 0^+$, every $\varepsilon_j \in (0, \delta_j^2)$ and every critical point u_{ε_j} of (1.1) in B_2 satisfying*

$$\begin{aligned} u_{\varepsilon_j}(x) - x_N &\leq \delta_j & \text{in } B_1 \cap \{u_{\varepsilon_j} \geq \vartheta_1\varepsilon_j\} \\ -\delta_j &\leq u_{\varepsilon_j}(x) - x_N & \text{in } B_1, \end{aligned} \tag{6.26}$$

with $u_{\varepsilon_j}(0) \in [\vartheta_1 \varepsilon_j, \vartheta_2 \varepsilon_j]$, then, setting

$$v_j(x) := \frac{u_{\varepsilon_j}(x) - x_N}{\delta_j},$$

the sequence of graphs

$$G_j = \{(x, v_j(x)) : x \in \{u_{\varepsilon_j} \geq \vartheta_1 \varepsilon_j\} \cap B_{1/4}\} \quad (6.27)$$

converge in the Hausdorff distance in \mathbb{R}^{N+1} to

$$G = \{(x, v(x)) : x \in \{x_N \geq 0\} \cap B_{1/4}\}, \quad (6.28)$$

as $j \rightarrow +\infty$, up to passing to a suitable subsequence.

Proof. Let $\alpha \in (0, 1)$ and $\kappa, C > 0$ as in Lemma 6.4. Let $\delta_j \rightarrow 0^+$, $\varepsilon_j \in (0, \delta_j^2)$ and set $U_j := \{u_{\varepsilon_j} > \vartheta_1 \varepsilon_j\} \cap B_{1/4}$, $H := \{x_N > 0\} \cap B_{1/4}$.

Step 1: Compactness. We show that there is v (harmonic in H and α -Hölder in \overline{H} and with L^∞ norm bounded by 1) such that for every $\sigma \in (0, 1/4)$,

$$\|v_j - v\|_{L^\infty(H_\sigma)} \rightarrow 0, \quad (6.29)$$

as $j \rightarrow +\infty$, up to passing to a suitable subsequence, where $H_\sigma := \{x_N > \sigma\} \cap B_{1/4}$.

By (6.26), there is $j_\sigma \in \mathbb{N}$, such that $H_{\sigma/2} \subset U_j$ and $\|v_j\|_{L^\infty(H_\sigma)} \leq 1$ (this follows (6.26) by δ_j) and every $j \geq j_\sigma$. In addition, v_j is harmonic in U_σ and thus, by standard elliptic estimates and a diagonal procedure, there exists a harmonic function v in H such that $v_j \rightarrow v$ locally uniformly in H , up to passing to a suitable subsequence. On the other hand, by (6.20), we have

$$|v_j(x) - v_j(y)| \leq C(\delta_j + |x - y|)^\alpha,$$

for every $x, y \in \overline{U_j}$, and thus, passing to the limit as $j \rightarrow +\infty$, we obtain that v can be continuously extended up to ∂U and $v \in C^\alpha(\overline{H})$ with $\|v\|_{L^\infty(H)} \leq 1$.

Step 2: Convergence of graphs. Fix $\sigma \in (0, \frac{1}{4})$, $x \in \overline{H}$, $p := (x, v(x)) \in G$ and set $q := (y, v(y))$, where $y \in H_{\sigma/2}$ is taken such that $|x - y| \leq \sigma$. Then, by the C^α estimate proved above, we obtain

$$|p - q|^2 = |x - y|^2 + |v(x) - v(y)|^2 \leq \sigma^2 + C^2 \sigma^{2\alpha} \leq C^2 \sigma^{2\alpha},$$

for some new $C > 0$. Now, if j is large enough, we have $H_{\sigma/2} \subset U_j$ and so

$$\text{dist}(q, G_j)^2 = \inf_{y' \in \overline{U_j}} |y - y'|^2 + |v(y) - v_j(y')|^2 \leq |v(y) - v_j(y)|^2 \leq \|v - v_j\|_{L^\infty(U_{\sigma/2})}^2,$$

from which we deduce

$$\text{dist}(p, G_j) \leq |p - q| + \text{dist}(q, G_j) \leq C\sigma^\alpha + \|v - v_j\|_{L^\infty(U_{\sigma/2})} \leq C\sigma^\alpha, \quad (6.30)$$

for some new $C > 0$, for every j large enough, in view of (6.29).

On the other hand, given any sequence $p_j = (x_j, v_j(x_j)) \in G_j$, if j is large enough we may take $y_j \in H_{\sigma/2}$ such that $\frac{\sigma}{2} \leq |x_j - y_j| \leq \sigma$ with j such that $\delta_j \leq \frac{\sigma}{2}$. Consequently, setting $q_j = (y_j, v_j(y_j))$, we have by (6.20)

$$|p_j - q_j|^2 = |x_j - y_j|^2 + |v_j(x_j) - v_j(y_j)|^2 \leq \sigma^2 + C^2\sigma^{2\alpha} \leq C^2\sigma^{2\alpha}.$$

Further, as above

$$\text{dist}(q_j, G) = \inf_{y' \in H} |y_j - y'|^2 + |v_j(y_j) - v(y')|^2 \leq |v_j(y_j) - v(y_j)|^2 \leq \|v - v_j\|_{L^\infty(U_{\sigma/2})},$$

and thus, by (6.29),

$$\text{dist}(p_j, G) \leq C\sigma^\alpha + \|v - v_j\|_{L^\infty(U_{\sigma/2})} \leq C\sigma^\alpha, \quad (6.31)$$

for j large enough. Since p , p_j and $\sigma > 0$ are arbitrary, the thesis follows by (6.30) and (6.31). \square

Proof of Theorem 2.1. By scaling, we may assume $R = 1$. Assume by contradiction that there are $\gamma \in (0, 1)$ and a sequence $\delta_j \rightarrow 0^+$ such that for every $\varrho_0 \in (0, 1)$, there is $\varepsilon_j \in (0, \delta_j^2)$, a solution $u_j := u_{\varepsilon_j}$ to (1.8) in B_1 satisfying

$$\begin{aligned} u_{\varepsilon_j}(x) - x_N &\leq \delta_j && \text{in } B_1 \cap \{u_{\varepsilon_j} \geq \vartheta_1 \varepsilon_j\} \\ -\delta_j &\leq u_{\varepsilon_j}(x) - x_N && \text{in } B_1, \end{aligned} \quad (6.32)$$

with $u_{\varepsilon_j}(0) \in [\vartheta_1 \varepsilon_j, \vartheta_2 \varepsilon_j]$, such that for every $\nu \in \mathbb{S}^{N-1}$, either

$$\begin{aligned} u_{\varepsilon_j}(x) - \nu \cdot x &\leq \delta_j \varrho_0^{1+\gamma} && \text{in } B_{\varrho_0} \cap \{u_{\varepsilon_j} \geq \vartheta_1 \varepsilon_j\} \\ -\delta_j \varrho_0^{1+\gamma} &\leq u_{\varepsilon_j}(x) - \nu \cdot x && \text{in } B_{\varrho_0} \cap \{u_{\varepsilon_j} \geq 0\} \end{aligned} \quad (6.33)$$

or

$$|\nu - e_N| \leq \sqrt{2n} \delta_j \quad (6.34)$$

fails for $j \in \mathbb{N}$ large enough.

Step 1: Compactness. By Lemma 6.5, we have that the sequence

$$v_j := v_{\varepsilon_j, \delta_j} = \frac{u_j - x_N}{\delta_j}$$

converge uniformly on compact sets of $U := \{x_N > 0\} \cap B_{1/4}$ to some limit function $v \in C^\alpha(\overline{U})$ which is harmonic in U and, further, the sequence of graphs G_j defined in (6.27) converge in the Hausdorff distance in \mathbb{R}^{N+1} to the graph G defined in (6.28). In addition, since $0 \in \{\vartheta_1 \varepsilon_j \leq u_j \leq \vartheta_2 \varepsilon_j\}$ and $\varepsilon_j \in (0, \delta_j^2)$, then

$$0 < v_j(0) \leq \vartheta_2 \delta_j^2,$$

for every j , and thus $v(0) = 0$. Before moving forward, we define the even reflection of v w.r.t. the hyperplane $\{x_N = 0\}$

$$\tilde{v}(x) = \begin{cases} v(x', x_N) & \text{in } x_N \geq 0 \\ v(x', -x_N) & \text{in } x_N < 0, \end{cases}$$

defined in the whole $B_{1/4}$ and satisfying $\tilde{v} \in C^\alpha(B_{1/4})$.

Step 2. In this step we prove that $\partial_N \tilde{v} \leq 0$ in $\{x_N = 0\}$ in the viscosity sense, that is for every $\varphi \in C^\infty(B_1)$ such that $\varphi \leq \tilde{v}$ in $B_{1/4}$ with equality only at some $z \in \{x_N = 0\} \cap B_{1/4}$, then $\partial_N \varphi(z) \leq 0$.

By contradiction, we assume there is $\varphi \in C^\infty(B_1)$ and $z \in \{x_N = 0\} \cap B_{1/4}$ as above, with $\partial_N \varphi(z) > 0$. For simplicity, we assume $z = 0$, $\varphi(z) = 0$ (the same proof work in the general case with minor modifications). In addition, we may take φ to be a polynomial of degree 2 (cf. [9, Chapter 2]) with the form

$$\varphi(x) = mx_N + m' \cdot x' + x^T \cdot M \cdot x, \quad x \in B_r \quad (6.35)$$

for some vector (m', m) with $m > 0$, some matrix $M \in \mathbb{R}^{N,N}$ with $\text{tr}(M) = 0$ and some $r > 0$. This can be easily obtained by modifying a generic polynomial of degree 2, taking r small enough and using the assumption $\partial_N \varphi(0) > 0$. Taking eventually r smaller, we may also assume $\varphi \leq \tilde{v} - \epsilon$ in ∂B_r , for some $\epsilon > 0$ depending on r .

Now, since $G_j \rightarrow G$ in the Hausdorff distance and $\tilde{v} \in C^\alpha(B_{1/4})$, then for every sequence $\sigma_j \rightarrow 0^+$ there is a sequence $r_j \rightarrow 0^+$, such that

$$|v_j(x) - \tilde{v}(y)| \leq \sigma_j, \quad \text{for every } x, y \in \overline{U}_j \text{ satisfying } |x - y| \leq r_j, \quad (6.36)$$

where $U_j := \{u_j > \vartheta_1 \varepsilon_j\} \cap B_{1/4}$. Since $\tilde{v} \geq \varphi$ in B_r with $\tilde{v} \geq \varphi + \epsilon$ in ∂B_r and $v(0) = \varphi(0) = 0$, we have $v_j \geq \varphi - \sigma_j$ in $\overline{U}_j \cap B_r$, $v_j \geq \varphi + \epsilon - \sigma_j$ in $\overline{U}_j \cap \partial B_r$ and $v_j \leq \sigma_j$ in $\overline{U}_j \cap B_{r_j}$, for every j . Let

$$t_j := \sup\{t \in \mathbb{R} : v_j \geq \varphi + t\sigma_j \text{ in } \overline{U}_j \cap B_r\}.$$

Since $v_j \leq \sigma_j$ in $\overline{U}_j \cap B_{r_j}$ and $\varphi > 0$ in $\{x_N > 0\} \cap \{x' = 0\} \cap B_{r_j}$, we have $t_j \in [-1, 2]$. So, setting $\tilde{\delta}_j := t_j \sigma_j \delta_j = o(\delta_j)$,

$$\phi_j(x) := x_N + \delta_j \varphi(x) + \tilde{\delta}_j,$$

and using the definition of t_j and v_j , we deduce

$$\begin{cases} u_j \geq \phi_j & \text{in } \overline{U}_j \cap B_r \\ u_j \geq \phi_j + \epsilon \delta_j & \text{in } \overline{U}_j \cap \partial B_r \\ u_j(x_j) = \phi_j(x_j) & \text{for some } x_j \in \overline{U}_j \cap B_r. \end{cases} \quad (6.37)$$

Further, by (6.20), we have

$$v_j(x) - v_j(y) \geq -\omega_j(x - y) \quad \forall x \in B_r, y \in \overline{U}_j, \quad (6.38)$$

where $\omega_j(x - y) := C(\delta_j + |x - y|)^\alpha$, and $C > 0$ and $\alpha \in (0, 1)$ are as in Lemma 6.4, for every j .

Now, given $x \in \{x_N > -\sqrt{\delta_j}\} \cap B_r$, since by assumption $\{x_N \geq -\delta_j\} \subset U_j$, we can take $y \in \overline{U}_j$ such that $|x - y| \leq 2\sqrt{\delta_j}$. Hence, using (6.38) we deduce

$$\begin{aligned} v_j(x) &\geq v_j(y) - \omega_j(x - y) \geq \varphi(y) - \sigma_j - C\delta_j^{\alpha/2} \\ &\geq \varphi(x) - C|x - y| - \sigma_j - C\delta_j^{\alpha/2} \\ &\geq \varphi(x) - 2C\delta_j^{1/2} - \sigma_j - C\delta_j^{\alpha/2}, \end{aligned} \quad (6.39)$$

for j large enough and a new constant $C > 0$. Consequently, noticing that

$$v_j(x) = \frac{u_j(x) - x_N}{\delta_j} \geq -\frac{x_N}{\delta_j} \geq \delta_j^{-1/2} \rightarrow +\infty \quad \text{in } B_r \cap \{x_N \leq -\sqrt{\delta_j}\}, \quad (6.40)$$

for large j , it follows

$$u_j \geq \phi_j \quad \text{in } B_r, \quad (6.41)$$

for j large enough, eventually taking $\tilde{\delta}_j = o(\delta_j)$ smaller.

Now, let us set $w^{\varepsilon_j} = w_{\varepsilon_j}^{\varepsilon_j}$. Combining the first inequality of (3.9) (with $\delta = 0$ and $\sigma \in (1/2, 3/4)$) with (6.41), we obtain $u_j > w^{\varepsilon_j}(\phi_j - \varepsilon_j^\sigma)$ in $B_r \cap \{w^{\varepsilon_j}(\phi_j - \varepsilon_j^\sigma) > 0\}$ and thus, since $u_j > 0$,

$$u_j > w^{\varepsilon_j}(\phi_j - \varepsilon_j^\sigma) \quad \text{in } \overline{B}_r, \quad (6.42)$$

for j large enough. Using (3.9) again and the last two inequalities in (6.37), it follows

$$\begin{cases} u_j > w^{\varepsilon_j}(\phi_j - \varepsilon_j^\sigma + \frac{\epsilon}{2}\delta_j) + \frac{\epsilon}{2}\delta_j & \text{in } \overline{U}_j \cap \partial B_r \\ u_j(x_j) < w^{\varepsilon_j}(\phi_j(x_j) + \varepsilon_j^\sigma), \end{cases} \quad (6.43)$$

for every j large enough. Now, let us set $w_\lambda := w^{\varepsilon_j}(\phi_j + \lambda \varepsilon_j^\sigma)$ and define

$$\lambda_* := \sup\{\lambda \in (-1, \infty) : w_\lambda < u_j \text{ in } B_r\}.$$

By definition of λ_* , we have

$$\begin{cases} u_j \geq w_{\lambda_*} & \text{in } B_r \\ u_j(y) = w_{\lambda_*}(y) & \text{for some } y \in \{w_{\lambda_*} > 0\} \cap \overline{B}_r, \end{cases} \quad (6.44)$$

while, following (6.15) and recalling that $\partial_N \varphi > 0$ in B_r and $\Delta \varphi = \text{tr}(M) = 0$, we easily find

$$\begin{aligned} \Delta w_{\lambda_*} &= \frac{1}{2} \Phi'_{\varepsilon_j}(w_{\lambda_*}) [1 + 2\delta_j \partial_N \varphi + \delta_j^2 |\nabla \varphi|^2] \\ &> \frac{1}{2} \Phi'_\varepsilon(w_{\lambda_*}) \quad \text{in } \{w_{\lambda_*} > 0\} \cap B_r. \end{aligned} \quad (6.45)$$

If $y \in \{w_{\lambda_*} > 0\} \cap B_r$, then $\Delta(u_j - w_{\lambda_*})(y) < 0$, in contradiction with (6.44). So, we are left to show that it cannot be $y \in \{w_{\lambda_*} > 0\} \cap \partial B_r$, obtaining a contradiction with the definition of λ_* .

To see this, we notice that $\lambda_* \in (-1, 1)$, thanks to (6.43) and the monotonicity of w^{ε_j} . Consequently, since for j large enough we have $2\delta_j^{2\sigma-1} < \frac{\varepsilon}{2}$, the first inequality in (6.43) yields

$$\begin{aligned} w_{\lambda_*} &= w^{\varepsilon_j}(\phi_j + \lambda_* \varepsilon_j^\sigma) \leq w^{\varepsilon_j}(\phi_j - \varepsilon_j^\sigma + 2\varepsilon_j^\sigma) \leq w^{\varepsilon_j}(\phi_j - \varepsilon_j^\sigma + 2\delta_j^{2\sigma}) \\ &\leq w^{\varepsilon_j}(\phi_j - \varepsilon_j^\sigma + \frac{\varepsilon}{2}\delta_j) < u_j - \frac{\varepsilon}{2}\delta_j \quad \text{in } \overline{U}_j \cap \partial B_r. \end{aligned}$$

Notice that the above inequality also implies $w_{\lambda_*} = 0$ in $\partial U_j \cap \partial B_r$, that is $\{w_{\lambda_*} > 0\} \cap \partial B_r \subset U_j \cap \partial B_r$, and our contradiction follows.

Step 3. Now we show that $\partial_N \tilde{v} \geq 0$ in $\{x_N = 0\}$ in the viscosity sense, that is for every $\varphi \in C^\infty(B_1)$ such that $\varphi \geq \tilde{v}$ in $B_{1/4}$ with equality only at some $z \in \{x_N = 0\} \cap B_{1/4}$, then $\partial_N \varphi(z) \geq 0$.

Proceeding as in *Step 2*, we assume by contradiction $\partial_N \varphi(0) < 0$ for some $\varphi \in C^\infty(B_1)$ as in (6.35) with $m < 0$ and $\text{tr}(M) = 0$.

By (6.36) and the assumptions on φ , we have $v_j \leq \varphi + \sigma_j$ in $\overline{U}_j \cap B_r$, $v_j \leq \varphi - \epsilon + \sigma_j$ in $\overline{U}_j \cap \partial B_r$ and $v_j \geq -\sigma_j$ in $\overline{U}_j \cap B_{r_j}$, for every j . So, similar to *Step 2*, we deduce

$$\begin{cases} u_j \leq \phi_j & \text{in } \overline{U}_j \cap B_r \\ u_j \leq \phi_j - \epsilon \delta_j & \text{in } \overline{U}_j \cap \partial B_r \\ u_j(x_j) = \phi_j(x_j) & \text{for some } x_j \in \overline{U}_j \cap B_r, \end{cases}$$

where $\phi_j(x) := x_N + \delta_j \varphi(x) + \tilde{\delta}_j$, for some $\tilde{\delta}_j = o(\delta_j)$. As above, by the second inequality in (3.10) (with $\sigma \in (1/2, 3/4)$ and $\delta = 0$), we obtain

$$u_j < w^{-\varepsilon_j}(\phi_j + \varepsilon_j^\sigma) \quad \text{in } \overline{U}_j \cap \overline{B}_r,$$

and

$$\begin{cases} u_j < w^{-\varepsilon_j}(\phi_j + \varepsilon_j^\sigma - \frac{\varepsilon}{2}\delta_j) - \frac{\varepsilon}{2}\delta_j & \text{in } \overline{U}_j \cap \partial B_r \\ u_j(x_j) > w_\varepsilon^{-\varepsilon}(\phi_j(x_j) - \varepsilon_j^\sigma). \end{cases} \quad (6.46)$$

Actually, we have

$$u_j < w^{-\varepsilon_j}(\phi_j + \varepsilon_j^\sigma) \quad \text{in } \overline{B}_r, \quad (6.47)$$

for j large enough. Indeed, exactly as in (6.7), u_j exponentially decays in $B_r \setminus V_j$, where

$$V_j := B_r \cap \{x : d(x, U_j) \leq \varepsilon_j^{3/4}\},$$

and thus, by (3.3), we have $u_j < w^{-\varepsilon_j}(\phi_j + \varepsilon_j^\sigma)$ in $B_r \setminus V_j$. Moreover, by monotonicity,

$$w^{-\varepsilon_j}(\phi_j + \varepsilon_j^\sigma) \geq w^{-\varepsilon_j}(\phi_j + \varepsilon_j^\sigma - \frac{\varepsilon}{2}\delta_j) \geq u_j + \frac{\varepsilon}{2}\delta_j \quad \text{in } \overline{U}_j \cap \partial B_r, \quad (6.48)$$

by the first inequality in (6.46). So, thanks to the comparison principle, we are left to check that

$$u_j < w^{-\varepsilon_j}(\phi_j + \varepsilon_j^\sigma) \quad \text{in } \partial B_r \cap (V_j \setminus U_j).$$

This follows exactly as the end of the proof of Lemma 6.2 (Step 1): by the inequality above and $\varepsilon_j \leq \delta_j^2$, we have

$$w^{-\varepsilon_j}(\phi_j + \varepsilon_j^\sigma) \geq \frac{\varepsilon}{2}\sqrt{\varepsilon_j} \quad \text{in } \overline{U}_j \cap \partial B_r,$$

and so, if $y \in \overline{B}_r$ is any point such that $w^{-\varepsilon_j}(\phi_j(y) + \varepsilon_j^\sigma) = \vartheta_2\varepsilon_j$ and $x \in \overline{U}_j \cap \partial B_r$, then it must be $|x - y| \geq c\sqrt{\varepsilon_j}$, for some $c > 0$ independent of j , which implies

$$w^{-\varepsilon_j}(\phi_j + \varepsilon_j^\sigma) \geq \vartheta_2\varepsilon_j \quad \text{in } \{x : d(x, U_j \cap \partial B_r) \leq \frac{\sqrt{\varepsilon_j}}{2}\}.$$

Finally, since $\varepsilon_j^{3/4} < \varepsilon_j^\sigma$ for every j large enough, we have

$$w^{-\varepsilon_j}(\phi_j + \varepsilon_j^\sigma) \geq \vartheta_2\varepsilon_j > \vartheta_1\varepsilon_j \geq u_j \quad \text{in } \partial B_r \cap (V_j \setminus U_j),$$

and (6.47) follows.

Now, similar to Step 2, we define $w_\lambda := w^{-\varepsilon_j}(\phi_j + \lambda\varepsilon_j^\sigma)$

$$\lambda_* := \inf\{\lambda \in (-\infty, 1) : u_j < w_\lambda \text{ in } B_r\},$$

which satisfies $\lambda_* \in (-1, 1)$ in view of the second inequality in (6.46). Further,

$$\begin{cases} u_j \leq w_{\lambda_*} & \text{in } B_r \\ u_j(y) = w_{\lambda_*}(y) & \text{for some } y \in \overline{B}_r, \end{cases} \quad (6.49)$$

and by (6.15)-(6.45), and that $\partial_N \varphi < \frac{m}{2}$ in B_r with $\text{Tr}(M) = 0$, there holds

$$\Delta w_{\lambda_*} = \frac{1}{2} \Phi'_{\varepsilon_j}(w_{\lambda_*}) [1 + 2\delta_j \partial_N \varphi + \delta_j^2 |\nabla \varphi|^2] + \delta_j w_{\lambda_*} \Delta \varphi < \frac{1}{2} \Phi'_{\varepsilon_j}(w_{\lambda_*}) \quad \text{in } B_r,$$

if j is large enough. Exactly as above, (6.49), the equation of u_j and the above differential inequality imply $y \in \partial B_r$. On the other hand, since $\sigma \in (1/2, 3/4)$, $\lambda_* \in (-1, 1)$ and using (6.48), we see that

$$w_{\lambda_*} = w^{-\varepsilon_j}(\phi_j + \varepsilon_j^\sigma - \frac{\varepsilon}{2} \delta_j + \frac{\varepsilon}{2} \delta_j - (1-\lambda) \varepsilon_j^\sigma) \geq w^{-\varepsilon_j}(\phi_j + \varepsilon_j^\sigma - \frac{\varepsilon}{2} \delta_j) \geq u_j + \frac{\varepsilon}{2} \delta_j \quad \text{in } \overline{U}_j \cap \partial B_r,$$

up to taking j larger. Further, $u_j < w_{\lambda_*}$ in $B_r \setminus V_j$ by exponential decay as $j \rightarrow +\infty$ by Lemma 3.6 (similar to (6.7)). The fact that $u_j < w_{\lambda_*}$ in $\partial B_r \cap (V_j \setminus U_j)$ follows exactly as in the proof of (6.47) (that is, the case $\lambda_* = 1$) and thus $u_j < w_{\lambda_*}$ in ∂B_r , in contradiction with $y \in \partial B_r$.

Step 4. As a consequence of *Step 2* and *Step 3*, we obtain that \tilde{v} is bounded and harmonic in $B_{1/4}$ and $\partial_N \tilde{v}|_{x_N=0} = \partial_N v|_{x_N=0} = 0$, $\tilde{v}(0) = v(0) = 0$. In particular, by standard elliptic estimates, $\tilde{v} \in C^\infty(B_\varrho)$ and

$$\sup_{x \in B_\varrho} |\tilde{v}(x) - \nabla v(0) \cdot x| \leq c_N \varrho^2,$$

every $\varrho \in (0, \frac{1}{4})$ and some $c_N > 0$. Proceeding as in (6.39), we have

$$v_j(x) \geq v_j(y) - \omega_j(x - y) \geq \tilde{v}(y) - \sigma_j - C\delta_j^{\alpha/2} \geq \tilde{v}(x) - 2C\delta_j^{1/2} - \sigma_j - C\delta_j^{\alpha/2},$$

for every $x \in \{x_N > -\sqrt{\delta_j}\} \cap B_r$, we take $y \in \overline{U}_j$ such that $|x - y| \leq 2\sqrt{\delta_j}$, while, by (6.40), $v_j(x) \geq \delta_j^{-1/2}$ in $\{x_N < -\sqrt{\delta_j}\} \cap B_r$. Consequently, by (6.36), for every $\varrho \in (0, \frac{1}{4})$, there is $j_\varrho > 0$ such that

$$\begin{aligned} v_j(x) - \nabla v(0) \cdot x &\leq c_N \varrho^2 & \text{in } B_\varrho \cap \overline{U}_j \\ -c_N \varrho^2 \leq v_j(x) - \nabla v(0) \cdot x && \text{in } B_\varrho \end{aligned} \quad (6.50)$$

for some new $c_N > 0$ and all $j \geq j_\varrho$. Now, let us define the unit vector

$$\nu := \frac{e_N + \delta_j \nabla v(0)}{|e_N + \delta_j \nabla v(0)|}.$$

Notice that, since $\partial_N v(0) = 0$, we have

$$|e_N + \delta_j \nabla \tilde{v}(0)|^2 = 1 + \delta_j^2 |\nabla \tilde{v}(0)|^2, \quad (6.51)$$

and so

$$\begin{aligned} |e_N - \nu|^2 &= \frac{\delta_j^2 |\nabla v(0)|^2 + (|e_N + \delta_j \nabla v(0)| - 1)^2}{1 + \delta_j^2 |\nabla v(0)|^2} = \frac{2\delta_j^2 |\nabla v(0)|^2 + 2(1 - |e_N + \delta_j \nabla v(0)|)}{1 + \delta_j^2 |\nabla v(0)|^2} \\ &\leq 2\delta_j^2 |\nabla \tilde{v}(0)|^2. \end{aligned}$$

Hence, recalling $\|\tilde{v}\|_{L^\infty(B_1)} = \|v\|_{L^\infty(B_1 \cap \{x_N > 0\})} \leq 1$ and using the standard gradient estimate for harmonic functions

$$|e_N - \nu| \leq \sqrt{2}\delta_j |\nabla \tilde{v}(0)| \leq \sqrt{2}\delta_j N \|\tilde{v}\|_{L^\infty(B_1)} \leq \sqrt{2}N\delta_j,$$

for and j large enough. On the other hand, since u_j is uniformly bounded in $B_{1/4}$ by (6.32), (6.51) yields

$$\begin{aligned} \frac{u_j(x) - \nu \cdot x}{\delta_j} &= \frac{u_j(x) \left(\sqrt{1 + \delta_j^2 |\nabla v(0)|^2} - 1 \right)}{\delta_j} + \frac{u_j(x) - (e_N + \delta_j \nabla v(0)) \cdot x}{\delta_j} \\ &= O(\delta_j) + v_j(x) - \nabla v(0) \cdot x, \end{aligned}$$

and thus, by (6.50),

$$\begin{aligned} u_j(x) - \nu \cdot x &\leq c_N \varrho^2 \delta_j && \text{in } B_\varrho \cap \overline{U}_j \\ -c_N \varrho^2 \delta_j &\leq u_j(x) - \nu \cdot x && \text{in } B_\varrho, \end{aligned} \quad (6.52)$$

for some new $c_N > 0$ and $j \geq j_\varrho$. Finally, given any $\gamma \in (0, 1)$ and taking $\varrho_0 \in (0, \frac{1}{4})$ such that $c_N \varrho_0^2 \leq \varrho_0^{1+\gamma}$, we obtain that both (6.33) and (6.34) are satisfied, a contradiction. \square

7. Proof of Theorem 1.4 and Corollary 1.6

The goal of this section is to prove Theorem 1.4 and Corollary 1.6. The former will be a consequence of Theorem 7.3 below, which is obtained combining Theorem 2.1 and a sliding argument in the spirit of [3,17]. The latter will be an immediate byproduct of Proposition 1.5, Theorem 1.4 and the classification of 1-homogeneous entire local minimizers of (1.5) established in [10,19].

We begin with two consequences of Theorem 2.1 that we will use in the proof of Theorem 7.3.

Corollary 7.1 (*Preservation of flatness*). *Fix $\gamma = 1/2$, and let $\delta_0 > 0$ and $\varrho_0 \in (0, 1/4)$ be the constants as in Theorem 2.1. Let $R_0 := 1/\varrho_0$. Given $\delta > 0$, we define*

$$j_\delta := \left\lceil \frac{|\log \delta^2|}{\log R_0} \right\rceil. \quad (7.1)$$

Let $u : \mathbb{R}^N \rightarrow \mathbb{R}_+$ be a critical point \mathcal{E} with $u(0) \in [\vartheta_1, \vartheta_2]$. If u satisfies $\text{Flat}_1(\nu_k, \delta, R_0^k)$ for some $\delta \in (0, \delta_0)$, $k \geq j_\delta$, and $\nu_k \in \mathbb{S}^{N-1}$, then for every i such that $j_\delta \leq i \leq k$, u satisfies $\text{Flat}_1(\nu_i, \delta, R_0^i)$ for some $\nu_i \in \mathbb{S}^{N-1}$.

Proof. The proof is by iterating Theorem 2.1. Indeed, thanks to (7.1) we have

$$\frac{1}{\delta^2 R_0^i} \leq \frac{1}{\delta^2 R_0^{j_\delta}} < 1 \quad \text{for all } i \geq j_\delta. \quad (7.2)$$

Thanks to Theorem 2.1 if u satisfies $\text{Flat}_1(\nu_i, \delta, R_0^i)$ for some $\nu_i \in \mathbb{S}^{N-1}$, and $i \geq j_\delta$ then u satisfies $\text{Flat}_1(\nu_{i-1}, R_0^\gamma \delta, R_0^{i-1})$ for some $\nu_{i-1} \in \mathbb{S}^{N-1}$. In particular u satisfies $\text{Flat}_1(\nu_{i-1}, \delta, R_0^{i-1})$. Iterating this the corollary follows. \square

Corollary 7.2 (Improvement of flatness). Fix $\gamma = 1/2$, and let $\delta_0 > 0$ and $\varrho_0 \in (0, 1/4)$ be the constants as in Theorem 2.1. Let $R_0 := 1/\varrho_0$. Let $k, n \in \mathbb{N}$ and $\delta > 0$ such that

$$(1 + 2\gamma)n \leq k - \frac{|\log \delta^2|}{\log R_0}. \quad (7.3)$$

Let $u : \mathbb{R}^N \rightarrow \mathbb{R}_+$ be a critical point of \mathcal{E} with $u(0) \in [\vartheta_1, \vartheta_2]$. If u satisfies $\text{Flat}_1(\nu_k, \delta, R_0^k)$ for some $\delta \in (0, \delta_0)$, $k \geq j_\delta$ and $\nu_k \in \mathbb{S}^{N-1}$, for every i such that $k - n \leq i \leq k$, u satisfies $\text{Flat}_1(\nu_i, R_0^{-\gamma(k-i)} \delta, R_0^i)$ for some $\nu_i \in \mathbb{S}^{N-1}$.

Proof. The proof is by iterating Theorem 2.1. Indeed, thanks to (7.3) we have

$$\frac{1}{(R_0^{-\gamma(k-i)} \delta)^2 R_0^i} \leq \frac{1}{\delta^2 R_0^{k-n-2\gamma n}} < 1 \quad \text{for all } i \geq j_\delta. \quad (7.4)$$

Thanks to Theorem 2.1 if u satisfies $\text{Flat}_1(\nu_i, R_0^{-\gamma(k-i)} \delta, R_0^i)$ for some $\nu_i \in \mathbb{S}^{N-1}$ (which is satisfied by assumption for $i = k$), and $i \geq k - n$ then u satisfies $\text{Flat}_1(\nu_{i-1}, R_0^{-\gamma(k-i+1)} \delta, R_0^{i-1})$ for some $\nu_{i-1} \in \mathbb{S}^{N-1}$. Iterating this, the corollary follows. \square

Theorem 7.3. Let $\gamma = 1/2$, and let $\varrho_0 \in (0, 1/4)$ be the constant in Theorem 2.1, and $R_0 := 1/\varrho_0 \geq 2$.

Suppose that $u : \mathbb{R}^N \rightarrow \mathbb{R}_+$ is a critical point of \mathcal{E} with $0 \in \{\vartheta_1 \leq u \leq \vartheta_2\}$ and let $\{u_\varepsilon\}_{\varepsilon \in (0,1)}$ be a blow-down family, where $u_\varepsilon := \varepsilon u(\cdot/\varepsilon)$.

Set $\varepsilon_j := R_0^{-j}$ and assume there exist $\nu \in \mathbb{S}^{N-1}$, and a sequence $j_l \rightarrow +\infty$ and $\delta_l \rightarrow 0$ (as $l \rightarrow +\infty$) for which

$$|u_{\varepsilon_{j_l}} - (\nu \cdot x)_+| \leq \delta_l \quad \text{in } B_2, \quad (7.5)$$

and

$$\{x : \nu \cdot x \leq -\delta_l\} \subset \{u_{\varepsilon_{j_l}} \leq \vartheta_1 \varepsilon_{j_l}\} \subset \{u_{\varepsilon_{j_l}} \leq \vartheta_2 \varepsilon_{j_l}\} \subset \{x : \nu \cdot x \leq \delta_l\} \quad \text{in } B_2, \quad (7.6)$$

for every $l \in \mathbb{N}$. Then u is 1D.

Proof. Throughout the proof δ_0 will denote the constant of Theorem 2.1. Observe that, by possibly replacing δ_l by some sequence with slower convergence towards 0, we may assume without loss of generality that $\vartheta_2 \varepsilon_{j_l} \leq \delta_l/2$.

Up to a rotation of the coordinate system, we may assume $\nu = e_N$. The proof is divided in several steps as follows.

Step 1. Fix $\delta \in (0, \delta_0)$ to be chosen later. We first show that

$$u \text{ satisfies } \text{Flat}_1(\nu_j, \delta, R_0^j) \quad \forall j \geq j_\delta := \left\lceil \frac{|\log(\delta^2)|}{\log R_0} \right\rceil, \quad (7.7)$$

for some $\nu_j \in \mathbb{S}^{n-1}$.

By (7.5) ($\nu = e_N$), we have

$$(x_N)_+ - \delta_l \leq u_{\varepsilon_{j_l}} \leq (x_N)_+ + \delta_l \quad \text{in } B_2. \quad (7.8)$$

Let us show that this implies

$$\begin{aligned} u_{\varepsilon_{j_l}}(x) - x_N &\leq \delta_l && \text{in } B_1 \cap \{u_{\varepsilon_{j_l}} \geq \vartheta_1 \varepsilon_{j_l}\} \\ -\delta_l &\leq u_{\varepsilon_{j_l}}(x) - x_N && \text{in } B_1, \end{aligned} \quad (7.9)$$

for all l sufficiently large.

Indeed on the one hand, (7.8) implies $u_{\varepsilon_{j_l}} \geq x_N - \delta_l$ in B_1 (for l large), which gives the inequality from below in (7.9).

To show the one from above, we set $v := u_{\varepsilon_{j_l}} - x_N - 2\delta_l$ and we show $v \leq 0$ in $B_1 \cap \{u_{\varepsilon_{j_l}} \geq \vartheta_1 \varepsilon_{j_l}\}$ using a comparison argument. Thanks to (7.8), using $(x_N)_+ - \delta_l \leq x_N$ in $\{x_N \geq -\delta_l\}$ we find (using $\vartheta_2 \varepsilon_{j_l} \leq \delta_l/2$)

$$v \leq u_{\varepsilon_{j_l}} - (x_N)_+ - \delta_l \leq \vartheta_2 \varepsilon_{j_l} - \delta_l \leq -\frac{\delta_l}{2} \quad \text{in } B_2 \cap \{u_{\varepsilon_{j_l}} \leq \vartheta_2 \varepsilon_{j_l}\} \cap \{x_N \geq -\delta_l\},$$

for every l large enough. Further, (7.8) automatically implies $v \leq -\delta_l \leq 0$ in $B_2 \cap \{x_N \geq 0\}$, since $(x_N)_+ = x_N$ there. Also, by (7.8) again, $v \leq \delta_j$ in $\overline{B_2} \cap \{u_{\varepsilon_{j_l}} \geq \vartheta_2 \varepsilon_{j_l}\} \cap \{|x_N| \leq \delta_j\}$.

On the other hand, $\Delta v = \Delta u_{\varepsilon_{j_l}} = \frac{1}{2} \Phi'_{\varepsilon_{j_l}}(u_{\varepsilon_{j_l}}) \geq 0$ in B_2 and thus the function

$$\underline{v} := \frac{v}{\delta_l} + A x_N^2$$

satisfies

$$\begin{cases} \Delta \underline{v} \geq 2A & \text{in } B_2 \cap \{-\delta_l \leq x_N < 0\} \\ \underline{v} \leq -\frac{1}{2} + A\delta_l^2 & \text{in } B_2 \cap \{x_N = -\delta_l\} \\ \underline{v} \leq 0 & \text{in } B_2 \cap \{x_N = 0\} \\ \underline{v} \leq 1 + A\delta_l^2 & \text{in } \partial B_2 \cap \{-\delta_l \leq x_N < 0\}, \end{cases}$$

for every $A > 0$. Now, consider the function $\bar{h}_{x_0}(x) = \frac{A}{N}|x - x_0|^2$. For every $x_0 \in B_1 \cap \{-\delta_l < x_N < 0\}$, we have

$$\bar{h}_{x_0} \geq \frac{A}{N} \quad \text{in } \partial B_2 \cap \{-\delta_l \leq x_N \leq 0\},$$

and taking $A := 2N$, we have $\bar{h}_{x_0} \geq 2$ in $\partial B_2 \cap \{-\delta_l \leq x_N \leq 0\}$. Then, for l large, we have $A\delta_l^2 \leq 1$ and

$$\begin{cases} \Delta \underline{v} \geq 2A = \Delta \bar{h}_{x_0} & \text{in } B_2 \cap \{-\delta_l \leq x_N < 0\} \\ \underline{v} \leq 0 \leq \bar{h}_{x_0} & \text{in } B_2 \cap \partial\{-\delta_l \leq x_N < 0\} \\ \underline{v} \leq 2 \leq \bar{h}_{x_0} & \text{in } \partial B_2 \cap \{-\delta_l \leq x_N < 0\}. \end{cases}$$

Then, by the maximum principle we obtain $\underline{v} \leq \bar{h}_{x_0}$. Consequently, since $\bar{h}_{x_0}(x_0) = 0$ and x_0 is arbitrary in $B_1 \cap \{-\delta_l < x_N < 0\}$, we have $\bar{v} \leq 0$ in $B_1 \cap \{-\delta_l < x_N < 0\}$ and so, by the definition of \underline{v} , we obtain $v \leq 0$ in $B_1 \cap \{-\delta_l < x_N < 0\}$. This proves (7.9). In other words, after scaling we have shown that (7.7) holds for $j = j_l$ and $\nu_{j_l} = \nu$, provided that l is sufficiently large. Hence, as a consequence of Corollary 7.1 we obtain that (7.7) holds for every integer j such that $j_\delta \leq j \leq j_l$ for some $\nu_j \in \mathbb{S}^{N-1}$. Observing that j_l can be taken arbitrarily large concludes the proof of (7.7).

Step 2. In this second step, we prove that there exists $C \geq 1$ such that for every $z \in \{\vartheta_1 \leq u \leq \vartheta_2\}$ and every $R \geq C$

$$u(z + \cdot) \quad \text{satisfies} \quad \text{Flat}_1(e_N, CR^{-1/2}, R) \quad \forall R \geq C. \quad (7.10)$$

Note that this is a really strong information since the constant C and the direction e_N of flatness are independent of z , which varies in an unbounded set!

To obtain (7.10), we first show the existence of some k_0 (independent of z) such that for every $k \geq k_0$ and every $z \in \{\vartheta_1 \leq u \leq \vartheta_2\}$, there are $\nu_{z,k} \in \mathbb{S}^{N-1}$ such that, for all $k \geq k_0$,

$$u(z + \cdot) \quad \text{satisfies} \quad \text{Flat}_1(\nu_{z,k}, \delta_0, R_0^k) \quad (7.11)$$

for some $\nu_{z,k} \in \mathbb{S}^{N-1}$. Indeed, given $z \in \{\vartheta_1 \leq u \leq \vartheta_2\}$ choose $i \in \mathbb{N}$ such that $|z| \leq \frac{\delta_0}{2} R_0^i$. Take $j = i + 1$ in (7.7), and choose δ such that $2\delta R_0 \leq \delta_0$. We then have

$$\begin{aligned} u(x) - \nu_{i+1} \cdot x &\leq \frac{\delta_0}{2} R_0^i && \text{in } B_{R_0^{i+1}}(0) \cap \{u \geq \vartheta_1\} \\ -\frac{\delta_0}{2} R_0^i &\leq u(x) - \nu_{i+1} \cdot x && \text{in } B_{R_0^{i+1}}(0). \end{aligned} \quad (7.12)$$

Now since $|z| \leq \frac{\delta_0}{2} R_0^i$ and $R_0 \geq 2$ we have $B_{R_0^i}(z) \subset B_{R_0^{i+1}}$ and

$$|u(x) - \nu_{i+1} \cdot (x - z)| \leq |u(x) - \nu_{i+1} \cdot x| + |z| \leq \delta_0 R_0^i \quad \text{in } B_{R_0^i}(z).$$

Thus, (7.12) implies

$$\begin{aligned} u(x) - \nu_{i+1} \cdot x &\leq \delta_0 R_0^i && \text{in } B_{R_0^i}(z) \cap \{u \geq \vartheta_1\} \\ -\delta_0 R_0^i &\leq u(x) - \nu_{i+1} \cdot x && \text{in } B_{R_0^i}(z). \end{aligned}$$

In other words, setting $\nu_{k,z} := \nu_{i+1}$, we see that (7.11) is satisfied for $k = i$ large enough (where i depends on z). But then thanks to Corollary 7.1 (applied with $\delta = \delta_0$ and to the “translated function” $u(z + \cdot)$) we obtain that (7.11) holds for all $k \geq k_0 := j_{\delta_0}$.

We will now use (7.11) and Corollary 7.2 (applied again to the translated function $u(z + \cdot)$) to show (7.10). Indeed, for given $j \in \mathbb{N}$ large enough, set

$$n := \left\lceil \frac{j - |\log \delta^2| / \log R_0}{2\gamma} \right\rceil$$

and

$$k := j + n$$

Then,

$$(1 + 2\gamma)n = n + 2\gamma \left\lceil \frac{j - |\log \delta^2| / \log R_0}{2\gamma} \right\rceil \leq k - \frac{|\log \delta_0^2|}{\log R_0}. \quad (7.13)$$

The above inequality implies that (7.3) is satisfied. By (7.11) (since we assume that $j \geq C$ sufficiently large we have $k \geq j \geq k_0$), we may apply Corollary 7.2 to $u(z + \cdot)$ to obtain that $u(z + \cdot)$ satisfies $\text{Flat}_1(\nu_{z,i}, R_0^{-\gamma(k-i)} \delta_0, R_0^i)$ for some $\nu_{z,i} \in \mathbb{S}^{N-1}$ for all $i = j, j+1, \dots, j+n$ (in particular for $i = j$). Hence using the definition of Flat_1 and that $k - i = n \geq \frac{j}{2\gamma} - C$, we obtain

$$\delta_0 R_0^{-\gamma(k-j)} R_0^j \leq \delta_0 R_0^{j-\gamma n} \leq C R_0^{j(1-1/2)}$$

and

$$\begin{aligned} u(x) - \nu_{z,i} \cdot (x - z) &\leq C R_0^{j/2} && \text{in } B_{R_0^j}(z) \cap \{u \geq \vartheta_1\} \\ -C R_0^{j/2} &\leq u(x) - \nu_{z,i} \cdot (x - z) && \text{in } B_{R_0^j}(z). \end{aligned} \quad (7.14)$$

Now, on the one hand, as a consequence of (7.14), we have

$$\max(0, \nu_{z,j} \cdot x - C R_0^{j/2}) \leq u(z + x) \leq \max(\vartheta_1, \nu_{z,j} \cdot x + C R_0^{j/2}) \quad \text{in } B_{R_0^j},$$

and thus, using this in two consecutive scales, we obtain

$$\max(0, \nu_{z,j} \cdot x - C R_0^{j/2}) \leq \max(\vartheta_1, \nu_{z,j+1} \cdot x + C R_0^{(j+1)/2}) \quad \text{in } B_{R_0^j}.$$

This implies (for j large)

$$|\nu_{z,j} - \nu_{z,j+1}| \leq C(R_0)R_0^{-j/2}, \quad (7.15)$$

where $C(R_0)$ is independent of z and j . This shows that $\nu_{z,j} \rightarrow \nu_z$ as $j \rightarrow \infty$ for all z . On the other hand, since for every two pair of points z_1, z_2 (7.14) applied at a scales $R_0^j \gg |z_1 - z_2|$ implies $(\nu_{z_1,j} - \nu_{z_2,j}) \rightarrow 0$, we see that $\nu_z \equiv \nu_*$ for all z , where ν_* is independent of z . On the other hand, assumption (7.5) (where $\nu = e_N$ as said in the beginning of the proof), forces $\nu_* = e_N$ and hence $\lim_{j \rightarrow \infty} \nu_{z,j} = e_N$ for all z . Finally, using again (7.15), triangle inequality, and summing the geometric series we obtain

$$|\nu_{z,j} - e_N| \leq |\nu_{z,j} - \lim_{j \rightarrow \infty} \nu_{z,j}| \leq C(R_0) \sum_{l=j}^{\infty} R_0^{-l/2} \leq CR_0^{-j/2},$$

for all $z \in \{\vartheta_1 \leq u \leq \vartheta_2\}$. Combining this information with (7.14) we conclude the proof of (7.10).

Step 3. We now observe that (7.10) has two significant consequences. First, it implies the existence of a function $G : \mathbb{R}^{N-1} \rightarrow \mathbb{R}$ with $G(0) = 0$ satisfying

$$|G(x') - G(y')| \leq C\sqrt{|x' - y'|}, \quad \forall x', y' \in \mathbb{R}^{N-1} \quad (7.16)$$

and

$$\{x_N \leq G(x') - C\} \subset \{u \leq \vartheta_1\} \subset \{u \leq \vartheta_2\} \subset \{x_N \leq G(x') + C\} \quad \text{in } \mathbb{R}^N. \quad (7.17)$$

Second, since $u - x_N$ is harmonic in $\{u > \vartheta_2\}$, standard elliptic estimates yield

$$\sup_{x \in B_{r/2}(y)} |\nabla(u(x) - x_N)| \leq \frac{c_N}{r} \sup_{x \in B_r(y)} |u(x) - x_N| \leq \frac{c_N C}{r} R^{j/2},$$

for every $B_r(y) \subset \{u > \vartheta_2\} \cap B_{R^j}$. Consequently, for every $j \in \mathbb{N}$ and every y such that $B_{R^j/2}(y) \subset \{u > \vartheta_2\} \cap B_{R^j}$, we have

$$\sup_{x \in B_{R^j/4}(y)} |\nabla(u(x) - x_N)| \leq c_N C R^{-j/2}. \quad (7.18)$$

This easily implies that

$$|\nabla u| \leq C \quad \text{in } \mathbb{R}^N \quad (7.19)$$

Indeed, if $x = (x', x_N)$ is a point in \mathbb{R}^N and let $R_0 \geq C$ to be chosen. We consider two complementary cases: either $x_N - G(x') \leq R_0$ or $x_N - G(x') > R_0$. In the first case, by (7.10) with $R = 2R_0$, we obtain $|u| \leq C$ in $B_{R_0}(x)$ (with a possibly larger C).

Now, since u is a bounded solution of the semilinear equation $\Delta u = \frac{1}{2}\Phi'(u)$ standard elliptic estimates yield $|\nabla u(x)| \leq C$. In the second case, using (7.17)–(7.16) we obtain that, if R_\circ is chosen large enough and $R := \text{dist}(x, \partial\{u > \vartheta_2\}) \geq R_\circ$, then it also follows $|\nabla u(x)| \leq C$, thanks to (7.18).

Step 4. We now perform a sliding argument à la Caffarelli-Berestycki-Nirenberg. We fix $\sigma > 0$, $e' \in \mathbb{S}^{N-1} \cap \{x_N = 0\}$, and define, for any given $\lambda > 0$,

$$e := (e', \sigma), \quad u^\lambda(x) := u(x - \lambda e). \quad (7.20)$$

Choose $\lambda_\sigma > 0$ such that $C\sqrt{\lambda_\sigma} + 2C \leq \sigma\lambda_\sigma$, where C is the constant in (7.17)–(7.18). Let us show that

$$u^\lambda \leq u \quad \text{in } \mathbb{R}^N \quad \text{for every } \lambda \geq \lambda_\sigma. \quad (7.21)$$

To prove so, we first observe that, for every $\lambda \geq \lambda_\sigma$

$$\{u \leq \vartheta_2\} \subset \{u^\lambda \leq \vartheta_1\}. \quad (7.22)$$

Indeed, let $x \in \{u \leq \vartheta_2\}$ and notice that (7.16) yields

$$\begin{aligned} (x - \lambda e)_N - G((x - \lambda e)') + C &= x_N - \sigma\lambda - G(x' - \lambda e') + C \\ &\leq -\sigma\lambda + G(x') - G(x' - \lambda e') + 2C \\ &\leq -\sigma\lambda + C\sqrt{\lambda} + 2C \leq 0, \end{aligned}$$

for every $\lambda \geq \lambda_\sigma$, provided λ_σ is chosen large enough.

Now, we set $v := u - u^\lambda$ and we show that $v \geq 0$ in \mathbb{R}^N for every $\lambda \geq \lambda_\sigma$, that is (7.22).

To do so, we first notice that $\mathbb{R}^N = \Omega_1 \cup \Omega_2 := \{u \geq \vartheta_2\} \cup \{u^\lambda < \vartheta_1\}$ for every $\lambda \geq \lambda_\sigma$, thanks to (7.22). Further, in the domain Ω_1 the function v satisfies $\Delta v = 0$ in $\{u^\lambda > \vartheta_2\}$ and $u - u^\lambda \geq \vartheta_2 - \vartheta_2 = 0$ in $u^\lambda \leq \vartheta_2$. Hence the negative part of v_- is subharmonic in Ω_1 .

Also, thanks to (7.22), the boundary $\partial\{u > \vartheta_2\}$ of Ω_1 is contained in $\{u^\lambda \leq \vartheta_1\}$ and hence $u - u^\lambda \geq \vartheta_2 - \vartheta_1 > 0$ on $\partial\{u > \vartheta_2\}$. In other words v_- is subharmonic and vanishes on the boundary of Ω_1 . Since —thanks to (7.16) and (7.17)— the complement of Ω_1 contains a cone with nonempty interior, and —thanks to (7.19) v (and in particular v_-) is bounded in all of \mathbb{R}^N , we deduce $v_- = 0$ in Ω_1 from the comparison principle in unbounded domains which contain a cone (see for instance [3, Lemma 2.1]).

Similarly inside Ω_2 , either $u \geq \vartheta_1$ and $u^\lambda < \vartheta_1$ and so $v \geq 0$ or both u and u^λ are smaller than ϑ_1 . In that second case, recalling that Φ' is increasing in $(0, \vartheta_1)$, we have $\Delta v = \Phi'(u) - \Phi'(u^\lambda) \leq 0$ at points where $v = u - u^\lambda \leq 0$. Hence v_- is again a subharmonic function in Ω_2 . Similarly as before we can show that $v_- = 0$ on $\partial\Omega_2$ and

that it is bounded. And again the complement of Ω_2 contains a cone, so we may conclude $v_- = 0$ everywhere (that is $v \geq 0$).

Step 5. Let C and G as in (7.17). Define

$$\overline{C} := C + (1 + \lambda_\sigma) \|\nabla G\|_{L^\infty(\mathbb{R}^{N-1})}, \quad (7.23)$$

and

$$\mathcal{G} := \{x = (x', x_N) \in \mathbb{R}^N : |x_N - G(x')| \leq \overline{C}\}. \quad (7.24)$$

We prove that for every $\lambda > 0$

$$u^\lambda \leq u \text{ in } \mathcal{G} \quad \Rightarrow \quad u^\lambda \leq u \text{ in } \mathbb{R}^N. \quad (7.25)$$

In light of (7.21), it is enough to treat the case $\lambda \in (0, \lambda_\sigma)$. Following the ideas of *Step 4*, we observe that

$$\{x_n \leq G(x') - \bar{C}\} \subset \{u \leq \vartheta_1\}$$

Indeed, let x satisfy

$$x_N \leq G(x') - \bar{C}.$$

Consequently, by (7.23), the above inequality and the definition of e , we obtain

$$\begin{aligned} (x - \lambda e)_N - G((x - \lambda e)') &\leq x_N - G(x') + \lambda \|\nabla G\|_{L^\infty(\mathbb{R}^{N-1})} \\ &\leq x_N - G(x') + \lambda_\sigma \|\nabla G\|_{L^\infty(\mathbb{R}^{N-1})} \leq -\bar{C} + \overline{C} - C = -C, \end{aligned}$$

and thus, by (7.17), we have $x - \lambda e \in \{u \leq \vartheta_1\}$. In a very similar way, we show

$$\{u^\lambda \leq \vartheta_2\} \subset \{x_N \leq G(x') + \overline{C}\}, \quad (7.26)$$

for every $\lambda \in (0, \lambda_\sigma)$. To complete the proof of (7.25), it is enough to consider $v = u - u^\lambda$, notice that $v \geq 0$ in $\partial\mathcal{G}$ (by assumption) and repeat the arguments of *Step 5*.

Step 6. In this step we show that for every $\lambda > 0$

$$u^\lambda \leq u \quad \text{in } \mathcal{G}. \quad (7.27)$$

Notice that, as a consequence of (7.25), (7.27) implies that for every $\lambda > 0$

$$u^\lambda \leq u \quad \text{in } \mathbb{R}^N. \quad (7.28)$$

To verify (7.27), we let

$$\lambda_* := \inf\{\lambda \geq 0 : u^\lambda \leq u \text{ in } \mathcal{G}\} \leq \lambda_\sigma,$$

and show that $\lambda_* = 0$. Assume by contradiction that $\lambda_* \in (0, \lambda_\sigma)$.

By definition of λ_* , we have $u^{\lambda_*} \leq u$ in \mathcal{G} and, in addition, there exists $x_j \in \mathcal{G}$ such that $u(x_j) - u^{\lambda_*}(x_j) \leq 1/j$. Set $u_j(x) := u(x + x_j)$, $u_j^{\lambda_*}(x) := u^{\lambda_*}(x + x_j)$, $v_j := u_j - u_j^{\lambda_*}$ and $\mathcal{G}_j := \mathcal{G} - x_j$.

We have

$$\mathcal{G}_j = \{(x', x_N) \in \mathbb{R}^N : |x_N - G_j(x')| \leq \overline{C}\},$$

where $G_j(x') := G(x' + x'_j) - x_{j,N}$ (here $x_{j,N} := (x_j)_N$) with

$$\begin{aligned} |G_j(x')| &\leq |G(x' + x'_j) - G(x'_j)| + |G(x'_j) - x_{j,N}| \leq C\sqrt{|x'|} + 2\overline{C} \\ \|\nabla G_j\|_{L^\infty(\mathbb{R}^{N-1})} &\leq \|\nabla G\|_{L^\infty(\mathbb{R}^{N-1})}, \end{aligned} \quad (7.29)$$

for every j in view of (7.16) and that $x_j \in \mathcal{G}$. As a consequence, we deduce the existence of a locally bounded function $\overline{G} : \mathbb{R}^N \rightarrow \mathbb{R}$ such that $G_j \rightarrow \overline{G}$ locally uniformly in \mathbb{R}^N and $\mathcal{G}_j \rightarrow \overline{\mathcal{G}}$ locally Hausdorff in \mathbb{R}^N (up to subsequence), where $\overline{\mathcal{G}} := \{(x', x_N) \in \mathbb{R}^N : |x_N - \overline{G}(x')| \leq \overline{C}\}$.

In particular, thanks to (7.17), we have

$$\{x_N \leq \overline{G}(x') - 2\overline{C}\} \subset \{u_j \leq \vartheta_1\} \subset \{u_j \leq \vartheta_2\} \subset \{x_N \leq \overline{G}(x') + 2\overline{C}\} \quad \text{in } \mathbb{R}^N, \quad (7.30)$$

for every j large enough. On the other hand, using (7.29) for $x' = 0$ and recalling (7.19) we have

$$|u_j(0)| \leq 2\overline{C} \quad \text{and} \quad |\nabla u_j| \leq C \quad \text{in } \mathbb{R}^N,$$

thus, the sequence $\{u_j\}_{j \in \mathbb{N}}$ is locally uniformly bounded in \mathbb{R}^N .

Further, since $\Delta u_j = \frac{1}{2}\Phi'(u_j)$ in \mathbb{R}^N and Φ' is bounded, standard elliptic estimates and a diagonal argument yield $u_j \rightarrow \overline{u}$ in C_{loc}^2 as $j \rightarrow +\infty$, for some $\overline{u} \in C_{loc}^2(\mathbb{R}^N)$, up to passing to a subsequence. Similar, $\overline{u}_j \rightarrow \overline{u}^{\lambda_*}$ in C_{loc}^2 as $j \rightarrow +\infty$ and, since

$$\begin{cases} \Delta v_j = \frac{1}{2}(\Phi'(u_j) - \Phi'(u_j^{\lambda_*})) & \text{in } \mathbb{R}^N \\ v_j \geq 0 & \text{in } \mathcal{G}_j \\ v_j(0) \leq 1/j, \end{cases}$$

for every $j \in \mathbb{N}$, $v_j \rightarrow \overline{v}$ in C_{loc}^2 as $j \rightarrow +\infty$. By uniform convergence we have $\overline{v}(0) = 0$, $\Delta \overline{u} = \frac{1}{2}\Phi'(\overline{u})$ in \mathbb{R}^N , and $\overline{v} \geq 0$ in $\overline{\mathcal{G}}$. Therefore, using (7.25) applied to the function \overline{u} and with $\lambda = \lambda_*$, we deduce

$$\overline{v} \geq 0 \quad \text{in } \mathbb{R}^N. \quad (7.31)$$

On the other hand,

$$\Delta \bar{v} = \frac{1}{2}(\Phi'(\bar{u}) - \Phi'(\bar{u}^{\lambda_*})) \leq \|\Phi\|_{C^{1,1}(\mathbb{R})} \bar{v} \quad \text{in } \mathbb{R}^N,$$

and so, $\bar{v}(0) = 0$, (7.31) and the strong maximum principle yield $\bar{v} = 0$ in \mathbb{R}^N . Consequently, for every fixed $x \in \mathbb{R}^N$

$$\begin{aligned} \bar{u}(x) &= \bar{u}^{\lambda_*}(x) = \lim_{j \rightarrow +\infty} u^{\lambda_*}(x + x_j) = \lim_{j \rightarrow +\infty} u(x + x_j - e\lambda_*) \\ &= \lim_{j \rightarrow +\infty} u_j(x - e\lambda_*) = \bar{u}(x - e\lambda_*), \end{aligned}$$

that is, \bar{u} is λ_* -periodic along the direction e .

Now, fix $\vartheta \in (\vartheta_1, \vartheta_2)$ and take $\tilde{x}_j \in \mathcal{G}$ such that $\tilde{x}_{j,N} = x_{j,N}$ and $u(\tilde{x}_j) = \vartheta$, for every $j \in \mathbb{N}$ and set $\hat{x}_j := \tilde{x}_j - x_j$. By (7.17), (7.23) and (7.24), we have $|\hat{x}_j| \leq 2\bar{C}$ and thus, up to passing to a subsequence, $\hat{x}_j \rightarrow \hat{x} \in \mathcal{G}$ as $j \rightarrow +\infty$, and

$$\vartheta = \lim_{j \rightarrow +\infty} u(\tilde{x}_j) = \lim_{j \rightarrow +\infty} u(\hat{x}_j + x_j) = \lim_{j \rightarrow +\infty} u_j(\hat{x}_j) = \bar{u}(\hat{x}), \quad (7.32)$$

by uniform convergence. We also have

$$\{\bar{u} = \vartheta\} \subset \{x_N \geq -\bar{C}(1 + \sqrt{|x'|})\}, \quad (7.33)$$

up to taking $\bar{C} > 0$ larger. To see this, we take $y \in \{\bar{u} = \vartheta\}$ and we notice that $y \in \{\vartheta_1 \leq u_j \leq \vartheta_2\}$ for large j 's or, equivalently, $y + x_j \in \{\vartheta_1 \leq u \leq \vartheta_2\} \subset \{x_N \geq G(x') - C\}$, in light of (7.17). This, combined with the fact that $x_{j,N} \leq G(x'_j) + \bar{C}$ (since $x_j \in \mathcal{G}$) and (7.16) give

$$y_N \geq G(y' + x'_j) - x_{j,N} - C \geq G(y' + x'_j) - G(x'_j) - \bar{C} - C \geq -c\sqrt{|y'|} - \bar{C} - C,$$

which is (7.33), up to taking $c > 0$ large enough (depending on C and \bar{C}).

To complete the contradiction argument, we notice that by (7.32) and the λ_* -periodicity of \bar{u} (along the direction e), it must be $\hat{x} - ne\lambda_* \in \{\bar{u} = \vartheta\}$ for every $n \in \mathbb{Z}$, and thus, using (7.33), it follows $\hat{x} - ne\lambda_* \in \{x_N \geq -\bar{C}(1 + \sqrt{|x'|})\}$ for every $n \in \mathbb{Z}$. Using the definition of e and passing to the limit as $n \rightarrow +\infty$, we find

$$\begin{aligned} 0 &\leq (\hat{x} - ne\lambda_*)_N + \bar{C}(1 + \sqrt{|(\hat{x} - ne\lambda_*)'|}) \\ &= \hat{x}_N - n\sigma\lambda_* + \bar{C}(1 + \sqrt{|\hat{x}' - ne'\lambda_*|}) \rightarrow -\infty, \end{aligned}$$

as $n \rightarrow +\infty$, a contradiction, and (7.27) follows.

Step 7. By (7.28), we have $\partial_e u \geq 0$ in \mathbb{R}^N , independently of $\sigma > 0$ (cf. (7.20)) and so $\partial_{(e',0)} u \geq 0$ in \mathbb{R}^N , for every $e' \in \mathbb{S}^{N-1} \cap \{x_N = 0\}$. Since $\partial_{(-e',0)} u = -\partial_{(e',0)} u \leq 0$ in \mathbb{R}^N and e' is arbitrary, it must be $\partial_{(e',0)} u = 0$ in \mathbb{R}^N for every $e' \in \mathbb{S}^{N-1} \cap \{x_N = 0\}$, that is u is 1D. \square

Proof of Theorem 1.4. Let $u : \mathbb{R}^N \rightarrow \mathbb{R}_+$ be a critical point of \mathcal{E} in \mathbb{R}^N satisfying (1.9) and (1.10) for some $R_k \rightarrow \infty$ and $\delta_k \rightarrow 0$. Setting $\varepsilon_k := R^{-k}$ and scaling, we immediately see that $u_{\varepsilon_k} := \varepsilon_k u(\cdot/\varepsilon_k)$ satisfies (7.5) and (7.6) for k large and so $u(x) = v(x_N)$ for some $v : \mathbb{R} \rightarrow \mathbb{R}$ (up to a rotation), by Theorem 7.3.

On the other hand, by Lemma 3.1, we know there are exactly three families of 1D solutions (cf. (i), (ii), (iii) of Lemma 3.1 with $\varepsilon = 1$). However, by (7.5), we have $u_{\varepsilon_k} \rightarrow (x_N)_+$ locally uniformly, up to a translation and a rotation, and thus v cannot be of class (ii) and (iii). The only possibility is that v is of class (i). Recalling that $v(0) = \vartheta_1$ by construction, a direct integration of (3.5) (with $A = 1$) yields (cf. (3.6))

$$v^{-1}(z) = \int_{\vartheta_1}^z \frac{d\zeta}{\sqrt{\Phi(\zeta)}},$$

for every $z \in \mathbb{R}$, which is (1.7), up to a shift. \square

Proof of Corollary 1.6. Let $u : \mathbb{R}^N \rightarrow \mathbb{R}_+$ be an entire local minimizer of \mathcal{E} in \mathbb{R}^N . Up to shift, we may assume $u(0) = \vartheta_1$. If $\{R_j\}_{j \in \mathbb{N}}$ is an arbitrary sequence satisfying $R_j \rightarrow +\infty$ as $j \rightarrow +\infty$ then, by Proposition 1.5, there exist sequences $R_{j_\ell} \rightarrow +\infty$, $\delta_\ell \rightarrow 0$ and a 1-homogeneous nontrivial entire local minimizer u_0 of (1.5) with $0 \in \partial\{u_0 > 0\}$, such that (1.11) and (1.12) hold true (with $k = j$). Consequently, since we know that

$$u_0(x) = (\nu \cdot x)_+,$$

for some $\nu \in \mathbb{S}^{N-1}$ (see [10,19]), we deduce that (1.9) and (1.10) are satisfied too, and thus u satisfies (1.7) by Theorem 1.4. \square

References

- [1] H.W. Alt, L. Caffarelli, Existence and regularity for a minimum problem with free boundary, *J. Reine Angew. Math.* 325 (1981) 105–144.
- [2] H.W. Alt, L. Caffarelli, A. Friedman, Variational problems with two phases and their free boundaries, *Trans. Am. Math. Soc.* 282 (1984) 431–461.
- [3] H. Berestycki, L. Caffarelli, L. Nirenberg, Monotonicity for elliptic equations in unbounded Lipschitz domains, *Commun. Pure Appl. Math.* 50 (1997) 1089–1111.
- [4] J.D. Buckmaster, G.S. Ludford, *Theory of Laminar Flames*, Cambridge Univ. Press, Cambridge, 1982.
- [5] X. Cabré, A. Figalli, X. Ros-Oton, J. Serra, Stable solutions to semilinear elliptic equations are smooth up to dimension 9, *Acta Math.* 224 (2020) 187–252.
- [6] L. Caffarelli, A Harnack inequality approach to the regularity of free boundaries. I. Lipschitz free boundaries are $C^{1,\alpha}$, *Rev. Mat. Iberoam.* 3 (1987) 139–162.
- [7] L. Caffarelli, A Harnack inequality approach to the regularity of free boundaries. II. Flat free boundaries are Lipschitz, *Commun. Pure Appl. Math.* 42 (1989) 55–78.
- [8] L. Caffarelli, A Harnack inequality approach to the regularity of free boundaries. III. Existence theory, compactness, and dependence on X , *Ann. Sc. Norm. Super. Pisa* 15 (1988) 583–602.
- [9] L.A. Caffarelli, X. Cabré, *Fully Nonlinear Elliptic Equations*, Colloquium Publications, vol. 43, AMS, 1995.

- [10] L.A. Caffarelli, D.S. Jerison, C.E. Kenig, Global energy minimizers for free boundary problems and full regularity in three dimensions, in: *Contemp. Math.*, vol. 350, Amer. Math. Soc., Providence RI, 2004, pp. 83–97.
- [11] L.A. Caffarelli, S. Salsa, *A Geometric Approach to Free Boundary Problems*, *Grad. Stud. Math.*, vol. 68, AMS, 2005.
- [12] L.A. Caffarelli, J.L. Vázquez, A free-boundary problem for the heat equation arising in flame propagation, *Trans. Am. Math. Soc.* 347 (1995) 411–441.
- [13] D. Danielli, A. Petrosyan, H. Shahgholian, A singular perturbation problem for the p -Laplace operator, *Indiana Univ. Math. J.* 52 (2003) 457–476.
- [14] E. De Giorgi, Convergence problems for functionals and operators, in: *Proceedings of the International Meeting on Recent Methods in Nonlinear Analysis*, Rome, 1978, Pitagora, Bologna, 1979, pp. 131–188.
- [15] D. De Silva, Free boundary regularity from a problem with right hand side, *Interfaces Free Bound.* 13 (2011) 223–238.
- [16] D. De Silva, D.S. Jerison, A singular energy minimizing free boundary, *J. Reine Angew. Math.* 635 (2009) 1–21.
- [17] S. Dipierro, J. Serra, E. Valdinoci, Improvement of flatness for nonlocal phase transitions, *Am. J. Math.* 142 (2020) 1083–1160.
- [18] X. Fernández-Real, X. Ros-Oton, On global solutions to semilinear elliptic equations related to the one-phase free boundary problem, *Discrete Contin. Dyn. Syst., Ser. A* 39 (2019) 6945–6959.
- [19] D.S. Jerison, O. Savin, Some remarks on stability of cones for the one phase free boundary problem, *Geom. Funct. Anal.* 25 (2015) 1240–1257.
- [20] D. Kinderlehrer, L. Nirenberg, J. Spruck, Regularity in elliptic free boundary problems, *J. Anal. Math.* 34 (1979) 86–119.
- [21] A. Petrosyan, N.K. Yip, Nonuniqueness in a free boundary problem from combustion, *J. Geom. Anal.* 18 (2007) 1098–1126.
- [22] O. Savin, Regularity of flat level sets in phase transitions, *Ann. Math.* 169 (2009) 41–78.
- [23] B. Velichkov, Regularity of the one-phase free boundaries, <https://cvgmt.sns.it/paper/4367/>, 2019.
- [24] G.S. Weiss, Partial regularity for weak solutions of an elliptic free boundary problem, *Commun. Partial Differ. Equ.* 23 (1998) 439–455.
- [25] G.S. Weiss, Partial regularity for a minimum problem with free boundary, *J. Geom. Anal.* 9 (2) (1999) 317–326.
- [26] G.S. Weiss, A homogeneity improvement approach to the obstacle problem, *Invent. Math.* 138 (1) (1999) 23–50.
- [27] G.S. Weiss, A singular limit arising in combustion theory: fine properties of the free boundary, *Calc. Var. Partial Differ. Equ.* 17 (2003) 311–340.