

A hybrid modeling environment to describe aggregates of cells heterogeneous for genotype and behavior with possible phenotypic transitions

Original

A hybrid modeling environment to describe aggregates of cells heterogeneous for genotype and behavior with possible phenotypic transitions / Chiari, Giulia; Delitala, Marcello Edoardo; Morselli, David; Scianna, Marco. - In: INTERNATIONAL JOURNAL OF NON-LINEAR MECHANICS. - ISSN 0020-7462. - ELETTRONICO. - 144:(2022), p. 104063. [10.1016/j.ijnonlinmec.2022.104063]

Availability:

This version is available at: 11583/2964765 since: 2022-05-26T19:07:03Z

Publisher:

Elsevier

Published

DOI:10.1016/j.ijnonlinmec.2022.104063

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

Elsevier postprint/Author's Accepted Manuscript

© 2022. This manuscript version is made available under the CC-BY-NC-ND 4.0 license
<http://creativecommons.org/licenses/by-nc-nd/4.0/>. The final authenticated version is available online at:
<http://dx.doi.org/10.1016/j.ijnonlinmec.2022.104063>

(Article begins on next page)

Evaluation of the accuracy of convection-permitting sub-daily extreme precipitation simulations over Italy

Paola Mazzoglio¹, Marco Borga², Marco Lompi³, Francesco Marra^{4,5}, Eleonora Dallan², Roberto Deidda⁶, Pierluigi Claps¹, Salvatore Manfreda⁷, Valerio Noto⁸, Alberto Viglione¹, Mario Raffa⁹, Enrica Caporali³

- (1) Department of Environment, Land and Infrastructure Engineering, Politecnico di Torino, paola.mazzoglio@polito.it
- (2) Department of Land, Environment, Agriculture and Forestry, University of Padova
- (3) Department of Civil and Environmental Engineering, University of Florence
- (4) Department of Geosciences, University of Padova
- (5) Institute of Atmospheric Sciences and Climate, National Research Council of Italy
- (6) Department of Civil, Environmental and Architecture Engineering, University of Cagliari
- (7) Department of Civil, Building and Environmental Engineering, University of Naples Federico II
- (8) Department of Engineering, University of Palermo
- (9) Regional Model and geo-Hydrological Impacts division, CMCC Foundation

Convection-permitting models have the potential to capture crucial processes in the climate system, presenting an opportunity to significantly enhance climate projections by providing more accurate representations of precipitation extremes.

In this work, we conduct an evaluation of the accuracy of sub-daily precipitation extremes obtained from VHR-PRO_IT (Very High-Resolution PROjections for ITaly) over the Italian peninsula, which are obtained with the downscaling of the CMCC global climate model at a convection-permitting scale (Raffa et al., 2023). Indeed, VHR-PRO_IT is generated through dynamic downscaling of the Italy8km-CM climate projection (spatial resolution of about 8 km; output frequency = 6 h; driven CMIP5 GCM = CMCC-CM) at about 2.2 km resolution under the IPCC RCP4.5 and RCP8.5 scenarios, employing the Regional Climate Model COSMO-CLM.

Gauged locations are used to assess the accuracy of VHR-PRO_IT in reproducing observed extremes. More specifically, the observed dataset used as ground truth for the comparison is I²-RED (Improved Italian – Rainfall Extreme Dataset; Mazzoglio et al., 2020). In this work, 742 rain gauges with a minimum of 30 years of short-duration (1, 3, 6, 12, 24 h) annual maximum rainfall depths recorded in the period from 1980 to 2022 are used (Figure 1). The dataset covers the entire peninsula, although some areas are characterized by a lower rain gauge density due to the complex station relocation performed about 30 years ago during the dismantlement of the national hydrological service (SIMN, Servizio Idrografico e Mareografico Nazionale). Conversely, the dataset derived from the VHR-PRO_IT climate projections includes annual maxima from a 30-year time series, connecting the historical period (1981-2005) with 5 years of the RCP8.5 scenario (2006-2010) of the CPM.




Figure 1 – Location of the 742 rain gauges with at least 30 years of data in the 1980-2022 period.

The comparison is made by assessing biases between quantiles of annual maximum rainfall obtained from the two empirical cumulative distribution frequencies. The considered quantiles are 0.1, 0.25, 0.5, 0.75 and 0.9. The biases are shown with respect to the orography and the proximity to the sea, to analyze potential correlation with their geographical position.

Further work will be performed to assess and quantify local biases between L-moments and GEV distributions. Also in this case, evaluating their dependence on the geographic position, orography, and proximity to the sea.

References

- Mazzoglio P., Butera I., Claps P. (2020). I²-RED: a massive update and quality control of the Italian annual extreme rainfall dataset. *Water*, 12, 3308.
- Raffa M., Adinolfi M., Reder A., Marras G.F., Mancini M., Scipione G., Santini M., Mercogliano P. (2023). Very High Resolution Projections over Italy under different CMIP5 IPCC scenarios. *Scientific Data*, 10, 238.