

Wiener-Hopf Solution of Diffraction by a PEC Wedge in Anisotropic Media

Original

Wiener-Hopf Solution of Diffraction by a PEC Wedge in Anisotropic Media / Daniele, V.; Lombardi, G.. - ELETTRONICO. - 2023 International Conference on Electromagnetics in Advanced Applications, ICEAA 2023:(2023), pp. 652-652. (Intervento presentato al convegno 25th International Conference on Electromagnetics in Advanced Applications, ICEAA 2023 tenutosi a Venice (Italy) nel 9-13 October 2023) [10.1109/ICEAA57318.2023.10297849].

Availability:

This version is available at: 11583/2987793 since: 2024-04-13T09:31:40Z

Publisher:

IEEE

Published

DOI:10.1109/ICEAA57318.2023.10297849

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

IEEE postprint/Author's Accepted Manuscript

©2023 IEEE. Personal use of this material is permitted. Permission from IEEE must be obtained for all other uses, in any current or future media, including reprinting/republishing this material for advertising or promotional purposes, creating new collecting works, for resale or lists, or reuse of any copyrighted component of this work in other works.

(Article begins on next page)



Wiener-Hopf Solution of Diffraction by a PEC Wedge in Anisotropic Media

Vito Daniele⁽¹⁾, Guido Lombardi⁽¹⁾

(1) Politecnico di Torino, Italy, e-mail: vito.daniele@polito.it, guido.lombardi@polito.it

In this work we present our recent work on novel Wiener-Hopf (WH) formulation for the analysis and the study of scattering of wedges [1] immersed in complex materials [2-3]. We start from an original study of the perfect electrically conducting (PEC) wedge immersed in uniaxial ($\epsilon_x=\epsilon_y\neq\epsilon_z$, $\mu_x=\mu_y\neq\mu_z$) and biaxial (all different ϵ_i , μ_i $i=x,y,z$) media illuminated by plane waves, where the wedge has an aperture angle of $2\pi-2\gamma$. The uniaxial case has been studied by Felsen in [4], however generalization and exploitation of this case has not been further carried out in literature.

Our formulation allows the analysis of uniaxial case and biaxial case by developing an original theory in the framework of Wiener-Hopf technique, which allows closed form analytical solution in spectral domain. The theory takes inspiration from the one presented in [5] for stratified media with a generalization to angular regions [2-3]. Moreover, in anisotropic media different from uniaxial and biaxial media, a closed form solution is not available, however our theory allows to resort to Fredholm factorization technique that gets approximate solutions preserving physical meaning [6],[1]. From transversalization of Maxwell's equations in oblique Cartesian coordinates, we obtain a matrix differential problem of first order in terms of the field components tangent at the boundaries. In spectral domain, by using Laplace transform, we apply the characteristic Green's function procedure to the system of differential equation yielding a spectral solution before imposing boundary conditions. The solutions represent functional equations that relate the spectral field components along two faces of the angular region. By imposing the boundary conditions the combination of functional equations of each angular region yields a set of GWHEs for the angular region problem ($-\gamma<\varphi<\gamma$). The imposition of PEC boundary conditions on the two faces of the wedge simplifies the problem (vanishing tangent electric field components), however the presence of biaxial medium yields two types of propagation constants. For this reason, the GWHEs present plus functions defined in a complex plane, and minus functions defined into two different complex planes (related to the two propagation constants). We note that the two complex planes of the minus functions are related to the complex plane of the plus functions. The proposed new and original extension of WH technique for the uniaxial and biaxial cases is based on a decoupling procedure among the equations due to the two propagation constants and on a suitable definition of spectral mappings to resort to two systems of Classical Wiener-Hopf Equations (CWHEs) in two new complex planes. Once completed this not easy procedure, simple closed form factorizations allow solutions in spectral domain. Further detail and numerical results will be presented at the conference. Moreover, the sets of derived WH equations are not limited to angular region problems, for example in [7] functional equations of different kind of regions are coupled together.

This work was supported by PRIN Grant 2017NT5W7Z GREEN TAGS and by EU-Next Generation EU within PNRR M4C2, Investimento 1.4-Avviso n.3138 16/12/2021-CN00000013 National Centre for HPC, Big Data and Quantum Computing(HPC)-CUP E13C22000990001-Multiscale modeling and Engineering App.

1. V. Daniele, G. Lombardi, *Scattering and Diffraction by Wedges 2: The Wiener-Hopf Solution - Advanced Applications*, Hoboken, NJ: John Wiley & Sons, Inc. London, UK: ISTE, 2020

2. V. Daniele, The Wiener-Hopf technique for wedge problems, Internal Rep. ELT-2004-2, DET, Politecnico di Torino, 2004, see <http://personal.det.polito.it/vito.daniele>

3. V. Daniele, G. Lombardi, "The Generalized Wiener-Hopf Equations for wave motion in angular regions: electromagnetic application," *Proc. R. Soc. A*, 477:20210040, n.2252, pp. 1-27, 2021, doi:10.1098/rspa.2021.0040

4. L. B. Felsen, "Propagation and diffraction in uniaxially anisotropic regions. i—theory. ii—applications," *Proc. Inst. Elect. Eng.*, vol. 111, no.3, pp. 445–464, Mar. 1964

5. A.D. Bresler, N. Marcuvitz, Operator methods in electromagnetic field theory, Report R-495-56, PIB-425, MRI Polytechnic Institute of Brooklyn, 1956.

6. V. Daniele, G. Lombardi, "Fredholm factorization of Wiener-Hopf scalar and matrix kernels," *Radio Science*, 42(06), pp. 1-19, 2007.

7. V. Daniele, G. Lombardi, "Radiation and Scattering of an Arbitrarily Flanged Dielectric-Loaded Waveguide," *IEEE Transactions on Antennas and Propagation*, vol. 67, no. 12, pp. 7569-7584, Dec. 2019