

On varieties with Ulrich twisted conormal bundle

Original

On varieties with Ulrich twisted conormal bundle / Antonelli, Vincenzo; Casnati, Gianfranco; Lopez, Angelo Felice; Raychaudhury, Debaditya. - In: PROCEEDINGS OF THE AMERICAN MATHEMATICAL SOCIETY. - ISSN 1088-6826. - ELETTRONICO. - 152:(2024), pp. 4645-4658. [10.1090/proc/16986]

Availability:

This version is available at: 11583/2990288 since: 2024-10-15T09:37:39Z

Publisher:

American Mathematical Society - AMS

Published

DOI:10.1090/proc/16986

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

AMS preprint/submitted version e/o postprint/Author's Accepted Manuscript

(Article begins on next page)

ON VARIETIES WITH ULRICH TWISTED CONORMAL BUNDLES

VINCENZO ANTONELLI, GIANFRANCO CASNATI, ANGELO FELICE LOPEZ AND DEBADITYA RAYCHAUDHURY

ABSTRACT. We study varieties $X \subset \mathbb{P}^r$ such that $N_X^*(k)$ is an Ulrich vector bundle for some integer k . We first prove that such an X must be a curve. Then we give several examples of curves with $N_X^*(k)$ an Ulrich vector bundle.

1. INTRODUCTION

Let $X \subset \mathbb{P}^r$ be a smooth variety of dimension $n \geq 1$ over an algebraically closed field F . Recall that a vector bundle \mathcal{E} on X is called Ulrich if $H^i(\mathcal{E}(-p)) = 0$ for all $i \geq 0$ and $1 \leq p \leq n$. The importance of Ulrich vector bundles is well-known (see for example [ES, B, CMRPL] and references therein). While the main general problem about Ulrich vector bundles is their conjectural existence, another line of research around them is what are the consequences on the geometry of X in the presence of an Ulrich vector bundle. In this vein, we continue our study of which natural bundles, associated to X and to its embedding in \mathbb{P}^r , can be Ulrich up to some twist.

In previous papers, the third and fourth authors analyzed normal and tangent bundles, see [L, LR] (see also [BMPT]). In the present paper we study the following question: for which integers k one has that the twisted conormal bundle $N_X^*(k)$ is Ulrich?

A first simple consequence can be drawn: if X is degenerate, that is contained in a hyperplane, then $(X, \mathcal{O}_X(1), k) = (\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(1), 1)$, see Lemma 5.2.

On the other hand, suppose that X is nondegenerate. While in previous cases [L, LR], examples of surfaces and threefolds appeared, we find a very different result for the conormal bundle. In fact we show that the answer to the above question is negative in dimension at least two.

Theorem 1. ($\text{char}(F)=0$)

Let $X \subset \mathbb{P}^r$ be a smooth nondegenerate variety such that $N_X^(k)$ is Ulrich. Then X is a curve.*

Now, for curves the situation is wide. First of all, there are many examples, at least in \mathbb{P}^3 , stemming from some classical works [EL, ElHi, BE] (see Examples 8.1 and 8.2).

We first prove that there is a bound, sharp in codimension 2, for the degree of a curve having Ulrich twisted conormal bundle.

Theorem 2.

Let $C \subset \mathbb{P}^{c+1}$ be a smooth nondegenerate curve of degree d and codimension $c \geq 1$ such that $N_C^(k)$ is Ulrich. Then $c \geq 2$ and*

$$(1.1) \quad d \geq \frac{c+2}{2k+c} \binom{k+c}{c+1}.$$

Moreover this bound is sharp for $c = 2$ and $k \equiv 1, 3 \pmod{6}$.

On the other hand, the examples mentioned above, Examples 8.1 and 8.2, are all subcanonical curves in \mathbb{P}^3 . We show that neither the fact of being subcanonical, nor of lying in \mathbb{P}^3 , is a necessary condition, by producing examples, for unbounded genus, of non-subcanonical curves in \mathbb{P}^3 and in \mathbb{P}^4 .

Theorem 3.

(i) Let $X \subset \mathbb{P}^3$ be a general nonspecial curve of genus g and degree $d = 2g - 2$. Then $N_X^(4)$ is Ulrich and X is not subcanonical.*

The first three authors are members of the GNSAGA group of INdAM. The third author was partially supported by PRIN “Advances in Moduli Theory and Birational Classification”.

Mathematics Subject Classification : Primary 14F06. Secondary 14H60, 14H10, 14J60.

(ii) Let $X \subset \mathbb{P}^4$ be a general curve of genus $g \geq 3$ and degree $d = 5g - 5$. Then $N_X^*(3)$ is Ulrich and X is not subcanonical.

Note that the nonspecial curve of genus 5 and degree 8 in (i) of the above theorem, is a non-subcanonical curve realizing equality in Theorem 2.

Here is a brief overview of the strategy of proofs of the theorems.

Assume that $N_X^*(k)$ is Ulrich. The first ingredient (Lemma 5.3 (iv)) is that X is contained in a hypersurface of degree k . Next, some vanishings hold for the cohomology of twists of \mathcal{O}_X (Lemma 5.3 (i)). In the case of curves, using Riemann-Roch this gives the bound in Theorem 2. In the case of higher dimensional varieties, since the property of $N_X^*(k)$ being Ulrich is preserved by hyperplane sections, one reduces to the surface section S . Also, S is projectively normal (Lemma 5.3 (ii)) and combining again the vanishings, Riemann-Roch and the Bogomolov inequality (because Ulrich bundles are semistable), one contradicts Theorem 2. As for Theorem 3, we use degenerations of nonspecial curves and the techniques of interpolation as in [ALY, LV].

We would like to thank the generous referee for the big contribution given to improve the hypotheses and shorten the proof of Theorem 3.

2. NOTATION

Throughout the paper we work over an algebraically closed field F . In some cases, when needed, we will specify that $\text{char}(F) = 0$. A *variety* is by definition an integral separated scheme of finite type over F . A *curve* (respectively a *surface*) is a variety of dimension 1 (resp. 2). Moreover, we henceforth establish the following:

Notation 2.1.

- $X \subset \mathbb{P}^r$ is a smooth closed variety of dimension $n \geq 1$ and codimension $c = r - n \geq 1$.
- H is a hyperplane divisor.
- $N_X := N_{X/\mathbb{P}^r}$ is the normal bundle.
- For any sheaf \mathcal{G} on X we set $\mathcal{G}(l) = \mathcal{G}(lH)$.
- $d = H^n$ is the degree of X .
- C is a general curve section of X under H .
- S is a general surface section of X under H , when $n \geq 2$.
- $g = g(C) = \frac{1}{2}[K_X H^{n-1} + (n-1)d] + 1$ is the sectional genus of X .
- For $1 \leq i \leq n-1$, let $H_i \in |H|$ be general divisors and set $X_n := X$ and $X_i = H_1 \cap \cdots \cap H_{n-i}$. In particular $X_1 = C, X_2 = S$.
- $s(X) = \min\{s \geq 1 : H^0(\mathcal{J}_{X/\mathbb{P}^r}(s)) \neq 0\}$.

We will also let $V = H^0(\mathcal{O}_{\mathbb{P}^r}(1))$ and consider the exact sequences

$$(2.1) \quad 0 \rightarrow \Omega_{\mathbb{P}^r|X}^1 \rightarrow V \otimes \mathcal{O}_X(-1) \rightarrow \mathcal{O}_X \rightarrow 0$$

and

$$(2.2) \quad 0 \rightarrow N_X^* \rightarrow \Omega_{\mathbb{P}^r|X}^1 \rightarrow \Omega_X^1 \rightarrow 0.$$

3. A GENERAL FACT ABOUT PROJECTIVE VARIETIES

We record here a simple but useful fact.

Lemma 3.1. *Let $X \subset \mathbb{P}^r$ be a smooth variety of dimension $n \geq 1$. If $H^0(N_X^*(l)) = 0$ and $\pi : X \rightarrow \bar{X} \subset \mathbb{P}^m$ is an isomorphic projection, then $l \leq \min\{s(X) - 1, s(\bar{X}) - 1\}$.*

Proof. Set $s = s(X)$ and suppose that $l \geq s$, so that $H^0(N_X^*(s)) = 0$. Now the exact sequence

$$0 \rightarrow \mathcal{J}_{X/\mathbb{P}^r}^2(s) \rightarrow \mathcal{J}_{X/\mathbb{P}^r}(s) \rightarrow N_X^*(s) \rightarrow 0$$

implies that $h^0(\mathcal{J}_{X/\mathbb{P}^r}^2(s)) = h^0(\mathcal{J}_{X/\mathbb{P}^r}(s)) > 0$, hence there is a hypersurface G of degree s such that $X \subseteq \text{Sing}(G)$. Hence there is a non-zero partial derivative of the equation of G , giving a hypersurface of degree $s - 1$ containing X , a contradiction. Therefore

$$(3.1) \quad l \leq s(X) - 1.$$

Now let $\pi : X \rightarrow \bar{X} \subset \mathbb{P}^m$ be an isomorphic projection, so that we have an exact sequence

$$(3.2) \quad 0 \rightarrow \pi^* N_{\bar{X}/\mathbb{P}^m}^* \rightarrow N_X^* \rightarrow \mathcal{O}_X(-1)^{\oplus(r-m)} \rightarrow 0.$$

Since $H^0(N_X^*(l)) = 0$ we get that $H^0(\bar{X}, N_{\bar{X}/\mathbb{P}^m}^*(l)) = H^0(X, \pi^* N_{\bar{X}/\mathbb{P}^m}^*(l)) = 0$. Hence applying (3.1) to $\bar{X} \subset \mathbb{P}^m$ we get that $l \leq s(\bar{X}) - 1$ and the lemma is proved. \square

4. GENERALITIES ON ULRICH BUNDLES

We collect here some well-known facts about Ulrich bundles, to be used sometimes later.

Definition 4.1. Let \mathcal{E} be a vector bundle on X . We say that \mathcal{E} is *Ulrich* for (X, H) if $H^i(\mathcal{E}(-p)) = 0$ for all $i \geq 0$ and $1 \leq p \leq n$.

We have

Lemma 4.2. *Let \mathcal{E} be a rank t Ulrich vector bundle for (X, H) . Then*

- (i) $c_1(\mathcal{E})H^{n-1} = \frac{t}{2}[K_X + (n+1)H]H^{n-1}$.
- (ii) $\mathcal{E}^*(K_X + (n+1)H)$ is also Ulrich for (X, H) .
- (iii) \mathcal{E} is globally generated.
- (iv) \mathcal{E} is arithmetically Cohen-Macaulay (aCM), that is $H^i(\mathcal{E}(j)) = 0$ for $0 < i < n$ and all $j \in \mathbb{Z}$.
- (v) $\mathcal{E}|_Y$ is Ulrich on a smooth hyperplane section Y of X .
- (vi) $\mathcal{O}_X(l)$ is Ulrich if and only if $(X, H, l) = (\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(1), 0)$.

Proof. Well-known. For (i)-(v) see for example [LR, Lemma 3.2]. As for (vi), it is obvious that $\mathcal{O}_{\mathbb{P}^n}$ is Ulrich for $(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(1))$. Vice versa, if $\mathcal{O}_X(l)$ is Ulrich, then it is globally generated by (iii), so that $l \geq 0$. But also $H^0(\mathcal{O}_X(l-1)) = 0$, hence $l = 0$. It follows by [LR, Lemma 3.2(vii)] that $d = h^0(\mathcal{O}_X) = 1$, so that $(X, H) = (\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(1))$. \square

Lemma 4.3. *Let $X \subset \mathbb{P}^r$ be a smooth variety of dimension $n \geq 3$. Let \mathcal{E} be a vector bundle on X and let Y be a smooth hyperplane section of X . If $\mathcal{E}|_Y$ is Ulrich, then \mathcal{E} is Ulrich.*

Proof. For $j \in \mathbb{Z}$ consider the exact sequence

$$(4.1) \quad 0 \rightarrow \mathcal{E}(j-1) \rightarrow \mathcal{E}(j) \rightarrow \mathcal{E}|_Y(j) \rightarrow 0.$$

If $2 \leq i \leq n-2$ we have that $H^{i-1}(\mathcal{E}|_Y(j)) = H^i(\mathcal{E}|_Y(j)) = 0$ for any $j \in \mathbb{Z}$ by Lemma 4.2(iv). Hence (4.1) gives that $h^i(\mathcal{E}(j-1)) = h^i(\mathcal{E}(j))$ for any $j \in \mathbb{Z}$. On the other hand $h^i(\mathcal{E}(j)) = 0$ for $j \gg 0$ and it follows that $h^i(\mathcal{E}(j)) = 0$ for any $j \in \mathbb{Z}$ and $2 \leq i \leq n-2$.

Suppose now that $i \in \{0, 1\}$ and $j \leq -1$. We have that $H^0(\mathcal{E}|_Y(j)) = 0$ and, since $n-1 \geq 2$, also that $H^1(\mathcal{E}|_Y(j)) = 0$ by Lemma 4.2(iv). Hence (4.1) gives that $h^i(\mathcal{E}(j-1)) = h^i(\mathcal{E}(j))$. On the other hand, by Serre duality, $h^i(\mathcal{E}(j)) = h^{n-i}(\mathcal{E}^*(K_X - jH)) = 0$ for $j \ll 0$ and therefore

$$(4.2) \quad h^i(\mathcal{E}(j)) = 0 \text{ for } i \in \{0, 1\} \text{ and } j \leq -1.$$

Now let $\mathcal{E}' = \mathcal{E}^*(K_X + (n+1)H)$. Then $\mathcal{E}'|_Y = \mathcal{E}_Y^*(K_Y + nH|_Y)$ is also Ulrich by Lemma 4.2(ii). Therefore (4.2) implies that $h^i(\mathcal{E}'(j)) = 0$ for $i \in \{0, 1\}$ and $j \leq -1$. By Serre duality we get that $h^{n-i}(\mathcal{E}(-n-1-j)) = h^i(\mathcal{E}^*(K_X + (n+1+j)H)) = h^i(\mathcal{E}'(j)) = 0$ for $i \in \{0, 1\}$ and $j \leq -1$. But this is the same as $h^s(\mathcal{E}(l)) = 0$ for $s \in \{n-1, n\}$ and $l \geq -n$.

Thus we have proved that $H^i(\mathcal{E}(-p)) = 0$ for $i \geq 0$ and $1 \leq p \leq n$, that is \mathcal{E} is Ulrich. \square

5. ULRICH CONORMAL BUNDLES

In this section we will draw some very useful consequences and facts for varieties $X \subset \mathbb{P}^r$ such that $N_X^*(k)$ is Ulrich.

The first one is a reduction via hyperplane sections (for the X_i 's see Notation 2.1).

Lemma 5.1. *Let $X \subset \mathbb{P}^{c+n}$ be a smooth variety of dimension n and codimension $c \geq 1$. If $n \geq 2$ and $N_X^*(k)$ is Ulrich, then $N_{X_i/\mathbb{P}^{c+i}}^*(k)$ is Ulrich for all $i \in \{1, \dots, n-1\}$. Vice versa, if $n \geq 3$ and $N_{X_i/\mathbb{P}^{c+i}}^*(k)$ is Ulrich for some $i \in \{2, \dots, n-1\}$, then $N_{X_j/\mathbb{P}^{c+j}}^*(k)$ is Ulrich for all $j \in \{2, \dots, n\}$ (hence in particular so is $N_X^*(k)$).*

Proof. Recall that if $Y \subset \mathbb{P}^m$ is smooth and Z is a smooth hyperplane section, then

$$(N_{Y/\mathbb{P}^m})|_Z \cong N_{Z/\mathbb{P}^{m-1}}.$$

Now if $N_X^*(k)$ is Ulrich, then so are all $N_{X_i/\mathbb{P}^{c+i}}^*(k)$ by Lemma 4.2(v).

Vice versa, if $N_{X_i/\mathbb{P}^{c+i}}^*(k)$ is Ulrich for some $i \in \{2, \dots, n-1\}$, then $N_{X_{i+1}/\mathbb{P}^{c+i+1}}^*(k)$ is Ulrich by Lemma 4.3. Repeating the argument we get that $N_{X_j/\mathbb{P}^{c+j}}^*(k)$ is Ulrich for all $j \in \{2, \dots, n\}$. \square

We now deal with X degenerate in \mathbb{P}^r .

Lemma 5.2. *Let $X \subset \mathbb{P}^r$ be a smooth degenerate variety of dimension $n \geq 1$. Then $N_X^*(k)$ is Ulrich if and only if $(X, H, k) = (\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(1), 1)$.*

Proof. If $(X, H) = (\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(1))$ then $N_X^*(1) = \mathcal{O}_{\mathbb{P}^n}^{\oplus c}$ is Ulrich.

Vice versa assume that $N_X^*(k)$ is Ulrich. Since X is degenerate, $N_X^*(k)$ has $\mathcal{O}_X(k-1)$ as a direct summand. Therefore also $\mathcal{O}_X(k-1)$ is Ulrich and Lemma 4.2(vi) gives that $(X, H, k) = (\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(1), 1)$. \square

In the sequel we will then consider only nondegenerate varieties.

We start by collecting some cohomological and numerical conditions.

Lemma 5.3. (cohomological conditions)

Let $X \subset \mathbb{P}^{c+n}$ be a smooth nondegenerate variety of dimension $n \geq 1$ and codimension $c \geq 1$. If $N_X^(k)$ is Ulrich, we have:*

- (i) $H^n(\mathcal{O}_X(l)) = 0$ for every $l \geq k - n - 1$.
- (ii) If $n \geq 2$, then $X \subset \mathbb{P}^{c+n}$ is projectively normal.
- (iii) If $n \geq 2$ then $q(X) = 0$.
- (iv) $k \leq \min\{s(X), s(\bar{X})\}$, where $\bar{X} \subset \mathbb{P}^r$ is any isomorphic projection of X .

Proof. By hypothesis $N_X^*(k)$ is Ulrich, hence it is aCM by Lemma 4.2(iv).

To see (i) we just need to prove that $H^n(\mathcal{O}_X(k - n - 1)) = 0$. Assume that $H^n(\mathcal{O}_X(k - n - 1)) \neq 0$, that is, by Serre duality, $H^0(K_X + (n + 1 - k)H) \neq 0$. Then we have an inclusion

$$N_X(-1) \hookrightarrow N_X(K_X + (n - k)H)$$

and, since $N_X(-1)$ is globally generated, we get that $h^0(N_X(K_X + (n - k)H)) \neq 0$. On the other hand, $N_X(K_X + (n + 1 - k)H)$ is Ulrich by Lemma 4.2(ii), hence $h^0(N_X(K_X + (n - k)H)) = 0$. This contradiction proves (i). To see (ii), let $V = H^0(\mathcal{O}_{\mathbb{P}^{c+n}}(1))$ and let $P^1(\mathcal{O}_X(1))$ be the sheaf of principal parts and consider, as in [E, Proof of Thm. 2.4], the following commutative diagram

$$\begin{array}{ccccccc} & & 0 & & 0 & & \\ & & \downarrow & & \downarrow & & \\ & & N_X^*(1) & \xrightarrow{\cong} & N_X^*(1) & & \\ & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & \Omega_{\mathbb{P}^{c+n}}^1(1)|_X & \longrightarrow & V \otimes \mathcal{O}_X & \longrightarrow & \mathcal{O}_X(1) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \cong \\ 0 & \longrightarrow & \Omega_X^1(1) & \longrightarrow & P^1(\mathcal{O}_X(1)) & \longrightarrow & \mathcal{O}_X(1) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \\ & & 0 & & 0 & & \end{array}$$

Pick an integer $l \geq 0$. Tensoring the above diagram by $\mathcal{O}_X(l)$ and observing that

$$P^1(\mathcal{O}_X(1)) \otimes \mathcal{O}_X(l) \cong P^1(\mathcal{O}_X(l + 1))$$

by [E, (2.2)], we get the commutative diagram

$$(5.1) \quad \begin{array}{ccc} V \otimes H^0(\mathcal{O}_X(l)) & & \\ \downarrow f_l & \searrow h_l & \\ H^0(P^1(\mathcal{O}_X(l+1))) & \xrightarrow{g_l} & H^0(\mathcal{O}_X(l+1)) \\ \downarrow & & \\ H^1(N_X^*(l+1)). & & \end{array}$$

Now we have that $H^1(N_X^*(l+1)) = 0$ since N_X^* is aCM and $n \geq 2$. Hence f_l is surjective for every $l \geq 0$ and so is g_l by [E, Prop. 2.3]. It follows by (5.1) that h_l is surjective for every $l \geq 0$. Moreover the commutative diagram

$$(5.2) \quad \begin{array}{ccc} V \otimes H^0(\mathcal{O}_{\mathbb{P}^{c+n}}(l)) & \longrightarrow & H^0(\mathcal{O}_{\mathbb{P}^{c+n}}(l+1)) \\ \downarrow \text{Id}_V \otimes r_l & & \downarrow r_{l+1} \\ V \otimes H^0(\mathcal{O}_X(l)) & \xrightarrow{h_l} & H^0(\mathcal{O}_X(l+1)) \end{array}$$

shows by induction that $r_l : H^0(\mathcal{O}_{\mathbb{P}^{c+n}}(l)) \rightarrow H^0(\mathcal{O}_X(l))$ is surjective for every $l \geq 0$, so that $X \subset \mathbb{P}^r$ is projectively normal, that is (ii).

To see (iii) observe that, (2.1) gives an exact sequence

$$(5.3) \quad 0 \longrightarrow H^0(\Omega_{\mathbb{P}^{c+n}}^1(1)|_X) \longrightarrow V \xrightarrow{f} H^0(\mathcal{O}_X(1))$$

and f is injective since X is nondegenerate, hence $H^0(\Omega_{\mathbb{P}^{c+n}}^1(1)|_X) = 0$. Now $N_X^*(k)$ is aCM and therefore $H^1(N_X^*(1)) = 0$. Then the exact sequence

$$(5.4) \quad 0 \rightarrow N_X^*(1) \rightarrow \Omega_{\mathbb{P}^{c+n}}^1(1)|_X \rightarrow \Omega_X^1(1) \rightarrow 0$$

shows that $H^0(\Omega_X^1(1)) = 0$, hence, in particular $q(X) = h^0(\Omega_X^1) = 0$. This proves (iii). Finally (iv) follows by Lemma 3.1 since, $N_X^*(k)$ being Ulrich, we have that $H^0(N_X^*(k-1)) = 0$. \square

Lemma 5.4. (numerical conditions)

Let $X \subset \mathbb{P}^{c+n}$ be a smooth nondegenerate variety of dimension $n \geq 1$ and codimension $c \geq 1$ and degree d . If $N_X^*(k)$ is Ulrich, we have:

- (i) $[(k-2)c-2]d = (c+2)(g-1)$.
- (ii) $c \geq 2$.
- (iii) $k \geq 3$.

Proof. By hypothesis $N_X^*(k)$ is Ulrich. By (2.1) and (2.2) we see that $c_1(N_X^*(k)) = -K_X - (c+n+1-kc)H$. Hence Lemma 4.2(i) implies

$$-(K_X + (c+n+1-kc)H)H^{n-1} = \frac{c}{2} (K_X H^{n-1} + (n+1)d)$$

and this gives $K_X H^{n-1} = (2k-n-3-\frac{4(k-1)}{c+2})d$. But also $K_X H^{n-1} = 2(g-1) - (n-1)d$ and we get (i). As for (ii), if $c=1$ then $N_X^*(k) = \mathcal{O}_X(k-d)$ and Lemma 4.2(vi) gives that $X \subset \mathbb{P}^r$ is a linear space, a contradiction. This proves (ii). To see (iii), since X is nondegenerate, we get from (2.1) that $H^0(\Omega_{\mathbb{P}^{c+n}}^1(1)|_X) = 0$. Now (2.2) gives that $h^0(\Omega_{\mathbb{P}^{c+n}}^1(1)|_X(k)) \geq h^0(N_X^*(k)) > 0$ by Lemma 4.2(iii). Hence $k \geq 2$. But if $k=2$ then (i) gives that $g=0$ and $d = \frac{c+2}{2}$. As it is well known, $d \geq c+1$, giving a contradiction. Thus (iii) holds. \square

6. PROPERTIES OF THE SURFACE SECTION

We deduce here some very useful properties of the surface section of some $X \subset \mathbb{P}^r$ such that $N_X^*(k)$ is Ulrich. We assume that $\text{char}(F) = 0$.

Lemma 6.1. *Let $X \subset \mathbb{P}^{c+n}$ be a smooth nondegenerate variety of dimension $n \geq 2$ and codimension $c \geq 1$. If $N_X^*(k)$ is Ulrich, the following inequality holds for the surface section S :*

$$\chi(\mathcal{O}_S) \geq \frac{d}{2(c+2)(c+3)(3c+4)(c+12)} [(3c^4 + 43c^3 + 86c^2 + 24c)k^2 - (15c^4 + 229c^3 + 624c^2 + 432c)k + 18c^4 + 296c^3 + 1070c^2 + 1368c + 576].$$

Proof. Note that $N_{S/\mathbb{P}^{c+2}}^*(k)$ is Ulrich by Lemma 5.1. Therefore, Lemma 5.3(iii) implies that

$$(6.1) \quad q(S) = 0.$$

Next, computing the Chern classes of $N_{S/\mathbb{P}^{c+2}}^*(k)$ and applying [C, (2.2)], we get

$$(6.2) \quad cK_S^2 + (c+12)c_2(S) = \frac{6d}{c+2} [c(c+2)k^2 - c(5c+12)k + 6c^2 + 20c + 12].$$

Now note that $N_{S/\mathbb{P}^{c+2}}^*(k)$ is semistable by [CH, Thm. 2.9], hence so is $N_{S/\mathbb{P}^{c+2}}^*$ and since $\text{rk}(N_{S/\mathbb{P}^{c+2}}^*) = c$, the Bogomolov inequality is

$$(6.3) \quad d(c+3)(4k-7-\frac{8(k-1)}{c+2}) + (c+1)K_S^2 \geq (2c)c_2(S).$$

Then (6.2) and (6.3) give

$$(6.4) \quad K_S^2 \geq \frac{d}{(c+2)(c+3)(3c+4)} [(12c^3 + 24c^2)k^2 - (64c^3 + 204c^2 + 144c)k + 79c^3 + 351c^2 + 486c + 216].$$

Finally the inequality on $\chi(\mathcal{O}_S)$ in the statement follows by (6.2), (6.4) and Noether's formula. \square

7. PROOFS OF MAIN THEOREMS

We start by proving Theorem 2.

Proof of Theorem 2. Since $N_C^*(k)$ is Ulrich, it follows that $c \geq 2$ by Lemma 5.4(ii) and $k \leq s(C)$ by Lemma 5.3(iv). Also Lemma 5.3(i) gives $H^1(\mathcal{O}_C(k-2)) = 0$. Therefore we have that $H^0(\mathcal{J}_{C/\mathbb{P}^{c+1}}(k-1)) = H^1(\mathcal{O}_C(k-1)) = 0$ and the exact sequence

$$0 \rightarrow \mathcal{J}_{C/\mathbb{P}^{c+1}}(k-1) \rightarrow \mathcal{O}_{\mathbb{P}^{c+1}}(k-1) \rightarrow \mathcal{O}_C(k-1) \rightarrow 0$$

together with Riemann-Roch, shows that

$$(7.1) \quad \binom{k+c}{c+1} = h^0(\mathcal{O}_{\mathbb{P}^{c+1}}(k-1)) \leq h^0(\mathcal{O}_C(k-1)) = d(k-1) - g + 1.$$

Also, $g-1 = \frac{(k-2)c-2}{c+2}d$ by Lemma 5.4(i) and replacing in (7.1) we get (1.1).

Finally, sharpness for $c=2$ and $k \equiv 1, 3 \pmod{6}$ follows by [BE, Examples, p. 88], see Example 8.2. \square

Next, we prove Theorem 1.

Proof of Theorem 1. Suppose that $n \geq 2$. In order to simplify the calculations we set

$$A = (3c^4 + 43c^3 + 86c^2 + 24c)k^2 - (15c^4 + 229c^3 + 624c^2 + 432c)k + 18c^4 + 296c^3 + 1070c^2 + 1368c + 576$$

so that it follows by Lemma 6.1 that

$$(7.2) \quad \chi(\mathcal{O}_S) \geq \frac{dA}{2(c+2)(c+3)(3c+4)(c+12)}.$$

Now Lemma 5.1 implies that $N_{S/\mathbb{P}^{c+2}}^*(k)$ is Ulrich. Hence $k \leq s(S)$ by Lemma 5.3(iv), $H^2(\mathcal{O}_S(k-2)) = 0$ by Lemma 5.3(i) and $S \subset \mathbb{P}^{c+2}$ is projectively normal by Lemma 5.3(ii). Therefore, we have that

$$H^0(\mathcal{J}_{S/\mathbb{P}^{c+2}}(k-2)) = H^1(\mathcal{J}_{S/\mathbb{P}^{c+2}}(k-2)) = H^2(\mathcal{O}_S(k-2)) = 0$$

and the exact sequence

$$0 \rightarrow \mathcal{J}_{S/\mathbb{P}^{c+2}}(k-2) \rightarrow \mathcal{O}_{\mathbb{P}^{c+2}}(k-2) \rightarrow \mathcal{O}_S(k-2) \rightarrow 0$$

together with Riemann-Roch and Lemma 5.4(i), shows that

$$(7.3) \quad \binom{k+c}{c+2} = h^0(\mathcal{O}_{\mathbb{P}^{c+2}}(k-2)) = h^0(\mathcal{O}_S(k-2)) = \chi(\mathcal{O}_S(k-2)) + h^1(\mathcal{O}_S(k-2)) \geq \\ \geq \chi(\mathcal{O}_S) + \frac{d(k-2)}{2} \left(3 - k + \frac{4(k-1)}{c+2}\right).$$

Setting

$$B = A + (k-2)(c+3)(3c+4)(c+12)[(3-k)(c+2) + 4k-4] = 2(c+12)(k-1)(8ck + 12k + 5c^2 + 7c)$$

we see, using (7.2), that (7.3) implies

$$(7.4) \quad \binom{k+c}{c+2} \geq \frac{dB}{2(c+2)(c+3)(3c+4)(c+12)} = \frac{d(k-1)(8ck + 12k + 5c^2 + 7c)}{(c+2)(c+3)(3c+4)}.$$

Since $k \geq 3$ by Lemma 5.4(iii), we see that (7.4) gives

$$d \leq \frac{(c+2)(c+3)(3c+4)}{(k-1)(8ck + 12k + 5c^2 + 7c)} \binom{k+c}{c+2}.$$

Since $N_X^*(k)$ is Ulrich, it follows by Lemma 5.1 that $N_{C/\mathbb{P}^{c+1}}^*(k)$ is Ulrich, hence, using Theorem 2, we find that

$$\frac{c+2}{2k+c} \binom{k+c}{c+1} \leq d \leq \frac{(c+2)(c+3)(3c+4)}{(k-1)(8ck + 12k + 5c^2 + 7c)} \binom{k+c}{c+2}$$

that is equivalent to $2c(c+1)(c+k+1) \leq 0$, a contradiction. \square

8. CURVES

In this section we construct some examples of curves $C \subset \mathbb{P}^{c+1}$ such that $N_C^*(k)$ is Ulrich.

First, we give a reinterpretation of two known cases.

Example 8.1. For every integer $d \geq 5$ there exists a smooth elliptic curve $C \subset \mathbb{P}^3$ such that $N_C^*(3)$ is Ulrich.

In fact, it follows by [EL, Prop. §8, page 278] and [ElHi, Thm. 2(b)] that for every integer $d \geq 5$ there exists a smooth elliptic curve $C \subset \mathbb{P}^3$ such that $H^0(N_C(-2)) = 0$. Since $\chi(N_C(-2)) = 0$ we get that $N_C(-2)$ is Ulrich and then also $N_C^*(3)$ is Ulrich by Lemma 4.2(ii).

Example 8.2. There are many subcanonical curves $X \subset \mathbb{P}^3$ with $H^0(N_X(-2)) = 0$ (see for example [BE, Examples, p. 88]), hence with $N_X(-1)$ Ulrich. Therefore, since $K_X = eH$ for some $e \in \mathbb{Z}$, we also have by Lemma 4.2(ii), that $N_X^*(e+3)$ is Ulrich.

We now describe the curves in [BE, Examples, p. 88]. Let $h \in \mathbb{Z}$ such that $h \geq 1$, let $c_2 = h(3h+2)$ and let $t = 3h+1$ (resp. $h \geq 0, c_2 = 3h^2 + 4h + 1$ and $t = 3h+2$). Then in loc. cit, there are examples of smooth curves $X \subset \mathbb{P}^3$ with $N_X^*(e+3)$ Ulrich, $d = t^2 + c_2$ and $g = (t-2)d + 1$. We have $e = \frac{2g-2}{d} = 2t-4$, hence $N_X^*(k)$ is Ulrich with $k = e+3 = 2t-1$. In the first case, $t = 3h+1$, we have $h = \frac{k-1}{6}, k \equiv 1 \pmod{6}$ and $d = \frac{1}{3}(k^2 + 2k)$. In the second case, $t = 3h+2$, we have $h = \frac{k-3}{6}, k \equiv 3 \pmod{6}$ and again $d = \frac{1}{3}(k^2 + 2k)$. In particular they have unbounded k .

The examples above are all subcanonical curves. In order to construct non-subcanonical ones, we will proceed below by degeneration applying the techniques of interpolation as in [ALY, LV]. For the reader's convenience, we recall some definitions and notation about elementary modifications of vector bundles (see [ALY, LV]).

Definition 8.3. Let Z be a scheme, let \mathcal{E} be a vector bundle on Z , let D be an effective Cartier divisor on Z and let \mathcal{F} be a subbundle of $\mathcal{E}|_D$. Then

$$\mathcal{E}[D \xrightarrow{-} \mathcal{F}] = \text{Ker}\{\mathcal{E} \rightarrow \mathcal{E}|_D/\mathcal{F}\}, \quad \mathcal{E}[D \rightarrow \mathcal{F}] = \mathcal{E}[D \xrightarrow{-} \mathcal{F}], \quad \mathcal{E}[D \xrightarrow{+} \mathcal{F}] = \mathcal{E}[D \xrightarrow{-} \mathcal{F}](D).$$

Definition 8.4. Let X be a smooth curve, let $f : X \rightarrow \mathbb{P}^r$ be an unramified morphism and let $N_f = \text{Ker}\{f^*\Omega_{\mathbb{P}^r} \rightarrow \Omega_X\}^\vee$ be the normal bundle of the map f . Given a subspace $\Lambda \subset \mathbb{P}^r$ of dimension λ , consider the projection $\pi_\Lambda : \mathbb{P}^r \dashrightarrow \mathbb{P}^{r-\lambda-1}$. Let U be the open subset of $X \setminus X \cap \Lambda$ where $\pi_\Lambda \circ f$

is unramified. Then $N_{f \rightarrow \Lambda}$ is the unique subbundle of N_f whose restriction to U is $\text{Ker}\{(N_f)|_U \rightarrow (N_{\pi_\Lambda \circ f})|_U\}$. Moreover,

$$N_f[D \xrightarrow{+} \Lambda] := N_f[D \xrightarrow{+} N_{f \rightarrow \Lambda}] \quad \text{and} \quad N_f[D \rightarrow \Lambda] := N_f[D \xrightarrow{-} N_{f \rightarrow \Lambda}].$$

When f is an embedding, we set $N_X[D \rightarrow \Lambda] = N_f[D \rightarrow \Lambda]$.

Next, we prove Theorem 3.

Proof of Theorem 3(i). From $d = 2g - 2$ we deduce that $g \geq 2$, hence, since X is nonspecial, $g - 1 = h^0(\mathcal{O}_C(1)) \geq 3$. Thus $g \geq 4$ and if equality holds, then C is a plane curve of degree 6, a contradiction. Therefore $g \geq 5$.

Let $E \subset \mathbb{P}^3$ be a general elliptic curve of degree $g - 1$ and let L_i be general chords of E with $L_i \cap E = \{p_i, q_i\}$, $1 \leq i \leq g - 1$. Let $Y = L_1 \cup \dots \cup L_{g-1}$ and let

$$X' = E \cup Y \subset \mathbb{P}^3.$$

Claim 8.5. *With the above notation, we have*

$$N_{X'|_Y} \otimes \omega_{X'}(-3) \cong \bigoplus_{i=1}^{g-1} \mathcal{O}_{L_i}(-1)^{\oplus 2}.$$

Proof. For each $i \in \{1, \dots, g - 1\}$ set $Z_i = E \cup L_1 \cup \dots \cup L_{i-1} \cup L_{i+1} \cup \dots \cup L_{g-1}$. By [LV, Prop. 3.5], if we choose points $r_i \in T_{p_i}Z_i \setminus \{p_i\}$, $r'_i \in T_{q_i}Z_i \setminus \{q_i\}$, we have

$$(8.1) \quad N_{X'|_{L_i}} \cong N_{L_i}[p_i \xrightarrow{+} r_i][q_i \xrightarrow{+} r'_i] = N_{L_i}[p_i \xrightarrow{-} r_i][q_i \xrightarrow{-} r'_i](p_i + q_i).$$

On the other hand, let $f : L_i \rightarrow \mathbb{P}^3$ be the embedding. Since $r_i \notin L_i$, we have that $f \circ \pi_{r_i} : L_i \rightarrow \mathbb{P}^2$ is an embedding, and therefore $N_{f \rightarrow \{r_i\}} = \mathcal{O}_{L_i}(1)$ and $N_{L_i}/N_{f \rightarrow \{r_i\}}$ is a line bundle on L_i . Therefore

$$(N_{L_i})|_{\{p_i\}} / (N_{f \rightarrow \{r_i\}})|_{\{p_i\}} \cong \mathcal{O}_{\{p_i\}}$$

and, similarly,

$$(N_{L_i})|_{\{q_i\}} / (N_{f \rightarrow \{r'_i\}})|_{\{q_i\}} \cong \mathcal{O}_{\{q_i\}}.$$

Hence

$$N_{L_i}[p_i \xrightarrow{-} r_i][q_i \xrightarrow{-} r'_i] = \text{Ker}\{N_{L_i} \rightarrow \mathcal{O}_{\{p_i\}} \oplus \mathcal{O}_{\{q_i\}}\} = \text{Ker}\{\mathcal{O}_{\mathbb{P}^1}(1)^{\oplus 2} \rightarrow \mathcal{O}_{\{p_i\}} \oplus \mathcal{O}_{\{q_i\}}\} \cong \mathcal{O}_{\mathbb{P}^1}^{\oplus 2}.$$

It follows by (8.1) that

$$N_{X'|_{L_i}} \cong \mathcal{O}_{\mathbb{P}^1}(p_i + q_i)^{\oplus 2} = \mathcal{O}_{\mathbb{P}^1}(2)^{\oplus 2}$$

and therefore

$$N_{X'|_Y} \otimes \omega_{X'}(-3) \cong \bigoplus_{i=1}^{g-1} \mathcal{O}_{L_i}(-1)^{\oplus 2}.$$

□

Claim 8.6. *With the above notation, we have that $H^0(N_{X'|_E}(-3)) = 0$.*

Proof. By [LV, Prop. 3.5], since $q_i \in L_i \setminus \{p_i\}$, $p_i \in L_i \setminus \{q_i\}$, we have

$$N_{X'|_E} \cong N_E[p_1 \xrightarrow{+} q_1] \cdots [p_{g-1} \xrightarrow{+} q_{g-1}] [q_1 \xrightarrow{+} p_1] \cdots [q_{g-1} \xrightarrow{+} p_{g-1}].$$

Next, specialize cyclically $q_1 \rightarrow p_2, q_2 \rightarrow p_3, \dots, q_{g-1} \rightarrow p_1$. Then, we get a flat family of vector bundles with central fiber

$$N_E[p_1 \xrightarrow{+} p_2] \cdots [p_{g-1} \xrightarrow{+} p_1] [p_2 \xrightarrow{+} p_1] \cdots [p_1 \xrightarrow{+} p_{g-1}].$$

We now claim that

$$(8.2) \quad N_E[p_1 \xrightarrow{+} p_2] \cdots [p_{g-1} \xrightarrow{+} p_1] [p_2 \xrightarrow{+} p_1] \cdots [p_1 \xrightarrow{+} p_{g-1}] \cong N_E(p_1 + \dots + p_{g-1}).$$

To see (8.2), let $f : E \rightarrow \mathbb{P}^3$ be the embedding. For every point $q \in E$, any modification of type p pointing towards the linear space $\{q\}$, is obtained by taking a quotient $(N_E)_{|\{p\}} / (N_{f \rightarrow \{q\}})_{|\{p\}}$. Since $N_E / N_{f \rightarrow \{q\}}$ is a line bundle on E , we have that

$$(N_E)_{|\{p\}} / (N_{f \rightarrow \{q\}})_{|\{p\}} \cong \mathcal{O}_{\{p\}}.$$

Therefore,

$$\begin{aligned} N_E[p_1 \xrightarrow{-} p_2] \cdots [p_{g-1} \xrightarrow{-} p_1] [p_2 \xrightarrow{-} p_1] \cdots [p_1 \xrightarrow{-} p_{g-1}] &= \\ &= \text{Ker}\{N_E \rightarrow \mathcal{O}_{\{p_1\}} \oplus \cdots \mathcal{O}_{\{p_{g-1}\}} \oplus \mathcal{O}_{\{p_2\}} \cdots \mathcal{O}_{\{p_{g-1}\}} \oplus \mathcal{O}_{\{p_1\}}\} = \\ &= \text{Ker}\{N_E \rightarrow \mathcal{O}_{\{p_1\}}^{\oplus 2} \oplus \cdots \mathcal{O}_{\{p_{g-1}\}}^{\oplus 2}\} = N_E(-p_1 - \cdots p_{g-1}). \end{aligned}$$

We deduce that

$$\begin{aligned} N_E[p_1 \xrightarrow{+} p_2] \cdots [p_{g-1} \xrightarrow{+} p_1] [p_2 \xrightarrow{+} p_1] \cdots [p_1 \xrightarrow{+} p_{g-1}] &= \\ &= N_E[p_1 \xrightarrow{-} p_2] \cdots [p_{g-1} \xrightarrow{-} p_1] [p_2 \xrightarrow{-} p_1] \cdots [p_1 \xrightarrow{-} p_{g-1}] (2p_1 + \cdots 2p_{g-1}) \cong N_E(p_1 + \cdots + p_{g-1}) \end{aligned}$$

and this proves (8.2).

Thus we have proved that $N_{X'|_E}(-3)$ specializes to $N_E(p_1 + \cdots + p_{g-1})(-3)$ and to prove the claim it suffices to show that $H^0(N_E(p_1 + \cdots + p_{g-1})(-3)) = 0$. Now, $\chi(N_E(p_1 + \cdots + p_{g-1})(-3)) = 0$ and $\mathcal{O}_E(p_1 + \cdots + p_{g-1})(-3)$ is a general line bundle of degree $-2(g-1)$, hence $H^0(N_E(p_1 + \cdots + p_{g-1})(-3)) = 0$ by interpolation [ALY, Thm. 1.2], [LV, Thm. 1.4]. \square

We now conclude the proof of Theorem 3(i). We have that $X' = E \cup Y$ is a nodal nonspecial curve with $\deg X' = 2g - 2$ and $p_a(X') = g$. In particular X' is smoothable in \mathbb{P}^3 . Moreover, the exact sequence

$$0 \rightarrow N_{X'|_E}(-3) \rightarrow N_{X'} \otimes \omega_{X'}(-3) \rightarrow N_{X'|_Y} \otimes \omega_{X'}(-3) \rightarrow 0$$

and Claims 8.5 and 8.6 show that $H^0(N_{X'} \otimes \omega_{X'}(-3)) = 0$. Then a general smoothing $X \subset \mathbb{P}^3$ of X' is such that $\deg X = 2g - 2$, $g(X) = g$ and $N_X^*(4)$ is Ulrich. Finally X is not subcanonical, for otherwise we would have that $K_C = H$, contradicting the fact that X is nonspecial. \square

Finally, we prove Theorem 3(ii).

Proof of Theorem 3(ii). Let $\overline{C} \subset \mathbb{P}^3$ be a general nonspecial curve of degree $3g - 3$ and genus $g \geq 3$. Since \overline{C} is nonspecial, we can embed it as a nonspecial curve $C \subset \mathbb{P}^4$ of degree $3g - 2$ and genus g , so that \overline{C} is just the projection from a general point $q \in C^1$. Let $p_i, 1 \leq i \leq 2g - 3$ be general points on C and let L_i be general 1-secant lines, each meeting C at p_i . Let $Y = L_1 \cup \cdots L_{2g-3}$, let $q_i \in L_i \setminus \{p_i\}$ and let

$$X' = C \cup Y \subset \mathbb{P}^4.$$

Note that X' is nonspecial, hence we can choose a flat family $\mathcal{X} \rightarrow \Delta$, with smooth total space, having as a general fiber X , a general curve in \mathbb{P}^4 of genus g and degree $5g - 5$ and special fiber X' . Since C is a Cartier divisor on \mathcal{X} we have that $\omega_{\mathcal{X}/\Delta}(-2)(C)$ is a line bundle on \mathcal{X} . Note that, since $C + Y$ is a fiber, we have that $C|_C = -Y|_C$. Setting

$$\mathcal{L} = \omega_{\mathcal{X}/\Delta}(-2)(C)|_{X'}$$

we have that

$$\mathcal{L}|_C = \omega_C(Y \cap C)(-2)(-Y|_C) = \omega_C(-2)$$

and, for each i ,

$$\mathcal{L}|_{L_i} = \omega_{L_i}(L_i \cap C)(-2)(L_i \cap C) = \mathcal{O}_{L_i}(-2).$$

Now we have

Claim 8.7. *Let $\mathcal{E} = N_C(p_1 + \cdots + p_{2g-3})[2p_1 \rightarrow q_1] \cdots [2p_{2g-3} \rightarrow q_{2g-3}] \otimes \omega_C(-2)$. If $H^0(\mathcal{E}) = 0$, then $H^0(N_{X'} \otimes \mathcal{L}) = 0$.*

¹For the construction of the curves C and \overline{C} we could also have appealed to [EiHa, Thm. 1], [CPJLLV].

Proof. Note that $\chi(\mathcal{E}) = 0$. In fact, we have that

$$\chi(N_C(p_1 + \dots + p_{2g-3})) = 20g - 18$$

so that, using [ALY, Prop. 3.3(a)], we get

$$\chi(N_C(p_1 + \dots + p_{2g-3})[2p_1 \rightarrow q_1] \cdots [2p_{2g-3} \rightarrow q_{2g-3}]) = \chi(N_C(p_1 + \dots + p_{2g-3})) - 4(2g-3) = 12g - 6$$

and therefore

$$\chi(\mathcal{E}) = 12g - 6 + 3 \deg(\omega_C(-2)) = 0.$$

Since $H^0(\mathcal{E}) = \chi(\mathcal{E}) = 0$, we have that $H^1(\mathcal{E}) = 0$ and for every divisor $D > 0$ on C we get that $H^0(\mathcal{E}(-D)) \subseteq H^0(\mathcal{E}) = 0$, hence \mathcal{E} satisfies interpolation. As in [ALY, Lemma 8.5] (see also [LV, Lemma 5.2]), we get that $N_{X'} \otimes \mathcal{L}$ satisfies interpolation, so in particular $H^1(N_{X'} \otimes \mathcal{L}) = 0$. On the other hand, $\chi(N_{X'} \otimes \mathcal{L}) = \chi(N_{X'}) + 3 \deg \mathcal{L} = 0$, and therefore $H^0(N_{X'} \otimes \mathcal{L}) = 0$. \square

In order to show that $H^0(\mathcal{E}) = 0$, we specialize each q_i to q , so that \mathcal{E} specializes to

$$\begin{aligned} N_C(p_1 + \dots + p_{2g-3})[2p_1 \rightarrow q] \cdots [2p_{2g-3} \rightarrow q] \otimes \omega_C(-2) &\cong \\ \cong N_C(p_1 + \dots + p_{2g-3})[2p_1 + \dots + 2p_{2g-3} \rightarrow q] \otimes \omega_C(-2). \end{aligned}$$

We have

Claim 8.8. *Set $\mathcal{E}_1 = N_C(p_1 + \dots + p_{2g-3})[2p_1 + \dots + 2p_{2g-3} \rightarrow q] \otimes \omega_C(-2)$. Then there is an exact sequence*

$$(8.3) \quad 0 \rightarrow \omega_C(-1)(2q + p_1 + \dots + p_{2g-3}) \rightarrow \mathcal{E}_1 \rightarrow N_{\overline{C}/\mathbb{P}^3}(q - p_1 - \dots - p_{2g-3}) \otimes \omega_C(-2) \rightarrow 0.$$

Proof. Set $D = p_1 + \dots + p_{2g-3}$. By [LV, page 10], the pointing (towards q) bundle exact sequence [LV, (3.4)], becomes

$$0 \rightarrow \mathcal{O}_C(1)(2q) \rightarrow N_C \rightarrow N_{\overline{C}/\mathbb{P}^3}(q) \rightarrow 0$$

hence, tensoring by $\mathcal{O}_C(D)$, we find an exact sequence

$$(8.4) \quad 0 \rightarrow \mathcal{O}_C(1)(2q + D) \rightarrow N_C(D) \rightarrow N_{\overline{C}/\mathbb{P}^3}(q + D) \rightarrow 0.$$

Applying the elementary modification $[2D \rightarrow q]$ we get the exact sequence

$$0 \rightarrow \mathcal{O}_C(1)(2q + D) \rightarrow N_C(D)[2D \rightarrow q] \rightarrow N_{\overline{C}/\mathbb{P}^3}(q + D)(-2D) \rightarrow 0$$

and tensoring by $\omega_C(-2)$ we get (8.3). \square

Since $\omega_C(-1)(2q + p_1 + \dots + p_{2g-3})$ is a general line bundle of degree $g - 1$ on C , we have that

$$H^0(\omega_C(-1)(2q + p_1 + \dots + p_{2g-3})) = 0.$$

Also, $\omega_C(-2)(q - p_1 - \dots - p_{2g-3})$ a general line bundle of degree $-6(g - 1)$ on C and

$$\chi(N_{\overline{C}/\mathbb{P}^3}(q - p_1 - \dots - p_{2g-3}) \otimes \omega_C(-2)) = 0$$

hence

$$H^0(N_{\overline{C}/\mathbb{P}^3}(q - p_1 - \dots - p_{2g-3}) \otimes \omega_C(-2)) = 0$$

by interpolation [ALY, Thm. 1.2], [LV, Thm. 1.4]. Then (8.3) implies that $H^0(\mathcal{E}_1) = 0$ and therefore, since \mathcal{E} specializes to \mathcal{E}_1 , also that $H^0(\mathcal{E}) = 0$. It follows by Claim 8.7 that $H^0(N_{X'} \otimes \mathcal{L}) = 0$ and then $H^0(N_X \otimes \omega_X(-2)) = 0$ by semicontinuity. Hence $N_X^*(3)$ is Ulrich and, clearly, X is not subcanonical. \square

REFERENCES

- [ALY] A. Atanasov, E. Larson, D. Yang. *Interpolation for normal bundles of general curves*. Mem. Amer. Math. Soc. **257** (2019), no. 1234, 105 pp. [2](#), [7](#), [9](#), [10](#)
- [B] A. Beauville. *An introduction to Ulrich bundles*. Eur. J. Math. **4** (2018), no. 1, 26-36. [1](#)
- [BE] E. Ballico, Ph. Ellia. *Some more examples of curves in \mathbb{P}^3 with stable normal bundle*. J. Reine Angew. Math. **350** (1984), 87-93. [1](#), [6](#), [7](#)
- [BMPT] V. Benedetti, P. Montero, Y. Prieto Montañez, S. Troncoso. *Projective manifolds whose tangent bundle is Ulrich*. J. Algebra **630** (2023), 248-273. [1](#)
- [C] G. Casnati. *Special Ulrich bundles on non-special surfaces with $p_g = q = 0$* . Internat. J. Math. **28** (2017), no. 8, 1750061, 18 pp. [6](#)
- [CH] M. Casanellas, R. Hartshorne. *Stable Ulrich bundles*. With an appendix by F. Geiss, F.-O. Schreyer. Internat. J. Math. **23** (2012), no. 8, 1250083, 50 pp. [6](#)
- [CPJLLV] K. Cook-Powell, D. Jensen, E. Larson, H. Larson, I. Vogt. *The embedding theorem in Hurwitz-Brill-Noether Theory*. Preprint 2023, arXiv:2303.15189. [9](#)
- [CMRPL] L. Costa, R. M. Miró-Roig, J. Pons-Llopis. *Ulrich bundles*. De Gruyter Studies in Mathematics, **77**, De Gruyter 2021. [1](#)
- [E] L. Ein. *Vanishing theorems for varieties of low codimension*. In: Algebraic geometry (Sundance, UT, 1986), 71-75, Lecture Notes in Math., **1311**, Springer, Berlin, 1988. [4](#), [5](#)
- [ELHi] G. Ellingsrud, A. Hirschowitz. *Sur le fibré normal des courbes gauches*. C. R. Acad. Sci. Paris Sér. I Math. **299** (1984), no. 7, 245-248. [1](#), [7](#)
- [EiHa] D. Eisenbud, J. Harris. *Divisors on general curves and cuspidal rational curves*. Invent. Math. **74** (1983), no. 3, 371-418. [9](#)
- [EL] G. Ellingsrud, D. Laksov. *The normal bundle of elliptic space curves of degree 5*. In: 18th Scandinavian Congress of Mathematicians (Aarhus, 1980), pp. 258-287, Progr. Math., **11**, Birkhäuser, Boston, Mass., 1981. [1](#), [7](#)
- [ES] D. Eisenbud, F.-O. Schreyer. *Resultants and Chow forms via exterior syzygies*. J. Amer. Math. Soc. **16** (2003), no. 3, 537-579. [1](#)
- [L] A. F. Lopez. *On varieties with Ulrich twisted normal bundles*. Preprint 2022 (arXiv:2205.06602). To appear on Perspectives on four decades of Algebraic Geometry: in Memory of Alberto Collino. Progress in Mathematics, volume 352, Birkhäuser. [1](#)
- [LR] A. F. Lopez, D. Raychaudhury. *On varieties with Ulrich twisted tangent bundles*. Ann. Mat. Pura Appl. (4) **203** (2024), no. 3, 1159-1193. [1](#), [3](#)
- [LV] E. Larson, I. Vogt. *Interpolation for Brill-Noether curves*. Forum Math. Pi **11** (2023), Paper No. e25, 90 pp. [2](#), [7](#), [8](#), [9](#), [10](#)

VINCENZO ANTONELLI, DIPARTIMENTO DI SCIENZE MATEMATICHE, POLITECNICO DI TORINO, C.SO DUCA DEGLI ABRUZZI 24, 10129 TORINO, ITALY. EMAIL: vincenzo.antonelli@polito.it

GIANFRANCO CASNATI, DIPARTIMENTO DI SCIENZE MATEMATICHE, POLITECNICO DI TORINO, C.SO DUCA DEGLI ABRUZZI 24, 10129 TORINO, ITALY. EMAIL: gianfranco.casnati@polito.it

ANGELO FELICE LOPEZ, DIPARTIMENTO DI MATEMATICA E FISICA, UNIVERSITÀ DI ROMA TRE, LARGO SAN LEONARDO MURIALDO 1, 00146, ROMA, ITALY. E-MAIL angelo.lopez@uniroma3.it

DEBADITYA RAYCHAUDHURY, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ARIZONA, 617 N SANTA RITA AVE., TUCSON, AZ 85721, USA. EMAIL: draychaudhury@math.arizona.edu

Da: Gregory G. Smith (via EditFlow) editflow+ggsmith@ams.msp.org 

Oggetto: Proc AMS 240610-orr86 - Accepted

Data: 13 giugno 2024 alle ore 16:13

A: vincenzo.antonelli@polito.it, gianfranco.casnati@polito.it, draychaudhury@math.arizona.edu, angelo.lopez@uniroma3.it

Cc: ggsmith@mast.queensu.ca

GS

Dear Professor Lopez,

I am pleased to inform you that your article

On varieties with Ulrich twisted conormal bundles
by Vincenzo Antonelli, Gianfranco Casnati, Angelo Felice Lopez and Debaditya Raychaudhury

has been accepted by the editorial board for publication in the Proceedings of the American Mathematical Society.

The PAMS editorial board accepts articles to be published in either the traditional PAMS, which is a green open access subscription journal, or PAMS, Series B, which is a gold open access journal supported by Article Processing Charges (APCs).

Now that your article has been accepted, please choose between the two publication options after consultation with your coauthors (if any) using the decision link near the bottom of this message.

To help inform your decision, information about AMS Open Access Journals, including APC and license information, is available at <http://www.ams.org/open-access>

As corresponding author, you are the only author receiving this message and are authorized to indicate the decision that you and your coauthors (if any) have made.

PLEASE INDICATE YOUR DECISION USING THIS FORM:

<https://www.ams.org/cgi-bin/ctpauthor/oachoice?m=122340&a=1075979&y=339C60347D>

If you have a need to correspond with the AMS about your decision, please send mail to openaccess@ams.org.

For any changes in your postal or email addresses, please keep the AMS informed at journals-query@ams.org.

Sincerely,

Gregory G. Smith

Sent via EditFlow by

Gregory G. Smith <ggsmith@mast.queensu.ca>

