

Wind tunnel study on the influence of vegetation density and wind direction on urban canyon ventilation

Original

Wind tunnel study on the influence of vegetation density and wind direction on urban canyon ventilation / Del Ponte, Annikavittoria; Fellini, Sofia; Marro, Massimo; Salizzoni, Pietro; Ridolfi, Luca. - (2024). (Intervento presentato al convegno EGU General Assembly 2024 tenutosi a Vienna, Austria nel 15-04-2024 al 19-04-2024).

Availability:

This version is available at: 11583/2990461 since: 2024-07-07T15:43:09Z

Publisher:

EGU General Assembly 2024

Published

DOI:

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)



Wind tunnel study on the influence of vegetation density and wind direction on urban canyon ventilation

Annika Vittoria Del Ponte^{1,2}, Sofia Fellini¹, Massimo Marro², Pietro Salizzoni², and Luca Ridolfi¹

¹Politecnico di Torino, Dipartimento di ingegneria per l'ambiente, il territorio e le infrastrutture (DIATI), Turin, Italy

²Univ Lyon, Ecole Centrale de Lyon, CNRS, Univ Claude Bernard Lyon 1, INSA Lyon, LMFA UMR5509, 69130, Ecully, France

Inserting vegetation within the urban environment mitigates the urban heat island effect, the flooding risk, and improves air quality. However, its aerodynamic effect has remarkable impact on the pollutant transport and, consequently, on human health comfort. Indeed, the presence of vegetation within an urban canyon leads to non-trivial patterns of pollutant concentration and mass fluxes, as a consequence of complex mean and turbulent velocity fields. In addition to the vegetation density, the flow structure within canyons is influenced by their geometry and by the wind direction.

The aim of the present study is to experimentally investigate the velocity field within a canyon, varying the vegetation density and the wind direction. We measured flow velocity statistics within an indefinitely long street canyon, with unit height-to-width ratio, subject to a neutrally stratified boundary layer modeled in the wind tunnel of École Centrale de Lyon. The aerodynamic impact of vegetation was reproduced by inserting plastic miniatures of trees along the two long sides of the canyon. We considered an empty canyon and a vegetated canyon, whose longitudinal axes are oriented with angles of 0°, 30°, and 60° with respect to the external wind flow.

Results reveal that when the canyon is inclined with respect to the external wind direction the mean flow follows a complex helicoidal structure. The presence of trees decreases significantly the mean longitudinal velocity and weakens the transversal circulation in the inclined canyon. The dampening effect of the mean longitudinal flow is more marked increasing the inclination angle of the canyon. Turbulent fluctuations are enhanced above the tree crowns, mostly when the wind blows parallel to the canyon axis. On the contrary, turbulent fluctuations decreases at tree trunk and crown levels, in particular when the canyon is inclined of 60° with respect to the external wind direction. Spectra of the velocity signal show that the presence of trees induces an evident shift of the energy peak towards high frequencies.

The collected data constitute a step forward to understand and modeling the urban microclimate.