

A new method to prioritize the QFDs' engineering characteristics inspired by the Law of Comparative Judgment

Original

A new method to prioritize the QFDs' engineering characteristics inspired by the Law of Comparative Judgment / Maisano, Domenico A.; Carrera, Giovanna; Mastrogiacomo, Luca; Franceschini, Fiorenzo. - In: RESEARCH IN ENGINEERING DESIGN. - ISSN 0934-9839. - 35:(2024), pp. 343-353. [10.1007/s00163-024-00436-8]

Availability:

This version is available at: 11583/2992213 since: 2024-10-19T16:45:57Z

Publisher:

Springer

Published

DOI:10.1007/s00163-024-00436-8

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)



A new method to prioritize the QFDs' engineering characteristics inspired by the Law of Comparative Judgment

Domenico A. Maisano¹ · Giovanna Carrera¹ · Luca Mastrogiacomo¹ · Fiorenzo Franceschini¹

Received: 27 July 2023 / Revised: 21 February 2024 / Accepted: 30 March 2024 / Published online: 5 May 2024
© The Author(s) 2024

Abstract

The primary module of *Quality Function Deployment* (QFD) is the *House of Quality* (HoQ), which supports the design of new products and services by translating *customer requirements* (CRs) into *engineering characteristics* (ECs). Within the HoQ framework, the traditional technique for prioritizing ECs is the *independent scoring method* (ISM), which aggregates the weights of the CRs and the relationships between CRs and ECs (i.e., *null*, *weak*, *medium*, and *high*) through a weighted sum. However, ISM incorporates two questionable operations: (i) an arbitrary numerical conversion of the relationships between CRs and ECs, and (ii) the “promotion” of these relationships from *ordinal* to *cardinal* scale. To address these conceptual shortcomings, this paper introduces a novel procedure for prioritizing ECs, inspired by the Thurstone's *Law of Comparative Judgment* (LCJ). This procedure offers a solution that is conceptually sound and practical, overcoming the conceptual shortcomings of ISM, while maintaining its simplicity, flexibility, and ease of implementation. The proposed approach is supported by a realistic application example illustrating its potential.

Keywords House of quality · Engineering characteristic · Prioritization · Law of comparative judgment · Scale promotion · Scale anchoring

Abbreviations

CR	Customer Requirement
EC	Engineering Characteristic
HoQ	House of Quality
ISM	Independence Scoring Method
LCJ	Law of Comparative Judgment
MCDM	Multi-Criteria Decision Making
QFD	Quality Function Deployment
VoC	Voice of the Customer

1 Introduction

Quality Function Deployment (QFD) is a commonly used methodology for facilitating customer-oriented design of new products and services (Franceschini 2001). The initial module of QFD, known as the *House of Quality* (HoQ), aims to translate the *customer requirements* (CRs) of a new product/service into *engineering characteristics* (ECs), i.e., technical design features that directly impact the CRs (Zare Mehrjerdi 2010). Figure 1a exemplifies the HoQ's *relationship matrix* with CRs and ECs indicated in rows and columns; the symbols in Fig. 1b represent the intensities of the relationships between ECs and CRs. Additionally, a weight (w_{CR_i}) is assigned to each i -th CR, reflecting its importance from three complementary perspectives: *final customer*, *corporate brand image*, and *improvement goals* for the new product/service in relation to existing counterparts in the market (Akao 1994; Franceschini and Maisano 2018).

Several actors participate in the QFD construction. The leading role is played by the members of the *QFD team*, i.e., a working group consisting of experts from the company of interest, with complementary skills ranging from marketing, design, quality, production, maintenance, etc. Another important role is played by a sample of interviewed

✉ Fiorenzo Franceschini
fiorenzo.franceschini@polito.it

Domenico A. Maisano
domenico.maisano@polito.it

Giovanna Carrera
giovanna.carrera@polito.it

Luca Mastrogiacomo
luca.mastrogiacomo@polito.it

¹ DIGEP Dept. of Management and Production Engineering,
Politecnico Di Torino, Corso Duca Degli Abruzzi 24,
10129 Turin, Italy

Fig. 1 **a** Relationship matrix and EC prioritization using the independence scoring method (ISM). **b** Intensity of the relationships between ECs and CRs, and conventional conversion into numerical coefficients (r_{ij})

(a) Relationship matrix					(b) Intensity of relationships between ECs and CRs		
	EC_1	EC_2	EC_3	w_{CR_i}	Intensity	Symbol	r_{ij}
CR_1	○	△		30%	None	∅	0
CR_2	△	●	△	25%	Low	△	1
CR_3			●	25%	Medium	○	3
CR_4			○	20%	High	●	9
w_{EC_j}	1.15	2.55	3.10				

respondents—i.e., (potential) end users who contribute to the collection of the so-called *voice of the customer* (VoC)—from which the QFD team extrapolates the HoQ’s CRs (Akao 1994; Franceschini and Rossetto 1995). To effectively identify the VoC, which is foundational to the whole QFD process, it is needed to meticulously select interviewees. This selection should be based on both the quantity and the quality of participants, ensuring that they possess a comprehensive understanding of the product or service under consideration, even if not from a technical standpoint. Concurrently, the QFD team members are expected to bring their specialized technical knowledge to the table, facilitating collaborative decision-making throughout the design and development stages. This approach, as outlined by Akao (1994) and Huang et al. (2022), is integral to the traditional HoQ construction method, which aims to equitably distribute tasks among all involved parties. Thus, while there may be room for enhancing the traditional HoQ, it would be ill-advised to alter the fundamental (largely manual) methods of data gathering that are familiar to the above-mentioned groups. This stance, however, does not preclude the possibility of refining the data processing stage, provided that the initial data collection methods remain unchanged.

Returning to the traditional QFD construction method, ECs should be prioritized considering their impact on the final customer; in general, ECs that have relatively intense relationships with CRs characterized by relatively high w_{CR_i} values deserve more attention during the design phase. The conventional approach for prioritizing ECs is the so-called *independence scoring method* (ISM), which is a weighted sum of the coefficients (r_{ij}) derived from the numerical conversion of the relationship matrix symbols (see Fig. 1b), utilizing the w_{CR_i} values as weights (Akao 1994). The resulting EC weight can be determined by Eq. 1, as exemplified in the lower part of Fig. 1a:

$$w_{EC_j} = \sum_{vi} (r_{ij} \cdot w_{CR_i}) \tag{1}$$

However, the ISM method incorporates two conceptually questionable operations: (i) an inherently arbitrary

conversion of the relationship intensities (defined on an *ordinal scale*: $\emptyset < \Delta < \circ < \bullet$) into conventional numerical coefficients (r_{ij} : $\emptyset \rightarrow 0, \Delta \rightarrow 1, \circ \rightarrow 3, \bullet \rightarrow 9$, cf. Figure 1b), and (ii) the aggregation of r_{ij} values through a weighted sum, which introduces an (undue) promotion to a *cardinal scale* with meaningful intervals (Franceschini et al. 2015). These questionable transformations can result in a distorted prioritization of ECs (Lyman 1990).

To address these conceptual shortcomings, this paper proposes a novel technique for prioritizing ECs based on Thurstone’s *Law of Comparative Judgment* (LCJ) (Thurstone 1927). Some advantages of the LCJ are that this technique is well-established, conceptually rigorous, effective and robust in practice (Maranell, 1974; Brown and Peterson 2009; Kelly et al. 2022). The integration of the LCJ into the new procedure and the integration of some “dummy” ECs (see Sect. 5) will enable the prioritization of ECs on a *cardinal scale* (Franceschini et al. 2022).

The remainder of this article is organized into five sections. Section 2 contains a brief review of the state-of-art techniques for prioritizing ECs that are alternative to ISM. Section 3 illustrates a case study that will accompany the description of the new procedure. Section 4 briefly recalls the LCJ, illustrating its underlying assumptions and practical application. Section 5 provides a step-by-step description of the proposed procedure, accompanied by an application example referring to the case study introduced in Sect. 3. The concluding section summarizes the major contributions of this research, its practical implications, limitations and insights for future development.

2 Literature review

Although ISM is the most widely used technique for EC prioritization, it has some shortcomings documented in a plurality of scientific contributions. In addition to (i) the arbitrary numerical conversion of the relationship matrix coefficients and (ii) the undue promotion of their *ordinal scale* into a *cardinal* one (cf. Figure 1b), there are other

Fig. 2 Relationship matrix referring to the design of a new climbing safety harness model, adapted from (Franceschini et al. 2015). The w_{EC_j} values are determined by converting the relationship matrix symbols into numerical coefficients (r_{ij} , cf. Figure 1b), and applying the ISM (Eq. 1)

		Meets safety standards	Harness weight	Webbing strength	No. of colours	No. of sizes	Padding thickness	No. of buckles	No. of gear loops	w_{CR_i}
		EC ₁	EC ₂	EC ₃	EC ₄	EC ₅	EC ₆	EC ₇	EC ₈	
Easy to put on	CR ₁					○		●		7%
Comfortable when hanging	CR ₂					○	●	○		20%
Fits over different clothes	CR ₃					○	○	●		3%
Accessible gear loops	CR ₄								●	12%
Does not restrict movement	CR ₅		○			○	●	○		27%
Lightweight	CR ₆		●	○			○	△	△	10%
Safe	CR ₇	●	○	●						14%
Attractive	CR ₈		△		●		△	△		7%

w_{EC_j}	1.26	2.20	1.56	0.63	1.71	4.69	2.48	1.18
------------	------	------	------	------	------	------	------	------

shortcomings, such as the fact that EC weights may not be consistent with CR weights. The latter shortcoming has inspired a corrective normalization (Lyman 1990) that, however, does not resolve the previous two.

The scientific literature includes multiple alternative techniques that aim to overcome the weaknesses of ISM and expand its scope. At the risk of oversimplifying, these techniques can be synthesized into three macro-categories (Franceschini et al. 2022).

1. *Rule-based techniques.* Practical techniques achieve a computationally simple, intuitive, and satisfactory prioritization of ECs, which is not necessarily the optimal one (assuming it exists). In some operational contexts, these techniques are classified as “heuristic” or “experience-based”. Let us recall, for example, Borda’s method or other techniques based on pairwise comparisons (Dym et al. 2002), and another technique (called *ordinal prioritization method*) based on the adaptation of a model proposed by Yager to the HoQ context (Franceschini et al. 2015; Galetto et al. 2018; Yager 2001).
2. *Multi-criteria decision-making (MCDM) techniques.* EC prioritization can be also seen as an MCDM problem involving conflicting CRs. Several MCDM techniques have been used in previous studies to improve the performance of QFD (Huang et al. 2022; Ping et al. 2020). Let us recall for example the application of the ELECTRE-II (*Élimination Et Choix Traduisant la Réalité*) method (Liu and Ma 2021) or the EDAS approach (*evaluation based on distance from average solution*) (Ghorabae et al. 2015; Mao et al. 2021).
3. *Techniques based on fuzzy logic.* These techniques, which could also be interpreted as advanced heuristics, take into account the inherent uncertainty in the formulation of CR-importance judgments (by interviewed

respondents) or in the formulation of relationship matrix intensities (by QFD team members). For example, the techniques proposed by Li et al. (2019) and Shi et al. (2022) apply to the *open design* contest, in which the intensity of relationships between CRs and ECs and the prioritization of ECs are inferred on the basis of linguistic variables and text-format information, describing the subjective imprecision of human cognition.

Most of the aforementioned techniques require additional information beyond that collected in the traditional QFD process. For example, techniques to handle information related to linguistic variables require verbatim transposition of interviews that were conducted in open-ended form (Li et al. 2019); most of the techniques based on fuzzy logic require some technical expertise in setting working parameters, such as thresholds or weights. The procedure proposed in this paper—which can be referred to as “distribution-based” since it relies on statistical assumptions regarding the distribution of the QFD team preferences (Franceschini et al. 2022)—fits the information contained in the classic HoQ and can also be implemented and automated by non-experts (cf. Section 5).

3 Test case

Referring to an application example adapted from the scientific literature (Franceschini et al. 2015), let us consider a company of mountain sports accessories that plans to design a new model of climbing safety harness through the QFD. Figure 2 illustrates the HoQ that was constructed by combining information obtained through interviews with a sample of respondents (i.e., potential end users) and from the




Fig. 3 Theoretical distributions of the position of two generic objects (i.e., EC_j and EC_k) in the psychological continuum. **b** Graphical representation of the quantity $(1 - p_{jk})$, being $p_{jk} = P[(EC_j - EC_k) \geq 0]$

technical examination of QFD team members (i.e., corporate staff with technical and design expertise). Eight CRs and eight ECs were identified.¹ The w_{CR_i} values inherently may take into account information concerning the CR importance to final customers, corporate brand image, and improvement goals, combining them through a multiplicative aggregation model (Franceschini 2001; Franceschini and Maisano 2018). For simplicity, Fig. 2 shows only the w_{CR_i} values, omitting the three CR-importance contributions mentioned above. Next, ECs are prioritized, determining the w_{EC_j} values reported in the lower part of Fig. 2 through the ISM (cf. Equation 1). This HoQ will be used as a test case to illustrate the new EC prioritization procedure step-by-step (cf. Section 5). The concluding part compares the results of the new prioritization procedure with those derived from the traditional ISM.

4 Basics of the law of comparative judgment

The very general problem in which Thurstone’s LCJ finds application is summarized as follows: a set of experts formulate their individual (subjective) judgments about a specific attribute of some objects and these judgments must be merged into a collective one (Maranell, 1974; Kelly et al. 2022). The attribute can be defined as “a specific feature of

objects, which evokes a subjective response in each expert”. Consider, for example, the intensity of the aroma (attribute) of some alternative coffee blends (objects), which are assessed by a panel of experienced café customers (experts).

In this scenario, Thurstone (1927) postulated the existence of a psychological continuum, i.e., “an abstract and unknown unidimensional scale, in which the position of the objects is directly proportional to their degree of the attribute of interest”. Although the psychological continuum is a unidimensional imaginary scale, the LCJ can be used to approximate the position of the objects of interest on it. According to the so-called case V of Thurstone’s LCJ, the position of a generic j -th object (EC_j) is in fact postulated to be distributed normally: $EC_j \sim N(\mu_{EC_j}, \sigma_{EC_j}^2)$, where μ_{EC_j} and $\sigma_{EC_j}^2$ are the unknown mean value and variance of that object’s attribute.² Figure 3 represents the hypothetical distributions of the position of any two generic objects, EC_j and EC_k . The distribution associated with a given object is characterized by a dispersion (or variance), which reflects the intrinsic expert-to-expert variability in positioning (albeit indirectly, as we will better understand below) that object on the psychological continuum. Let μ_{EC_j} and μ_{EC_k} correspond to the (unknown) expected values of the two objects and $\sigma_{EC_j}^2$ and $\sigma_{EC_k}^2$ the (unknown) variances. The difference $(EC_j - EC_k)$ will follow a normal distribution with parameters:

² This notation will facilitate understanding the description in Sect. 5, since the objects of the problem of interest are the HoQ’s ECs.

¹ The fact that the two quantities coincide is purely incidental.

$$\begin{aligned} \mu_{(EC_j - EC_k)} &= \mu_{EC_j} - \mu_{EC_k} \text{ and } \sigma_{(EC_j - EC_k)} \\ &= \sqrt{\sigma_{EC_j}^2 + \sigma_{EC_k}^2 - 2 \cdot \rho_{EC_j, EC_k} \cdot \sigma_{EC_j} \cdot \sigma_{EC_k}} \end{aligned} \tag{2}$$

where:

μ_{EC_j} and μ_{EC_k} denote the (unknown) mean values of EC_j and EC_k in the psychological continuum;

$\sigma_{EC_j}^2$ and $\sigma_{EC_k}^2$ denote the (unknown) variances of EC_j and EC_k ;

ρ_{EC_j, EC_k} denotes the (unknown) correlation between objects EC_j and EC_k .

Considering the area subtended by the distribution of $(EC_j - EC_k)$, let us draw a vertical line passing through the point with $EC_j - EC_k = 0$ (see Fig. 3b). The area to the right of the line depicts the observed proportion of times (p_{jk}) that $EC_j \geq EC_k$ or $(EC_j - EC_k) \geq 0$. Of course, the area to the left depicts the complementary proportion $(1 - p_{jk})$.

In addition, it is postulated that the variances of the objects are all equal ($\sigma_{EC_1}^2 = \sigma_{EC_2}^2 = \dots = \sigma^2$) and the inter-correlations (in the form of Pearson coefficients ρ_{EC_j, EC_k}) between pairs of objects (EC_j, EC_k) are all equal too ($\rho_{EC_j, EC_k} = \rho, \forall j, k$).

Having outlined this framework, let us now focus on the LCJ's application, which is based on the following four steps:

1. A set of *experts* (j_1, j_2, \dots) formulate their preferences for each object (EC_j) versus every other object (EC_k), considering all possible $C_2^m = m \cdot (m - 1) / 2$ pairs, m being the total number of objects. In practice, the following question needs to be answered: “How do you judge the degree of the attribute of EC_j compared to that of EC_k in relative terms?”. Judgments are expressed through relationships of *strict preference* (e.g., “ $EC_1 > EC_2$ ” or “ $EC_1 < EC_2$ ”) or *indifference* (e.g., “ $EC_1 \sim EC_2$ ”). Results are then aggregated into a proportion matrix (\mathbf{P}). Precisely, for each possible paired comparison between two objects (EC_j and EC_k), the conventional portion of experts who prefer the first object to the second one is determined as:

$$p_{jk} = 1 \cdot p_{jk}^{(>)} + 1/2 \cdot p_{jk}^{(\sim)} + 0 \cdot p_{jk}^{(<)}, \in [0, 1], \tag{3}$$

$p_{jk}^{(>)}$ being the portion of experts for whom $EC_j > EC_k$,

$p_{jk}^{(\sim)}$ being the portion of experts for whom $EC_j \sim EC_k$,

$p_{jk}^{(<)}$ being the portion of experts for whom $EC_j < EC_k$.

It can be noted that p_{jk} is the same quantity represented in Fig. 3b. The LCJ allows for its quantification on the basis of the (empirical) expert judgments.

The coefficient “1/2” multiplying $p_{jk}^{(\sim)}$ conventionally weighs the indifference relationship as an intermediate coefficient between that related to the “favourable”

(“ $EC_j > EC_k$ ”) strict preference relation (with coefficient “1”) and that related to the “unfavourable” (“ $EC_j < EC_k$ ”) strict preference relation (with coefficient “0”). Taking into account that $p_{jk}^{(\sim)} = p_{kj}^{(\sim)}$, this type of weighting guarantees the complementarity relationship:

$$p_{jk} = 1 - p_{kj}, \tag{4}$$

which can be demonstrated by combining Eq. 3 with the two following relationships:

$$p_{jk}^{(>)} + p_{jk}^{(\sim)} + p_{jk}^{(<)} = 1, \tag{5}$$

2. Next, p_{jk} values are transformed into z_{jk} values, through the relationship:

$$z_{jk} = \Phi^{-1}(1 - p_{jk}) \tag{6}$$

3. $\Phi^{-1}(\cdot)$ being the reverse of the cumulative distribution function (Φ) of the standard normal distribution. The element z_{jk} represents a unit normal deviate, which will be positive for all values of $(1 - p_{jk})$ over 0.50 and negative for all values of $(1 - p_{jk})$ under 0.50.
4. In general, objects are judged differently by experts. However, if all experts express the same preference for each outcome, the model is no more viable (p_{jk} values of 1.00 and 0.00 would correspond to z_{jk} values of $\pm\infty$). A simplified approach for tackling this problem is to associate those p_{jk} values ≤ 0.00135 with $z_{jk} = \Phi^{-1}(1 - 0.00135) = 3$ and those p_{jk} values ≥ 0.99865 with $z_{jk} = \Phi^{-1}(1 - 0.99865) = -3$. More sophisticated solutions to deal with this issue have been proposed (Franceschini et al. 2022).

Next, the z_{jk} values related to the possible paired comparisons are reported into a matrix \mathbf{Z} . The element z_{jk} is reported in the j -th row and k -th column. The relationship $z_{kj} = -z_{jk}$ holds, being unit normal deviates related to complementary cumulative probabilities (cf. Equation 4).

A *scaling* can be performed by (i) summing the values into each column of the matrix \mathbf{Z} and (ii) dividing these sums by m . It can be demonstrated that the result obtained for each k -th column (x_k) corresponds to the unknown average value (μ_{EC_k}) of the k -th object's attribute, up to a positive scale factor and an additive constant: $x_k = \sum_k(z_{jk}) / m = c_1 \cdot \mu_{EC_k} + c_2$. In other words, the LCJ results into an *interval scaling*, i.e., objects are defined on a scale (x) with arbitrary zero point and unit (Thurstone 1927; Franceschini et al. 2022).

5 Methodological approach

The LCJ can be adapted for prioritizing the ECs within the HoQ context. In this case, the focus is on the ECs (i.e., *objects*), which need to be prioritized according to the degree

(a) Relationship matrix				(b) EC rankings	
	EC_1	EC_2	EC_3	Without dummy ECs	With dummy ECs
CR_1	●			$\rightarrow EC_1 > (EC_2 \sim EC_3)$	$(EC_{\bullet} \sim EC_1) > EC_{\circ} > EC_{\Delta} > (EC_{\emptyset} \sim EC_2 \sim EC_3)$
CR_2	○	△	△	$\rightarrow EC_1 > (EC_2 \sim EC_3)$	$EC_{\bullet} > (EC_{\circ} \sim EC_1) > (EC_{\Delta} \sim EC_2 \sim EC_3) > EC_{\emptyset}$
...					

Fig. 4 Transformation of a relationship matrix into EC rankings, with or without dummy ECs (i.e., EC_{\bullet} , EC_{\circ} , EC_{Δ} , and EC_{\emptyset})

of intensity (i.e., *attribute*) of their relationships with each specific CR. Unlike the traditional application of LCJ, where individual experts formulate paired comparisons of objects, in this context paired comparisons emerge indirectly from the HoQ’s relationship matrix. This approach represents a departure from the traditional LCJ method, since experts do not make individual judgements that are then aggregated. Instead, in the new procedure, several expert judgements—derived from the collective compilation of the relationship matrix by the QFD team members (Zare Mehrjerdi 2010)—are then aggregated through the LCJ. This point is clarified in the explanation below, which is structured in four steps (*a*, *b*, *c* and *d*).

(a) *Transformation of relationship matrix into rankings* Focusing on the *i*-th specific row of the relationship matrix, a ranking of ECs can be determined according to the degree of their relationships with the *i*-th CR. For example, with reference to the relationship matrix in Fig. 2 and to “ CR_6 —Lightweight”, the following ranking can be obtained: •

$$\begin{aligned}
 (EC_{\bullet} \sim EC_2) &> (EC_{\circ} \sim EC_3 \sim EC_6) \\
 &> (EC_{\Delta} \sim EC_7 \sim EC_8) > (EC_{\emptyset} \sim EC_1 \sim EC_4 \sim EC_5)
 \end{aligned}
 \tag{7}$$

In addition to the “regular” ECs, i.e., EC_1 to EC_8 , the above ranking also includes four “dummy” or “anchor” ECs, i.e., EC_{\bullet} , EC_{\circ} , EC_{Δ} , and EC_{\emptyset} , which represent the degree of intensity of the relationships expressed in the relationship matrix, espresso in absolute terms (“None” $\rightarrow \emptyset$, “Low” $\rightarrow \Delta$, “Medium” $\rightarrow \circ$, and “High” $\rightarrow \bullet$; cf. Figure 1b). The introduction of these objects in the rankings, which by their nature are formulated in *relative* terms, is necessary in order not to lose part of the available information content: that is, the degree of intensity of the relationships in *absolute* terms. For example, referring to the new simplified relationship matrix in Fig. 4 (which is different from that in the test case in Fig. 2), the relationships of the three ECs (EC_1 , EC_2 and EC_3) with CR_1 (i.e., $EC_1 \rightarrow \bullet, EC_2 \rightarrow \emptyset, EC_3 \rightarrow \emptyset$) result into the ranking $EC_1 > (EC_2 \sim EC_3)$, which would be identical to that obtained by considering the relationships of the same ECs with CR_2 (i.e., $EC_1 \rightarrow \circ, EC_2 \rightarrow \Delta, EC_3 \rightarrow \Delta$). In this case, the transformation into *relative* rankings leads to losing information about the *absolute* degree of intensity

of individual relationships, which distinguishes the two configurations.

Referring to the LCJ framework, similarly to regular ECs, dummy ECs are assumed to project a normal distribution on the psychological continuum, with unknown mean value and unknown variance, equal to that of the other objects (cf. Section 4) (Franceschini and Maisano 2019). The dummy objects are also used to perform the so-called “anchoring” of the scaling resulting from the LCJ, as described below.

(b) *Transformation of rankings into paired comparison relationships.* Each ranking can be uniquely translated into paired comparisons. Figure 5 exemplifies this process for each CR in the relationship matrix. Since the test case includes twelve total ECs—i.e., eight regular (EC_1 to EC_8) and four dummy ones ($EC_{\bullet}, EC_{\circ}, EC_{\Delta}$, and EC_{\emptyset})—the total paired comparisons are $C_2^{12} = \binom{12}{2} = \frac{12 \cdot 11}{2} = 66$.

(c) *LCJ application.* Next, a proportion (p_{jk}) must be associated with each *jk*-th paired comparison. In the traditional LCJ (cf. Section 4), p_{jk} reflects the portion of expert population that expressed a preference of object EC_j over object EC_k ³. Although experts do not directly express their preferences between pairs of ECs, for each CR in the relationship matrix (which was collectively constructed by the QFD team members), an EC ranking can be determined as illustrated in Fig. 4; then, this ranking can be decomposed into paired comparisons between ECs.

Additionally, each CR in the HoQ corresponds to a certain percentage weight (i.e., w_{CR_i} , being $\sum_{v_i} w_{CR_i} = 1$), which describes its importance from different points of view (cf. Section 3). This weight can also be associated with the relative EC ranking and the paired comparison relationships resulting from it. For a given (*jk*-th) comparison between pairs of ECs, there are therefore as many paired comparison relationships as the number of CRs (with their associated weights). Figure 5 exemplifies the process of constructing paired comparison relationships from the relationship matrix in Fig. 2.

Adapting the LCJ (cf. Section 4), the p_{jk} proportions can be determined by aggregating the preceding comparisons by means

³ This preference combines the relationships of *strict preference* and *indifference* according to the conventional weighting in Eq. 3.

(a) Relationship matrix

	EC ₁	EC ₂	EC ₃	EC ₄	EC ₅	EC ₆	EC ₇	EC ₈
CR ₁					○		●	
CR ₂					○	●	○	
CR ₃					○	○	●	
CR ₄								●
CR ₅		○			○	●	○	
CR ₆		●	○			○	△	△
CR ₇	●	○	●					
CR ₈		△		●		△	△	

(b) Rankings (including dummy ECs: EC_●, EC_○, EC_△ and EC_◊)

- CR₁: (EC_●~EC₇) > (EC_○~EC₅) > EC_△ > (EC_◊~EC₁~EC₂~EC₃~EC₄~EC₆~EC₈)
- CR₂: (EC_●~EC₆) > (EC_○~EC₅~EC₇) > EC_△ > (EC_◊~EC₁~EC₂~EC₃~EC₄~EC₈)
- CR₃: (EC_●~EC₇) > (EC_○~EC₅~EC₆) > EC_△ > (EC_◊~EC₁~EC₂~EC₃~EC₄~EC₈)
- CR₄: (EC_●~EC₈) > EC_○ > EC_△ > (EC_◊~EC₁~EC₂~EC₃~EC₄~EC₅~EC₆~EC₇)
- CR₅: (EC_●~EC₆) > (EC_○~EC₂~EC₅~EC₇) > EC_△ > (EC_◊~EC₁~EC₃~EC₄~EC₈)
- CR₆: (EC_●~EC₂) > (EC_○~EC₃~EC₆) > (EC_△~EC₇~EC₈) > (EC_◊~EC₁~EC₄~EC₅)
- CR₇: (EC_●~EC₁~EC₃) > (EC_○~EC₂) > EC_△ > (EC_◊~EC₄~EC₅~EC₆~EC₇~EC₈)
- CR₈: (EC_●~EC₄) > EC_○ > (EC_△~EC₂~EC₆~EC₇) > (EC_◊~EC₁~EC₃~EC₅~EC₈)

(c) Paired-comparison relationships

EC pairs	CR ₁	CR ₂	CR ₃	CR ₄	CR ₅	CR ₆	CR ₇	CR ₈
EC ₁ , EC ₂	~	~	~	~	<	<	>	<
EC ₁ , EC ₃	~	~	~	~	~	<	~	~
EC ₁ , EC ₄	~	~	~	~	~	~	~	~
EC ₁ , EC ₅	<	<	<	~	<	~	>	~
EC ₁ , EC ₆	~	<	<	~	<	<	>	<
EC ₁ , EC ₇	<	<	<	~	<	<	>	<
EC ₁ , EC ₈	~	~	~	~	~	<	>	~
EC ₁ , EC _◊	~	~	~	~	~	~	>	~
EC ₁ , EC _△	<	<	<	<	<	<	>	<
EC ₁ , EC _○	<	<	<	<	<	<	>	<
EC ₁ , EC _●	<	<	<	<	<	<	~	<
EC ₂ , EC ₃	~	~	~	~	>	>	<	<
EC ₂ , EC ₄	~	~	~	~	>	>	<	<
EC ₂ , EC ₅	<	<	<	~	>	>	>	>
EC ₂ , EC ₆	~	<	<	~	<	>	>	~
EC ₂ , EC ₇	<	<	<	~	~	>	>	~
EC ₂ , EC ₈	~	~	~	<	>	>	>	>
EC ₂ , EC _◊	~	~	~	~	>	>	>	>
EC ₂ , EC _△	<	<	<	<	>	>	>	~
EC ₂ , EC _○	<	<	<	<	~	>	~	<
EC ₂ , EC _●	<	<	<	<	<	~	<	<
EC ₃ , EC ₄	~	~	~	~	~	>	>	<
EC ₃ , EC ₅	<	<	<	~	<	>	>	~
EC ₃ , EC ₆	~	<	<	~	<	~	>	~
EC ₃ , EC ₇	<	<	<	~	<	>	>	<
EC ₃ , EC ₈	~	~	~	<	~	>	>	~
EC ₃ , EC _◊	~	~	~	~	~	>	>	~
EC ₃ , EC _△	<	<	<	<	<	>	>	<
EC ₃ , EC _○	<	<	<	<	<	~	>	<
EC ₃ , EC _●	<	<	<	<	<	<	~	<
EC ₄ , EC ₅	<	<	<	~	<	~	~	>
EC ₄ , EC ₆	~	<	<	~	<	<	~	>
EC ₄ , EC ₇	<	<	<	~	<	<	~	>
EC ₄ , EC ₈	~	~	~	~	>	>	>	>
EC ₄ , EC _◊	<	<	<	<	<	<	<	<
EC ₄ , EC _△	<	<	<	<	<	<	<	<
EC ₄ , EC _○	<	<	<	<	<	<	<	<
EC ₄ , EC _●	<	<	<	<	<	<	<	<
EC ₅ , EC ₆	>	<	~	~	<	<	~	<
EC ₅ , EC ₇	<	~	<	~	~	<	~	<
EC ₅ , EC ₈	>	>	>	<	>	<	~	~
EC ₅ , EC _◊	>	>	>	~	>	<	~	~
EC ₅ , EC _△	>	>	>	<	>	<	<	<
EC ₅ , EC _○	~	~	~	<	~	<	<	<
EC ₅ , EC _●	<	<	<	<	<	<	<	<
EC ₆ , EC ₇	<	>	<	~	>	>	~	~
EC ₆ , EC ₈	~	>	>	~	>	>	~	>
EC ₆ , EC _◊	~	>	>	~	>	>	~	>
EC ₆ , EC _△	<	>	>	<	>	>	<	~
EC ₆ , EC _○	<	>	~	<	>	~	<	<
EC ₆ , EC _●	<	~	<	<	<	<	<	<
EC ₇ , EC ₈	>	>	>	<	>	~	~	>
EC ₇ , EC _◊	>	>	>	~	>	>	~	>
EC ₇ , EC _△	>	>	>	<	>	~	<	~
EC ₇ , EC _○	>	~	>	<	~	<	<	<
EC ₇ , EC _●	~	~	~	<	<	<	<	<
EC ₈ , EC _◊	~	~	~	>	~	>	~	~
EC ₈ , EC _△	<	<	<	>	<	~	<	<
EC ₈ , EC _○	<	<	<	>	<	<	<	<
EC ₈ , EC _●	<	<	<	~	<	<	<	<
EC _◊ , EC _△	<	<	<	<	<	<	<	<
EC _◊ , EC _○	<	<	<	<	<	<	<	<
EC _◊ , EC _●	<	<	<	<	<	<	<	<
EC _△ , EC _○	<	<	<	<	<	<	<	<
EC _△ , EC _●	<	<	<	<	<	<	<	<
EC _○ , EC _●	<	<	<	<	<	<	<	<

Fig. 5 Transformation of the relationship matrix in (a) into EC rankings (b) and paired comparison relationships (c)

of a weighted sum; before showing this aggregation, for the sake of clarity, it is appropriate to take a step back.

With reference to each *i*-th CR and each *jk*-th paired comparison (i.e., EC_{*j*} versus EC_{*k*}), the following three binary coefficients are defined:

$$\begin{aligned}
 c_{i,jk}^{(>)} & \text{ being equal to 1 if } EC_j > EC_k \text{ (otherwise 0),} \\
 c_{i,jk}^{(\sim)} & \text{ being equal to 1 if } EC_j \sim EC_k \text{ (otherwise 0),} \\
 c_{i,jk}^{(<)} & \text{ being equal to 1 if } EC_j < EC_k \text{ (otherwise 0),}
 \end{aligned}
 \tag{8}$$

It can be seen that these coefficients are mutually exclusive and the complementarity relationship holds: $c_{i,jk}^{(>)} + c_{i,jk}^{(\sim)} + c_{i,jk}^{(<)} = 1$. A general coefficient expressing the degree of preference of EC_{*j*} over EC_{*k*} from the perspective of the *i*-th CR can be defined as:

$$c_{i,jk} = 1 \cdot c_{i,jk}^{(>)} + 1/2 \cdot c_{i,jk}^{(\sim)} + 0 \cdot c_{i,jk}^{(<)} \in \left\{ 0, 1/2, 1 \right\}
 \tag{9}$$

Note the similarity between Eqs. 3 and 9; by construction, it results that: $c_{i,jk} = 1 - c_{i,kj}$. Figure 6 exemplifies the

EC pairs	CR1 (wCR ₁ =7%)				CR2 (wCR ₂ =20%)				CR3 (wCR ₃ =3%)				CR4 (wCR ₄ =12%)				CR5 (wCR ₅ =27%)				CR6 (wCR ₆ =10%)				CR7 (wCR ₇ =14%)				CR8 (wCR ₈ =7%)				P _{jk}
	c _{1,jk} ^(>)	c _{1,jk} ^(~)	c _{1,jk} ^(<)	c _{1,jk}	c _{2,jk} ^(>)	c _{2,jk} ^(~)	c _{2,jk} ^(<)	c _{2,jk}	c _{3,jk} ^(>)	c _{3,jk} ^(~)	c _{3,jk} ^(<)	c _{3,jk}	c _{4,jk} ^(>)	c _{4,jk} ^(~)	c _{4,jk} ^(<)	c _{4,jk}	c _{5,jk} ^(>)	c _{5,jk} ^(~)	c _{5,jk} ^(<)	c _{5,jk}	c _{6,jk} ^(>)	c _{6,jk} ^(~)	c _{6,jk} ^(<)	c _{6,jk}	c _{7,jk} ^(>)	c _{7,jk} ^(~)	c _{7,jk} ^(<)	c _{7,jk}	c _{8,jk} ^(>)	c _{8,jk} ^(~)	c _{8,jk} ^(<)	c _{8,jk}	
EC ₁ , EC ₂	0	1	0	0.5	0	1	0	0.5	0	1	0	0.5	0	1	0	0.5	0	0	1	0.0	0	0	1	0.0	1	0	0	1.0	0	0	1	0.0	0.350
EC ₁ , EC ₃	0	1	0	0.5	0	1	0	0.5	0	1	0	0.5	0	1	0	0.5	0	1	0	0.5	0	0	1	0.0	0	1	0	0.5	0	1	0	0.5	0.450
EC ₁ , EC ₄	0	1	0	0.5	0	1	0	0.5	0	1	0	0.5	0	1	0	0.5	0	1	0	0.5	0	1	0	0.0	1	0	0	1.0	0	0	1	0.0	0.535
EC ₁ , EC ₅	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	1	0	0.5	0	0	1	0.0	0	1	0	0.5	1	0	0	1.0	0	1	0	0.5	0.285
EC ₁ , EC ₆	0	1	0	0.5	0	0	1	0.0	0	0	1	0.0	0	1	0	0.5	0	0	1	0.0	0	0	1	0.0	1	0	0	1.0	0	0	1	0.0	0.235
EC ₁ , EC ₇	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	1	0	0.5	0	0	1	0.0	0	0	1	0.0	1	0	0	1.0	0	0	1	0.0	0.200
EC ₁ , EC ₈	0	1	0	0.5	0	1	0	0.5	0	1	0	0.5	0	0	1	0.0	0	1	0	0.5	0	0	1	0.0	1	0	0	1.0	0	1	0	0.5	0.460
EC ₁ , EC _∅	0	1	0	0.5	0	1	0	0.5	0	1	0	0.5	0	1	0	0.5	0	1	0	0.5	0	1	0	0.5	1	0	0	1.0	0	1	0	0.5	0.570
EC ₁ , EC _∆	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	1	0	0	1.0	0	0	1	0.0	0.140
EC ₁ , EC _∅	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	1	0	0	1.0	0	0	1	0.0	0.140
EC ₁ , EC _•	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	1	0	0.5	0	0	1	0.0	0.070
EC ₂ , EC ₃	0	1	0	0.5	0	1	0	0.5	0	1	0	0.5	0	1	0	0.5	1	0	0	1.0	1	0	0	1.0	0	0	1	0.0	1	0	0	1.0	0.650
EC ₂ , EC ₄	0	1	0	0.5	0	1	0	0.5	0	1	0	0.5	0	1	0	0.5	1	0	0	1.0	1	0	0	1.0	1	0	0	1.0	0	0	1	0.0	0.720
EC ₂ , EC ₅	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	1	0	0.5	0	1	0	0.5	1	0	0	1.0	1	0	0	1.0	1	0	0	1.0	0.505
EC ₂ , EC ₆	0	1	0	0.5	0	0	1	0.0	0	0	1	0.0	0	1	0	0.5	0	0	1	0.0	1	0	0	1.0	1	0	0	1.0	0	1	0	0.5	0.370
EC ₂ , EC ₇	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	1	0	0.5	0	1	0	0.5	1	0	0	1.0	1	0	0	1.0	0	1	0	0.5	0.470
EC ₂ , EC ₈	0	1	0	0.5	0	1	0	0.5	0	1	0	0.5	0	0	1	0.0	1	0	0	1.0	1	0	0	1.0	1	0	0	1.0	1	0	0	1.0	0.730
EC ₂ , EC _∅	0	1	0	0.5	0	1	0	0.5	0	1	0	0.5	0	1	0	0.5	1	0	0	1.0	1	0	0	1.0	1	0	0	1.0	1	0	0	1.0	0.790
EC ₂ , EC _∆	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	1	0	0	1.0	1	0	0	1.0	1	0	0	1.0	0	1	0	0.5	0.545
EC ₂ , EC _∅	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	1	0	0.5	1	0	0	1.0	0	1	0	0.5	0	0	1	0.0	0.305
EC ₂ , EC _•	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0.050
EC ₃ , EC ₄	0	1	0	0.5	0	1	0	0.5	0	1	0	0.5	0	1	0	0.5	0	1	0	0.5	1	0	0	1.0	1	0	0	1.0	0	0	1	0.0	0.585
EC ₃ , EC ₅	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	1	0	0.5	0	0	1	0.0	1	0	0	1.0	1	0	0	1.0	0	1	0	0.5	0.335
EC ₃ , EC ₆	0	1	0	0.5	0	0	1	0.0	0	0	1	0.0	0	1	0	0.5	0	0	1	0.0	0	1	0	0.5	1	0	0	1.0	0	0	1	0.0	0.285
EC ₃ , EC ₇	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	1	0	0.5	0	0	1	0.0	1	0	0	1.0	1	0	0	1.0	0	0	1	0.0	0.300
EC ₃ , EC ₈	0	1	0	0.5	0	1	0	0.5	0	1	0	0.5	0	0	1	0.0	0	1	0	0.5	1	0	0	1.0	1	0	0	1.0	0	1	0	0.5	0.560
EC ₃ , EC _∅	0	1	0	0.5	0	1	0	0.5	0	1	0	0.5	0	1	0	0.5	0	1	0	0.5	1	0	0	1.0	1	0	0	1.0	0	1	0	0.5	0.620
EC ₃ , EC _∆	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	1	0	0	1.0	1	0	0	1.0	0	0	1	0.0	0.240
EC ₃ , EC _∅	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	1	0	0.5	1	0	0	1.0	0	0	1	0.0	0.190
EC ₃ , EC _•	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	1	0	0.5	0	0	1	0.0	0.070
EC ₄ , EC ₅	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	1	0	0.5	0	0	1	0.0	0	1	0	0.5	0	1	0	0.5	1	0	0	1.0	0.250
EC ₄ , EC ₆	0	1	0	0.5	0	0	1	0.0	0	0	1	0.0	0	1	0	0.5	0	0	1	0.0	0	0	1	0.0	0	1	0	0.5	1	0	0	1.0	0.235
EC ₄ , EC ₇	0	0	1	0.0	0	0	1	0.0	0	0	1	0.0	0	1	0	0.5	0	0	1	0.0	0	0	1	0.0	0	1	0	0.5	1	0	0	1.0	0.200

Fig. 6 Calculation of the c_{ijk} values (bolded) by combining the binary coefficients $c_{ijk}^{(>)}, c_{ijk}^{(~)}, c_{ijk}^{(<)}$ obtained from the paired comparison relationships in Fig. 4, through Eq. 9. The relevant p_{jk} values (in the last column) are calculated by applying Eq. 10

calculation of the binary coefficients and the corresponding c_{ijk} values, for the paired comparison relationships in Fig. 5c.

Finally, the coefficients c_{ijk} are aggregated in the following weighted sum, which determines the p_{jk} value:

$$p_{jk} = \sum_{\forall i} (c_{i,jk} \cdot w_{CR_i}), \in [0, 1] \tag{10}$$

This value expresses the weighted fraction of CR for which the j -th EC has a greater influence than the k -th one. For example, the paired comparison “ EC_1, EC_2 ” (at the top of Fig. 6) would result in:

$$p_{12} = 0.5 \cdot 7\% + 0.5 \cdot 20\% + 0.5 \cdot 3\% + 0.5 \cdot 12\% + 0 \cdot 27\% + 0 \cdot 10\% + 1 \cdot 14\% + 0 \cdot 7\% = 0.350 \tag{11}$$

The p_{jk} values can be aggregated into a \mathbf{P} matrix of proportions, which—consistently with the LCJ—is made up of elements that are symmetrical with respect to the main diagonal and complementary to each other with respect to the unit⁴. Figure 7a contains the \mathbf{P} matrix that results from the paired comparison relationships in Fig. 5c. Henceforth, the traditional

LCJ (cf. Section 4) is applied, determining the \mathbf{Z} matrix (see Fig. 7b) and, subsequently, the *interval* scaling (x) of the ECs (see Fig. 7c).

(d) *Scale anchoring.* Taking inspiration from the methodology developed in (Franceschini and Maisano 2019), the interval scaling resulting from the LCJ can be “anchored” with respect to the (unknown) *psychological continuum*, using two of the previously introduced dummy objects: EC_{\emptyset} , i.e., an anchor object corresponding to the absence of relationship, and EC_{\bullet} , i.e., an anchor object corresponding to the maximum possible degree of intensity of the relationship with the CR of interest. Therefore, the (interval) scale (x) is transformed into a new one (y), defined in the conventional range $[0, 10]$, through the following linear transformation:

$$\frac{y_k - 0}{10 - 0} = \frac{x_k - x_{\emptyset}}{x_{\bullet} - x_{\emptyset}} \rightarrow y_k = 10 \cdot \frac{x_k - x_{\emptyset}}{x_{\bullet} - x_{\emptyset}}, \tag{12}$$

where.

⁴ By combining the relationships $c_{ijk}^{(>)} + c_{ijk}^{(~)} + c_{ijk}^{(<)} = 1$, $c_{i,jk} = 1 - c_{i,kj}$, and $\sum_{\forall i} w_{CR_i} = 1$, it can be easily deduced that $p_{jk} = 1 - p_{kj}$.

(a) Matrix P

	EC_1	EC_2	EC_3	EC_4	EC_5	EC_6	EC_7	EC_8	EC_\emptyset	EC_Δ	EC_\circ	EC_\bullet
EC_1	0.500	0.650	0.550	0.465	0.715	0.765	0.800	0.540	0.430	0.860	0.860	0.930
EC_2	0.350	0.500	0.350	0.280	0.495	0.630	0.530	0.270	0.210	0.455	0.695	0.950
EC_3	0.450	0.650	0.500	0.415	0.665	0.715	0.700	0.440	0.380	0.760	0.810	0.930
EC_4	0.535	0.720	0.585	0.500	0.750	0.765	0.800	0.575	0.465	0.930	0.930	0.965
EC_5	0.285	0.505	0.335	0.250	0.500	0.785	0.635	0.325	0.215	0.430	0.715	1.000*
EC_6	0.235	0.370	0.285	0.235	0.215	0.500	0.265	0.225	0.165	0.365	0.465	0.765
EC_7	0.200	0.470	0.300	0.200	0.365	0.735	0.500	0.240	0.130	0.345	0.665	0.950
EC_8	0.460	0.730	0.560	0.425	0.675	0.775	0.760	0.500	0.390	0.830	0.880	0.940
EC_\emptyset	0.570	0.790	0.620	0.535	0.785	0.835	0.870	0.610	0.500	1.000*	1.000*	1.000*
EC_Δ	0.140	0.545	0.240	0.070	0.570	0.635	0.655	0.170	0.000*	0.500	1.000*	1.000*
EC_\circ	0.140	0.305	0.190	0.070	0.285	0.535	0.335	0.120	0.000*	0.000*	0.500	1.000*
EC_\bullet	0.070	0.050	0.070	0.035	0.000*	0.235	0.050	0.060	0.000*	0.000*	0.000*	0.500

(b) Matrix Z

	EC_1	EC_2	EC_3	EC_4	EC_5	EC_6	EC_7	EC_8	EC_\emptyset	EC_Δ	EC_\circ	EC_\bullet
EC_1	0.000	0.385	0.126	-0.088	0.568	0.722	0.842	0.100	-0.176	1.080	1.080	1.476
EC_2	-0.385	0.000	-0.385	-0.583	-0.013	0.332	0.075	-0.613	-0.806	-0.113	0.510	1.645
EC_3	-0.126	0.385	0.000	-0.215	0.426	0.568	0.524	-0.151	-0.305	0.706	0.878	1.476
EC_4	0.088	0.583	0.215	0.000	0.674	0.722	0.842	0.189	-0.088	1.476	1.476	1.812
EC_5	-0.568	0.013	-0.426	-0.674	0.000	0.789	0.345	-0.454	-0.789	-0.176	0.568	3.000
EC_6	-0.722	-0.332	-0.568	-0.722	-0.789	0.000	-0.628	-0.755	-0.974	-0.345	-0.088	0.722
EC_7	-0.842	-0.075	-0.524	-0.842	-0.345	0.628	0.000	-0.706	-1.126	-0.399	0.426	1.645
EC_8	-0.100	0.613	0.151	-0.189	0.454	0.755	0.706	0.000	-0.279	0.954	1.175	1.555
EC_\emptyset	0.176	0.806	0.305	0.088	0.789	0.974	1.126	0.279	0.000	3.000	3.000	3.000
EC_Δ	-1.080	0.113	-0.706	-1.476	0.176	0.345	0.399	-0.954	-3.000	0.000	3.000	3.000
EC_\circ	-1.080	-0.510	-0.878	-1.476	-0.568	0.088	-0.426	-1.175	-3.000	-3.000	0.000	3.000
EC_\bullet	-1.476	-1.645	-1.476	-1.812	-3.000	-0.722	-1.645	-1.555	-3.000	-3.000	-3.000	0.000

(c) Scaling

	EC_1	EC_2	EC_3	EC_4	EC_5	EC_6	EC_7	EC_8	EC_\emptyset	EC_Δ	EC_\circ	EC_\bullet
Σ_k	-6.116	0.336	-4.167	-7.989	-1.627	5.202	2.161	-5.794	-13.545	0.183	9.025	22.330
x_k	-0.510	0.028	-0.347	-0.666	-0.136	0.434	0.180	-0.483	-1.129	0.015	0.752	1.861
y_k	2.1	3.9	2.6	1.5	3.3	5.2	4.4	2.2	0.0	3.8	6.3	10.0

Fig. 7 a P matrix, b Z matrix, and c scaling resulting from the application of the proposed procedure to the test case. Items marked with “*” in the matrix P are associated with values of ± 3 in the matrix Z (cf. Section 4). Σ_k is the summation of the values reported

in the k -th column of the matrix Z ; the x_k values concern the interval scaling resulting from the LCJ; the y_k values concern the ratio scaling downstream of the anchoring in Eq. 12




Fig. 8 Comparison between the results of the new prioritization procedure (y-scaling) and the traditional one (ISM), with reference to the test case in Sect. 3. Note that the ISM results have been previously calculated in Fig. 2, while the y-scaling has been previously determined in Fig. 7c

x_{\emptyset} and x_{\bullet} are the scale values of EC_{\emptyset} and EC_{\bullet} respectively, resulting from the LCJ;

x_k is the scale value of a generic k -th EC, resulting from the LCJ;

y_k is the relevant transformed scale value in the conventional range $[0, 10]$.

The new scale (y) has a conventional unit and a zero point (which corresponds to the absence of the attribute); it can therefore be considered as a *ratio* scale.

Although the mathematical operations required to implement the LCJ might seem complex, they are in fact computationally straightforward and can be entirely automated using a basic spreadsheets such as MS Excel (Brown and Peterson 2009; Franceschini and Maisano 2019).

6 Concluding remarks

This paper introduced a new procedure for prioritizing the HoQ's ECs, based on Thurstone's LCJ. Besides being conceptually more rigorous and overcoming some shortcomings of the traditional ISM, the new procedure offers several other advantages. First, it can be integrated into the traditional procedure for constructing the HoQ without requiring additional or extra works from participants. This means that interviewees providing the VoC and the QFD team members responsible for creating the relationship matrix do not have to change their traditional work, as the mathematical implementation of the LCJ is fully automatable with a simple spreadsheet. Second, it prioritizes the ECs in the form of a *ratio* scaling, anchoring the LCJ solution through several dummy ECs. In addition, it

is easy to implement, flexible, and adaptable to other response modes than the traditional one (e.g., those in which judgments are inherently uncertain or incomplete). The effectiveness and the robustness of the new EC prioritization are ensured by the LCJ itself, which is a well-established technique that has been used and tested in multiple contexts (Franceschini et al. 2022).

With reference to the exemplified test case, it is interesting to note that the proposed procedure produces results quite in line with those of the ISM, as shown in Fig. 8. This diagram denotes a relatively high correlation (i.e., Pearson correlation coefficient $R^2 \approx 0.8602$) between the two approaches, which produce two very similar final rankings of ECs (only a *rank reversal* between EC_1 and EC_8 is observed):

ISM : $EC_6 > EC_7 > EC_2 > EC_5 > EC_3 > EC_1 > EC_8 > EC_4$

New procedure : $EC_6 > EC_7 > EC_2 > EC_5 > EC_3 > EC_8 > EC_1 > EC_4$.

(13)

Other tests have confirmed some agreement between the proposed procedure and the ISM although there are specific situations in which the two approaches may produce different results. Precisely for these situations, the authors claim the superiority of the new procedure, as it avoids questionable and potentially distorting operations. This aspect will be further investigated in future studies.

In conclusion, the new procedure represents an additional tool in the QFD-team's toolbox, to expand the perspective of analysis. The limitations of the proposed procedure are those inherent in the LCJ, namely some postulates concerning the (normal) distribution of judgments in the psychological continuum (cf. Section 4).

Regarding the future, possible adaptations of the proposed procedure to problems characterized by uncertain and/or incomplete formulation of relationships between ECs and CRs will be investigated. Additionally, a structured comparison will be made between the results of the new procedure and those of other alternative procedures to ISM, which are already present in the scientific literature.

Acknowledgements This research was carried out under the MICS (Made in Italy, Circular and Sustainable) Extended Partnership and partially funded by the European Union Next-GenerationEU (Piano Nazionale di Ripresa e Resilienza—Missione 4, Componente 2, Investimento 1.3, D.D. 1551.11-10-2022, PE00000004).

Authors contributions The authors have provided an equal contribution to the drafting of the paper.

Funding Open access funding provided by Politecnico di Torino within the CRUI-CARE Agreement.

Data availability Not applicable.

Declarations

Conflict of interests The authors do not have conflict of interest.

Ethical approval The authors respect the Ethical Guidelines of the Journal.

Consent to participate Not applicable.

Consent to publish Not applicable.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Akao Y (1994) Development history of quality function deployment. *Cust Driven Approach Qual Plan Deploy* 339:90
- Brown TC, Peterson GL (2009) An enquiry into the method of paired comparison: reliability, scaling, and Thurstone's Law of Comparative Judgment. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. General Technical Report RMRS-GTR-216WWW, U.S. Forest Service Fort Collins, Colorado
- Dym CL, Wood WH, Scott MJ (2002) Rank ordering engineering designs: pairwise comparison charts and Borda counts. *Res Eng Design* 13:236–242
- Franceschini F (2001) *Advanced quality function deployment*. CRC Press
- Franceschini F, Maisano D (2018) A new proposal to improve the customer competitive benchmarking in QFD. *Qual Eng* 30(4):730–761
- Franceschini F, Maisano D (2019) Fusing incomplete preference rankings in design for manufacturing applications through the ZM_{II} -technique. *Int J Adv Manuf Technol* 103:3307–3322
- Franceschini F, Rossetto S (1995) QFD: the problem of comparing technical/engineering design requirements. *Res Eng Design* 7:270–278
- Franceschini F, Galetto M, Maisano D, Mastrogiacomo L (2015) Prioritisation of engineering characteristics in QFD in the case of customer requirements orderings. *Int J Prod Res* 53(13):3975–3988
- Franceschini F, Maisano DA, Mastrogiacomo L (2022) Rankings and decisions in engineering. Springer International Publishing, Cham, Switzerland
- Galetto M, Franceschini F, Maisano D, Mastrogiacomo L (2018) Engineering characteristics prioritisation in QFD using ordinal scales: a robustness analysis. *Eur J Ind Eng* 12(2):151–174
- Ghorabae MK, Zavadskas EK, Olfat L, Turskis Z (2015) Multi-criteria inventory classification using a new method of evaluation based on distance from average solution (EDAS). *Informatica* 26:435–451
- Huang J, Mao LX, Liu HC, Song MS (2022) Quality function deployment improvement: a bibliometric analysis and literature review. *Qual Quant* 56(3):1347–1366
- Kelly KT, Richardson M, Isaacs T (2022) Critiquing the rationales for using comparative judgement: a call for clarity. *Assess Educ Princ Polic Pract* 29(6):674–688. <https://doi.org/10.1080/0969594X.2022.2147901>
- Li S, Tang D, Wang Q (2019) Rating engineering characteristics in open design using a probabilistic language method based on fuzzy QFD. *Comput Ind Eng* 135:348–358
- Liu X, Ma Y (2021) A method to analyze the rank reversal problem in the ELECTRE II method. *Omega* 102:102317
- Lyman D (1990) Deployment normalization. In: *Transactions from the Second Symposium on Quality Function Deployment*. Automotive Division of the American Society for Quality Control, the American Supplier Institute, Inc., Dearborn, MI, and GOAL/QPC, Methuen, MA, pp. 307–315
- Mao LX, Liu R, Mou X, Liu HC (2021) New approach for quality function deployment using linguistic Z-numbers and EDAS method. *Informatica* 32(3):565–582
- Maranell G (ed) (1974) *Scaling: a sourcebook for behavioral scientists*, 1st edn. Routledge, New York
- Ping YJ, Liu R, Lin W, Liu HC (2020) A new integrated approach for engineering characteristic prioritization in quality function deployment. *Adv Eng Inform* 45:101099
- Shi H, Mao LX, Li K, Wang XH, Liu HC (2022) Engineering characteristics prioritization in quality function deployment using an improved ORESTE method with double hierarchy hesitant linguistic information. *Sustainability* 14(15):9771
- Thurstone LL (1927) A law of comparative judgments. *Psychol Rev* 34(4):273
- Yager RR (2001) Fusion of multi-agent preference orderings. *Fuzzy Sets Syst* 117(1):1–12
- Zare Mehrjerdi Y (2010) Quality function deployment and its extensions. *Int J Qual Reliab Manag* 27(6):616–640