

Rhythmic response of cave animals to external cycles

Original

Rhythmic response of cave animals to external cycles / Lapadula, S.; Balestra, V.; Barzaghi, B.; Falaschi, M.; Ficaretola, G. F.; Manenti, R.. - ELETTRONICO. - (2024), pp. 1138-1138. (Intervento presentato al convegno SGI-SIMP Congress "Geology for a sustainable management of our Planet" tenutosi a Bari nel 2-5 September 2024).

Availability:

This version is available at: 11583/2992308 since: 2024-09-08T22:47:32Z

Publisher:

Società Geologica Italiana

Published

DOI:

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)



Bari, 2-5 September 2024

ABSTRACT BOOK

a cura della Società Geologica Italiana



Geology for a sustainable management of our Planet



Politecnico di Bari



Rhythmic response of cave animals to external cycles

Lapadula S.¹, Balestra V.*², Barzaghi B.¹, Falaschi M.¹, Ficetola G.F.¹ & Manenti R.¹

¹ Dipartimento di Scienze e Politiche Ambientali, Università di Milano

² Dipartimento di Ingegneria dell'Ambiente, del Territorio e delle Infrastrutture, Politecnico di Torino.

Corresponding author email: stefano.lapadula@unimi.it

Keywords: biodiversity, cave biology, springs.

Organisms inhabiting caves are usually described as lacking biological cycles, since they exploit environments with largely constant conditions. However, recent evidence is emerging on the rhythmicity of cave-dwelling animals, with experimental, observational and genetic analyses. There are still many uncertainties, as some organisms may have regular or irregular rhythms, and some others seem to lack them. In this study, we analysed the activity of two groundwater-dwelling species: the European cave salamander (*Proteus anguinus*) and the crustacean isopod *Monolistra pavani*. Both species are usually thought to be arrhythmic and exclusively found in caves; however, for the cave salamanders, recent studies have described the non-random activity in surface freshwater springs. We used GLMMs and N-mixture models to test the relationship between abundance and detection probability of cave species and both external cycles (circadian, lunar, and seasonal) and flooding events (upstream cumulated rain), analysing in two different models the surface and underground sites. We found that cave salamanders are generally more abundant during the night. Their detection probability in springs was higher during the night, with higher moon illumination (i.e., full moon) and after upstream rains; in caves, detection probability was higher during winter. For isopods, the detection probability was higher in early summer and in the evening. Our results suggest that depigmented cave-dwelling organisms are vulnerable to sunlight and tend to avoid it, exploiting the spring environment during the night, when ecological conditions are more similar to caves. Higher activity in springs may as well be related to food availability, as floods may foster it. Future studies to clarify these patterns are needed, and a molecular approach may be necessary to finally disentangle the role of adaptations and plasticity. These results are in accordance with previous studies on cave-dwelling invertebrates. In conclusion, cave animals may exhibit a rhythmic behaviour both in springs and in caves.