

Design of a Digital Twin of the Heart for the Management of Heart Failure Patients

Original

Design of a Digital Twin of the Heart for the Management of Heart Failure Patients / Scotto, A., Giordano, N., Rosati, S., Balestra, G.. - ELETTRONICO. - 316:(2024), pp. 875-876. (Medical Informatics Europe (MIE) 2024 Athens (Greece) 25-29 August 2024) [10.3233/shti240551].

Availability:

This version is available at: 11583/2992407 since: 2024-09-12T13:20:38Z

Publisher:

IOS Press

Published

DOI:10.3233/shti240551

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Design of a Digital Twin of the Heart for the Management of Heart Failure Patients

Andrea SCOTTO^{a,1}, Noemi GIORDANO^a, Samanta ROSATI^a and Gabriella BALESTRA^a

^aDepartment of Electronics and Telecommunication, Politecnico di Torino, Italy

ORCID ID: Andrea Scotto <https://orcid.org/0000-0002-3317-3843>

Noemi Giordano <https://orcid.org/0000-0002-9265-6538>

Samanta Rosati <https://orcid.org/0000-0003-0620-594X>

Gabriella Balestra <https://orcid.org/0000-0003-2717-648X>

Abstract. Heart failure poses a significant global health burden with high prevalence and mortality rates. A promising possibility in this context is the constant monitoring of the patients through telemedicine. The aim of this work is to present a digital twin of a patient at risk of heart failure. Applying machine learning to the recorded data of the patient, the system is able to early detect potential issues and improve the outcome.

Keywords. Digital Twin, personalized medicine, heart failure, telemedicine

1. Introduction

Heart failure (HF) is a complex clinical syndrome that results from any structural or functional impairment of ventricular filling or ejection of blood [1]. HF is associated with high morbidity, mortality [2] and hospital readmission rate [3]. In 2017 the estimated prevalence was around 64 million people worldwide [4], expected to grow even more due to overall aging of the population, thus, leading to more hospitalizations.

The Covid-19 pandemic resulted in a rapid-shift to “remote-by-default” healthcare, increasing interest in telemedicine. Its application in the HF context facilitates patients to rapidly access quality care when needed and minimizes the frequency of clinical visits [5], reducing the hospitalization rates.

Another emerging tool are digital twins, virtual replicas of a physical system that, constantly updates with real-time data, mimics its real-world counterpart.

The aim of this work is to present the main features of a Digital Twin, tailored to the characteristics of HF patients to remotely monitor them and improve their outcome.

2. The Digital Twin

In figure 1 the block diagram of the proposed Digital Twin (DT) is presented. The heart of the DT is a cloud, which stores the patient data captured either daily through wearable

¹ Corresponding Author: Andrea Scotto, Department of Electronics and Telecommunications, Politecnico di Torino, Corso Duca degli Abruzzi 24, 10129 Torino; E-mail: andrea.scotto@polito.it.

devices in a home environment, from clinical reports acquired during periodical visits or produced by the DT itself. On the top-left there is a coupled electro-mechanical (EM) model of the human heart, able to generate synchronous electrocardiograms (ECG) and phonocardiograms (PCG) based on the characteristics of the patients. Its role is to generate a theoretical ECG and each component of the associated acoustic signal to train a machine learning (ML) tool to automatically separate the two elements of each heart tone (i.e. M1 and T1 for the first tone and P2 and A2 for the second tone) from real PCGs. Eventually, the aim is to improve the accuracy in the detection of cardiac time intervals [6], proved to be associated with a worsening in the status of HF patients [7]. On the top right there is a second tool, which leverages ML algorithms to analyze the data collected in the cloud. Its aim is to forecast potential health issues, enabling preemptive measures to prevent hospital readmissions and adverse outcomes or to suggest if patients can be discharged successfully or not if they are already hospitalized. Another feature is the ability of the DT to send an alert whenever it detects a worsening in the condition of the patient, allowing the clinician to quickly react and help the patient.




Figure 1. Block diagram of the Digital Twin.

3. Conclusions

In this work we introduce a Digital Twin. Its advantage is that it combines different tools to effectively monitor either at-risk patients or those who are already affected by the pathology to quickly detect a deterioration in their health and react promptly.

References

- [1] Heidenreich PA et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation* 2022;145(18), doi: 10.1161/CIR.0000000000001063
- [2] Emmons-Bell S, Johnson C, Roth G. Prevalence, incidence and survival of heart failure: a systematic review. *Heart* 2022;108(17):1351-1360, doi: 10.1136/heartjnl-2021-320131
- [3] Jencks SF, Williams MV, Coleman EA. Rehospitalizations among patients in the Medicare fee-for-service program. *N Engl J Med.* 2009;360:1418-1428, doi: 10.1056/NEJMs0803563
- [4] Savasere G et al. Global burden of heart failure: a comprehensive and updated review of epidemiology. *Cardiovascular Research* 2022;118(17):3272-3287, doi: 10.1093/cvr/cvac013
- [5] Singhal A, Riley JP, Cowie MR. Benefits and challenges of telemedicine for heart failure consultations: a qualitative study. *BMC Health Services Research* 2023;23(847), doi: 10.1186/s12913-023-09872-z
- [6] Giordano N, Rosati S, Balestra G, Knaflitz M. A Wearable Multi-Sensor Array Enables the Recording of Heart Sounds in Homecare. *Sensors* 2023;2(13), doi: 10.3390/s23136241
- [7] Shitara J et al. Temporal changes of cardiac acoustic biomarkers and cardiac function in acute decompensated heart failure. *ESC Hear. Fail.* 2021;8(5):4037-4047, doi: 10.1002/ehf2.13492