

Microplastic and microfibre pollution in Greenland: a preliminary study

*Original*

Microplastic and microfibre pollution in Greenland: a preliminary study / Balestra, Valentina; Akyildiz, SINEM HAZAL; Wadhams, Peter; Bellopede, Rossana. - ELETTRONICO. - (2024). ( MICRO2024 Plastic Pollution from Macro to Nano Lanzarote 23-27 September 2024).

*Availability:*

This version is available at: 11583/2993219 since: 2024-10-09T15:00:55Z

*Publisher:*

MICRO Community

*Published*

DOI:

*Terms of use:*

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

(Article begins on next page)

---

# Microplastic and microfibre pollution in Greenland: a preliminary study

Valentina Balestra\*<sup>†1</sup>, Sinem Hazal Akyildiz<sup>1</sup>, Peter Wadhams<sup>1,2</sup>, and Rossana Bellopede<sup>1</sup>

<sup>1</sup>Department of Environment, Land and Infrastructure Engineering - Politecnico di Torino (DIATI - PoliTO) – Politecnico di Torino - Corso Duca degli Abruzzi, 24 10129 Torino, Italy

<sup>2</sup>Department of Applied Mathematics and Theoretical Physics - University of Cambridge (DAMTP - University of Cambridge) – The Old Schools Trinity Lane Cambridge CB2 1TN United Kingdom, United Kingdom

## Abstract

Microplastics (MPs) and microfibrils (MFs) contaminate the environment worldwide. However, MP pollution is less studied in remote and extreme environments and MF monitoring is at early stage. Atmospheric currents are one of the most important factors that contribute to the transport of these micropollutants, especially in remote areas, therefore, monitoring these environments is useful to understand the seriousness of the problem from an environmental point of view. Within a project of the Raw Materials Engineering group of DIATI, Politecnico di Torino and Extreme E, preliminary investigations in Greenland glaciers were done to understand MP and MF pollution. Samples were collected in different areas, along a track crossed by members of Extreme E during the electric off-road racing series' visit to the region. Collected samples were pre-treated with HO and filtered. Particles on filters were counted and characterized with microscopy techniques, with and without UV light. Preliminary results indicated that MPs and MFs were present in all examined samples. Micropollutant abundances increased considering that smaller size and fibre-shape were the most present factors. Spectroscopic analyses showed different kinds of plastics and a high amount of cellulosic fibres from anthropogenic origin. However, these data could be affected by several factors, especially linked to climatic factors, such as snow fusion, seasonal variations, and weather. Therefore, other detailed analyses will be necessary. Moreover, this preliminary monitoring in extreme cold environments has highlighted different problems, especially in sampling, useful to the improvement of the methods for future analyses.

**Keywords:** microplastics, microfibrils, greenland, pollution, ice

---

\*Speaker

<sup>†</sup>Corresponding author: [valentina.balestra@polito.it](mailto:valentina.balestra@polito.it)