

Correction to: Predictability of orthodontic tooth movement with aligners: effect of treatment design

Original

Correction to: Predictability of orthodontic tooth movement with aligners: effect of treatment design / Castroflorio, Tommaso; Sedran, Ambra; Parrini, Simone; Garino, Francesco; Reverdito, Matteo; Capuozzo, Riccardo; Mutinelli, Sabrina; Grybauskas, Simonas; Vaitieknas, Mantas; Deregibus, Andrea. - In: PROGRESS IN ORTHODONTICS. - ISSN 2196-1042. - 24:1(2023). [10.1186/s40510-023-00499-8]

Availability:

This version is available at: 11583/2993657 since: 2024-10-24T13:17:14Z

Publisher:

Springer

Published

DOI:10.1186/s40510-023-00499-8

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

CORRECTION

Open Access



Correction to: Predictability of orthodontic tooth movement with aligners: effect of treatment design

Tommaso Castroflorio¹, Ambra Sedran^{1*} , Simone Parrini², Francesco Garino³, Matteo Reverdito⁴, Riccardo Capuozzo⁵, Sabrina Mutinelli⁶, Simonas Grybauskas⁷, Mantas Vaitiekūnas⁸ and Andrea Deregibus¹

Correction to: *Progress in Orthodontics* (2023) 24:2
<https://doi.org/10.1186/s40510-022-00453-0>

Following publication of the original article [1], the authors identified an error in two sentences in the Discussion section.

The sentences currently read:

Similar results were obtained by Goh et al. [33] for the lower first molar only. Differences among the studies could be related to the fact that patients considered by Goh et al. were treated with a maximum number of 14 aligners, therefore with simplest malocclusions with respect to the ones analyzed by our team, and to different applied methodologies for data collection.

The sentences should read:

Similar results were obtained by Goh et al. [33] for the lower first molar only. Differences among the studies could be related to different applied methodologies for data collection, however, the number of aligners and treatment time of the study groups considered is comparable so careful monitoring of the vestibular torque prescription for these dental elements is strongly recommended.

The indicated sentences in the Discussion section have been updated above and the original article [1] has been corrected.

Published online: 24 October 2023

The original article can be found online at <https://doi.org/10.1186/s40510-022-00453-0>.

*Correspondence:

Ambra Sedran
ambra.sedran@gmail.com

¹ Department of Surgical Sciences, Dental School of the University of Torino, Via Nizza 230, 10126 Turin, Italy

² Poggibonsi, Italy

³ Turin, Italy

⁴ Cuneo, Italy

⁵ Caserta, Italy

⁶ Department of Neuroscience, School of Dentistry, Section of Pedodontics, University of Padova, Via VII Febbraio 2, 35122 Padua, Italy

⁷ University of Ferrara, Vilnius, Lithuania

⁸ Kaunas University of Technology, Kaunas, Lithuania

References

1. Castroflorio T, Sedran A, Parrini S, et al. Predictability of orthodontic tooth movement with aligners: effect of treatment design. *Prog Orthod*. 2023;24:2. <https://doi.org/10.1186/s40510-022-00453-0>.
33. Goh S, Dreyer C, Weir T. The predictability of the mandibular curve of Wilson, buccolingual crown inclination, and transverse expansion expression with Invisalign treatment. *Am J Orthod Dentofacial Orthop*. 2022. <https://doi.org/10.1016/j.ajodo.2021.09.020>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.