POLITECNICO DI TORINO Repository ISTITUZIONALE

Correction to: Predictability of orthodontic tooth movement with aligners: effect of treatment design

Original

Correction to: Predictability of orthodontic tooth movement with aligners: effect of treatment design / Castroflorio, Tommaso; Sedran, Ambra; Parrini, Simone; Garino, Francesco; Reverdito, Matteo; Capuozzo, Riccardo; Mutinelli, Sabrina; Grybauskas, Simonas; Vaitieknas, Mantas; Deregibus, Andrea. - In: PROGRESS IN ORTHODONTICS. - ISSN 2196-1042. - 24:1(2023). [10.1186/s40510-023-00499-8]

Availability: This version is available at: 11583/2993657 since: 2024-10-24T13:17:14Z

Publisher: Springer

Published DOI:10.1186/s40510-023-00499-8

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Castroflorio et al. Progress in Orthodontics

https://doi.org/10.1186/s40510-023-00499-8



Correction to: Predictability of orthodontic tooth movement with aligners: effect of treatment design

(2023) 24:47

Tommaso Castroflorio¹, Ambra Sedran^{1*}, Simone Parrini², Francesco Garino³, Matteo Reverdito⁴, Riccardo Capuozzo⁵, Sabrina Mutinelli⁶, Simonas Grybauskas⁷, Mantas Vaitiekūnas⁸ and Andrea Deregibus¹

Correction to: Progress in Orthodontics (2023) 24:2 https://doi.org/10.1186/s40510-022-00453-0

Following publication of the original article [1], the authors identified an error in two sentences in the Discussion section.

The sentences currently read:

Similar results were obtained by Goh et al. [33] for the lower first molar only. Differences among the studies could be related to the fact that patients considered by Goh et al. were treated with a maximum number of 14 aligners, therefore with simplest malocclusions with respect to the ones analyzed by our team, and to different applied methodologies for data collection.

The original article can be found online at https://doi.org/10.1186/s40510-022-00453-0.

*Correspondence:

- Ambra Sedran
- ambra.sedran@gmail.com
- ¹ Department of Surgical Sciences, Dental School of the University of Torino, Via Nizza 230, 10126 Turin, Italy
- ² Poggibonsi, Italy
- ³ Turin, Italy
- [†] Cuneo, Italy
- ⁵ Caserta, Italy
- ⁶ Department of Neuroscience, School of Dentistry, Section
- of Pedodontics, University of Padova, Via VII Febbraio 2, 35122 Padua, Italy ⁷ University of Ferrara, Vilnius, Lithuania
- ⁸ Kaunas University of Technology, Kaunas, Lithuania



© The Author(s) 2023. Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

The sentences should read:

Similar results were obtained by Goh et al. [33] for the lower first molar only. Differences among the studies could be related to different applied methodologies for data collection, however, the number of aligners and treatment time of the study groups considered is comparable so careful monitoring of the vestibular torque prescription for these dental elements is strongly recommended.

The indicated sentences in the Discussion section have been updated above and the original article [1] has been corrected.

Published online: 24 October 2023

References

- Castroflorio T, Sedran A, Parrini S, et al. Predictability of orthodontic 1. tooth movement with aligners: effect of treatment design. Prog Orthod. 2023;24:2. https://doi.org/10.1186/s40510-022-00453-0.
- 33. Goh S, Dreyer C, Weir T. The predictability of the mandibular curve of Wilson, buccolingual crown inclination, and transverse expansion expression with Invisalign treatment. Am J Orthod Dentofacial Orthop. 2022. https:// doi.org/10.1016/j.ajodo.2021.09.020.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.