

Castelnuovo–Mumford regularity and splitting criteria for logarithmic bundles over rational normal scroll surfaces

Original

Castelnuovo–Mumford regularity and splitting criteria for logarithmic bundles over rational normal scroll surfaces / Di Gennaro, R.; Malaspina, F.. - In: INDAGATIONES MATHEMATICAE. - ISSN 0019-3577. - (2024), pp. 1-17. [10.1016/j.indag.2024.10.002]

Availability:

This version is available at: 11583/2995430 since: 2024-12-16T11:17:03Z

Publisher:

Elsevier

Published

DOI:10.1016/j.indag.2024.10.002

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)



Castelnuovo–Mumford regularity and splitting criteria for logarithmic bundles over rational normal scroll surfaces

R. Di Gennaro^a, F. Malaspina^{b,*}

^a *Università degli Studi di Napoli Federico II, Complesso Universitario Monte Sant'Angelo, Via Cinthia, 80126 Napoli, Italy*

^b *Politecnico di Torino, Corso Duca degli Abruzzi 24, 10129 Torino, Italy*

Received 6 May 2024; received in revised form 12 September 2024; accepted 5 October 2024

Communicated by Cecilia Salgado

Abstract

We introduce and study a notion of Castelnuovo–Mumford regularity suitable for rational normal scroll surfaces. In this setting we prove analogs of some classical properties. We prove splitting criteria for coherent sheaves and a characterization of Ulrich bundles. Finally we study logarithmic bundles associated to arrangements of lines and rational curves.

© 2024 The Author(s). Published by Elsevier B.V. on behalf of Royal Dutch Mathematical Society (KWG). This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Keywords: Castelnuovo–Mumford regularity; Rational normal scroll; Splitting criteria; Logarithmic bundles

1. Introduction

In chapter 14 of [18] Mumford introduced the concept of regularity for a coherent sheaf on a projective space \mathbb{P}^n . It was soon clear that it was a key notion and a fundamental tool in many areas of algebraic geometry and commutative algebra.

From the perspective of algebraic geometry, regularity measures the complexity of a sheaf: the regularity of a coherent sheaf is an integer that estimates the smallest twist for which the sheaf is generated by its global sections. In Castelnuovo's much earlier version, if X is a closed subvariety of projective space and H is a general hyperplane, one uses linear systems (seen now as a precursor of sheaf cohomology) to get information about X from information about the intersection of X with H plus other geometrical or numerical assumptions on X .

* Corresponding author.

E-mail addresses: digennar@unina.it (R. Di Gennaro), francesco.malaspina@polito.it (F. Malaspina).

<https://doi.org/10.1016/j.indag.2024.10.002>

0019-3577/© 2024 The Author(s). Published by Elsevier B.V. on behalf of Royal Dutch Mathematical Society (KWG). This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

From the computational and commutative algebra point of view, the regularity is one of the most important invariants of a finitely generated graded module over a polynomial ring. Roughly, it measures the amount of computational resources that working with that module requires. More precisely the regularity of a module bounds the largest degree of the minimal generators and the degree of syzygies.

Over the years, extensions of this notion have been proposed to handle other ambient varieties beyond projective space, including Grassmannians [1], quadrics [3], multiprojective spaces [4,9,15], n -dimensional smooth projective varieties with an n -block collection [9], and abelian varieties [19]. For a different approach to multigraded regularity from a commutative algebra point of view, see [5,6].

The aim of this paper is to introduce a very simple and natural concept of regularity on a rational normal scroll surface.

The interesting fact is that on $\mathcal{Q}_2 \cong \mathbb{P}^1 \times \mathbb{P}^1$ our definition of regularity coincides with this definition of regularity on $\mathbb{P}^1 \times \mathbb{P}^1$ given in [3,4,9,15] and we are able to prove that every regular coherent sheaf is globally generated, as done by Mumford in the classical case \mathbb{P}^n . Maclagan and Smith [17] gave a variant of multigraded Castelnuovo–Mumford regularity, motivated by toric geometry. Our notion requires only three cohomological vanishing conditions and is strongly linked to the canonical sheaf.

The second aim of this paper is to apply our notion of regularity in order to investigate under what circumstances a vector bundle can be decomposed into a direct sum of line bundles. In particular, in the second section splitting criteria of vector bundle on a rational normal scroll surface are given, generalizing some analogous result already known for $\mathbb{P}^1 \times \mathbb{P}^1$ [3,4]. In [13] the authors give some splitting criteria for vector bundles of rank 2 in terms of Chern classes and vanishing of certain cohomology groups using Beilinson type spectral sequence. They also remark that their results are the best possible without analysing the differentials in the spectral sequence. Our splitting criteria work for vector bundles of arbitrary rank thanks to the use of regularity and without the use of spectral sequences and with only a small number of cohomological vanishing conditions. In [7,17] are given splitting criteria in a far more general context but with an infinite number of cohomological vanishing conditions (in the first case it is required a table of cohomology and in the second are considered the twists with all the ACM bundles). Furthermore in [12] Theorem B is given a complete classification of Ulrich bundles on rational normal scroll surfaces. Here we give an alternative and simpler proof without relying on derived category techniques.

Finally, the last section focuses on the logarithmic bundle of divisors on a rational normal scroll. It fits in the classical topic of the study of normal crossing divisors on a smooth complex variety X . When D is a normal crossing divisor, Deligne [20] constructed a mixed Hodge structure on $U = X \setminus D$ using the logarithmic de Rham complex $\Omega_X^\bullet(-\log D)$. Following this idea, in [21] Saito defined the sheaf $T_X(D)$ of derivations tangent to D and (dually) the sheaf of logarithmic one-forms with pole along D , the logarithmic bundle $\Omega^1_X(\log D)$.

The module of tangent derivations is a sheaf of \mathcal{O}_X -modules, such that if $f \in \mathcal{O}_{X,p}$ is a local defining equation for D at p , then

$$(T_X(-\log D))_p = \{\theta \in T_X \mid \theta(f) \in \langle f \rangle\}.$$

When D is a normal-crossing divisor, $T_X(-\log D)$ is always locally free.

The module of derivations tangent to D is a reflexive sheaf. So, since a reflexive sheaf on a surface is always locally free, so it is interesting to understand when it (or its dual) splits as free module, in which case the divisor is said to be a *free*. In general, free divisors are difficult

to find. We find some classes of free divisors (precisely free arrangements of lines and rational curves) on a two dimensional rational normal scroll.

We thank M. Aprodu, A.P. Rao, G. Casnati and J. Pons-Llopis for helpful discussions and fundamental observations.

2. Regularity on $S(a_0, a_1)$

Throughout this article, our base field is algebraically closed with characteristic 0. Let $X = S(a_0, a_1)$ be a smooth rational normal scroll, the image of $\mathbb{P}(\mathcal{E})$ via the morphism defined by $\mathcal{O}_{\mathbb{P}(\mathcal{E})}(1)$, where $\mathcal{E} \cong \mathcal{O}_{\mathbb{P}^1}(a_0) \oplus \mathcal{O}_{\mathbb{P}^1}(a_1)$ is a vector bundle of rank 2 on \mathbb{P}^1 with $0 < a_0 \leq a_1$. Letting $\pi : \mathbb{P}(\mathcal{E}) \rightarrow \mathbb{P}^1$ be the projection, we may denote by H and f , the hyperplane section corresponding to $\mathcal{O}_{\mathbb{P}(\mathcal{E})}(1)$ and the fibre corresponding to $\pi^*\mathcal{O}_{\mathbb{P}^1}(1)$, respectively. Following [14, Notation 2.8.1] $S(a_0, a_1)$ is a Hirzebruch (ruled) surface with $C_0 = H - a_1 f$, $C_0^2 = a_0 - a_1$ and $C_0 \cdot f = 1$, so $H^2 = a_0 + a_1$. Then we have $Pic(X) \cong \mathbf{Z}\langle H, f \rangle \cong \mathbf{Z}\langle C_0, f \rangle$ and $\omega_X \cong \mathcal{O}_X(-2H + (c - 2)f)$, where $c := a_0 + a_1 > 1$ is the degree of X .

For the computational purpose, we use the following lemma.

Lemma 2.1 ([10]). *For any $i = 0, 1, 2$, we have*

- (i) $H^i(X, \mathcal{O}_X(aH + bf)) \cong H^i(\mathbb{P}^1, \text{Sym}^a \mathcal{E} \otimes \mathcal{O}_{\mathbb{P}^1}(b))$ if $a \geq 0$;
- (ii) $H^i(X, \mathcal{O}_X(-H + bf)) = 0$ for any b ;
- (iii) $H^i(X, \mathcal{O}_X(aH + bf)) \cong H^{2-i}(\mathbb{P}^1, \text{Sym}^{-a-2} \mathcal{E} \otimes \mathcal{O}_{\mathbb{P}^1}(c - b - 2))$ if $a \leq -2$.

Recall the dual of the relative Euler exact sequence of X :

$$0 \rightarrow \mathcal{O}_X(-H + cf) \rightarrow \mathcal{O}_X(a_0 f) \oplus \mathcal{O}_X(a_1 f) \rightarrow \mathcal{O}_X(H) \rightarrow 0. \tag{1}$$

The pullback of the Euler sequence in \mathbb{P}^1 is

$$0 \rightarrow \mathcal{O}_X(-f) \rightarrow \mathcal{O}_X^2 \rightarrow \mathcal{O}_X(f) \rightarrow 0, \tag{2}$$

and we obtain

$$0 \rightarrow \mathcal{O}_X(-H + (c-2)f) \rightarrow \mathcal{O}_X^2(-H + (c-1)f) \rightarrow \mathcal{O}_X(a_0 f) \oplus \mathcal{O}_X(a_1 f) \rightarrow \mathcal{O}_X(H) \rightarrow 0, \tag{3}$$

We give a definition of regularity on X :

Definition 2.2. A coherent sheaf F on X is said to be (p, p') -regular if, denoting $E = F(pH + p'f)$,

$$h^2(E(-H + (c - 2)f)) = h^1(E(-H + (c - 1)f)) = h^1(E(-f)) = 0.$$

We will say that F is *regular* if it is $(0, 0)$ -regular. We will say that F is p -regular if it is $(p, 0)$ -regular. We define the *regularity* of F , $Reg(F)$, as the least integer p such that F is p -regular. We set $Reg(F) = -\infty$ if there is no such integer.

Remark 2.3. When $a_0 = a_1 = 1$, we get $c = 2$ and C_0 is a line; X is the quadric $\mathbb{P}^1 \times \mathbb{P}^1$ and this notion of regularity coincides with the notions of Castelnuovo–Mumford regularity given in [3,4,15] since $h^2(E(-H + (c - 2)f)) = h^2(E(-1, -1))$, $h^1(E(-H + (c - 1)f)) = h^1(E(-1, 0))$, $h^1(E(-f)) = h^1(E(0, -1))$.

Lemma 2.4. *If F is a regular coherent sheaf on X , then $h^1(F|_f((a-1)H + bf)) = 0$ for any $a \geq 0$ and for any integer b .*

Proof. Let us consider this exact cohomology sequence:

$$\dots \rightarrow H^1(F(-H + (c-1)f)) \rightarrow H^1(F|_f(-H + (c-1)f)) \rightarrow H^2(F(-H + (c-2)f)) \rightarrow \dots$$

Since the first and the third groups vanish by hypothesis, then also the middle group vanishes. $H^1(\mathcal{O}|_f(-H + (c-1)f)) \cong H^1(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(-1))$ so $H^1(F|_f((a-1)H + bf)) = 0$ for any $a \geq 0$ and for any integer b . \square

Lemma 2.5. *If F is a regular coherent sheaf on X , then $h^2(F((a-1)H + (c-2+b)f)) = 0$ for any $a, b \geq 0$ and $h^1(F(tf)) = 0$ for any $t \geq -1$.*

Proof. From (2) we get $h^2(F(-H + (c-2+t)f)) = 0$ for any $t \geq 0$. From (1) tensored by $F((c-2)f)$ we get $h^2(F((c-2+t)f)) = 0$ and again by (2) we obtain $h^2(F((c-2+t)f)) = 0$ for $t \geq 0$. In the same way $h^2(F((a-1)H + (c-2+b)f)) = 0$ for any $a \geq 0$ and for any $b \geq 0$. From

$$0 \rightarrow F(-f) \rightarrow F \rightarrow F|_f \rightarrow 0,$$

we deduce that $h^1(F(tf)) = 0$ for any $t \geq -1$. \square

Lemma 2.6. *If F is a regular coherent sheaf on X ,*

- (i) $H^1(F|_H((c-1)+bf)) = 0$ for any $b \geq 0$.
- (ii) $H^1(F|_H((a+1)H + (b-1)f)) = 0$ for any $a, b \geq 0$.

Proof. Let us consider this exact cohomology sequence:

$$\dots \rightarrow H^1(F((c-1)f)) \rightarrow H^1(F|_H((c-1)f)) \rightarrow H^2(F(-H + (c-1)f)) \rightarrow \dots$$

Since $H^1(F((c-1)f)) = H^2(F(-H + (c-1)f)) = 0$ we get $h^1(F|_H((c-1)f)) = h^1(\mathbb{P}^1, F_H(c-1)) = 0$ and also $h^1(\mathbb{P}^1, F|_H(c-1+t)) = h^1(F|_H((c-1+t)f)) = 0$ for $t \geq 0$. So (i) is proved. $H^1(\mathcal{O}|_H((a+1)H + (b-1)f)) \cong H^1(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}((a+1)c + b - 1))$ and when $a, b \geq 0$ we get $(a+1)c + b - 1 \geq c - 1$ so $H^1(F|_H((a+1)H + (b-1)f)) = 0$ for any $a, b \geq 0$. So also (ii) is proved. \square

Proposition 2.7. *Let F be a regular coherent sheaf on X then*

1. $F(pH + p'f)$ is regular for $p, p' \geq 0$.
2. $H^0(F(f))$ is spanned by

$$H^0(F) \otimes H^0(\mathcal{O}(f));$$

and $H^0(F(H))$ it is spanned by

$$H^0(F(a_0f)) \oplus H^0(F(a_1f)).$$

Proof. (1) Let F be a regular coherent sheaf, we want to show that also $F(H)$ is regular. $h^2(F((c-2)f)) = 0$ by Lemma 2.5. In order to show that $h^1(F((c-1)f)) = 0$ let us consider the exact cohomology sequence:

$$\dots \rightarrow H^1(F(-H + (c-1)f)) \rightarrow H^1(F((c-1)f)) \rightarrow H^1(F|_H((c-1)f)) \rightarrow \dots$$

We notice that the first group vanishes by hypothesis and the third group vanishes by (i) of [Lemma 2.6](#). Then also the middle group vanishes. It remains to show that $h^1(F(H - f)) = 0$. By the exact cohomology sequence:

$$\dots \rightarrow H^1(F(-f)) \rightarrow H^1(F(H - f)) \rightarrow H^1(F|_H(H - f)) \rightarrow \dots$$

since the first group vanishes by hypothesis and the third group vanishes by (ii) of [Lemma 2.6](#) we obtain that also the middle group vanishes. Let F be a regular coherent sheaf, we want show that also $F(f)$ is regular. $h^2(F(-H + (c - 1)f)) = h^1(F) = 0$ by [Lemma 2.5](#). In order to show that $h^1(F(-H + cf)) = 0$ let us consider the exact cohomology sequence:

$$\dots \rightarrow H^1(F(-H + (c - 1)f)) \rightarrow H^1(F(-H + cf)) \rightarrow H^1(F|_f(-H + cf)) \rightarrow \dots$$

We notice that the first group vanishes by hypothesis and the third group vanishes by [Lemma 2.4](#). Then also the middle group vanishes. (2) Let us consider (2) tensored by F :

$$0 \rightarrow F(-f) \rightarrow F^2 \rightarrow F(f) \rightarrow 0.$$

Since $H^1(F(-f)) = 0$, we obtain

$$H^0(F) \otimes H^0(\mathcal{O}_X(f)) \rightarrow H^0(F(f)) \rightarrow 0.$$

Now let us consider (3) tensored by F :

$$0 \rightarrow F(-H + (c - 2)f) \rightarrow F^2(-H + (c - 1)f) \rightarrow F(a_0f) \oplus F(a_1f) \rightarrow F(H) \rightarrow 0.$$

Since $H^2(F(-H + (c - 2)f)) = H^1(F(-H + (c - 1)f)) = 0$, we obtain

$$H^0(F(a_0f)) \oplus H^0(F(a_1f)) \rightarrow H^0(F(H)) \rightarrow 0. \quad \square$$

Remark 2.8. If F is a regular coherent sheaf on X then it is globally generated.

In fact by the above proposition we have the following surjections:

$$H^0(F)^q \rightarrow H^0(F(a_0f)) \oplus H^0(F(a_1f)) \rightarrow H^0(F(H)),$$

for a suitable positive integer q . So also the map

$$H^0(F)^q \rightarrow H^0(F(H))$$

is a surjection.

Moreover we can consider a sufficiently large twist l such that $F(lH)$ is globally generated. For a suitable positive integer q' the commutativity of the diagram

$$\begin{array}{ccc} H^0(F)^{q'} \otimes \mathcal{O}_X & \rightarrow & H^0(F(lH)) \otimes \mathcal{O}_X \\ \downarrow & & \downarrow \\ H^0(F)^{q'} \otimes \mathcal{O}(lH) & \rightarrow & F(lH) \end{array}$$

and the surjectivity of the top horizontal map and the two vertical maps yield the surjectivity of $H^0(F) \otimes \mathcal{O}(lH) \rightarrow F(lH)$, which implies that F is generated by its sections.

Remark 2.9. $\mathcal{O}_X(aH - bf)$ is regular if and only if $a \geq 0$ and $b \leq aa_0$.

Remark 2.10. In particular $\mathcal{O}_X, \mathcal{O}_X(f), \mathcal{O}_X(H - f)$ are regular but not -1 -regular so $Reg(\mathcal{O}_X) = Reg(\mathcal{O}_X(f)) = Reg(\mathcal{O}_X(H - f)) = 0$.

3. Splitting criteria and ulrich bundles

It is possible to use this notion of regularity in order to prove splitting criteria for vector bundles:

Theorem 3.1. *Let E be a rank r vector bundle on X .*

Then following conditions are equivalent:

1. *for any integer t ,*

$$h^1(E(tH + (c - 1)f)) = h^1(E(tH - f)) = 0,$$

2. *There are r integer t_1, \dots, t_r such that $E \cong \bigoplus_{i=1}^r \mathcal{O}_X(t_i H)$.*

Proof. (1) \Rightarrow (2). Let assume that t is an integer such that $E(tH)$ is regular but $E((t - 1)H)$ not. By the definition of regularity and (1) we can say that $E((t - 1)H)$ is not regular if and only if $H^2(E((t - 2)H + (c - 2)f)) \neq 0$. By Serre duality we have that $H^0(E^\vee(-tH)) \neq 0$. Now since $E(tH)$ is globally generated by Remark 2.8 and $H^0(E^\vee(-tH)) \neq 0$ we can conclude that \mathcal{O}_X is a direct summand of $E(tH)$. By iterating these arguments we get (2). (2) \Rightarrow (1). $h^1(\mathcal{O}_X(tH + (c - 1)f)) = h^1(\mathcal{O}_X(tH - f)) = 0$, for any integer t , so if $E \cong \bigoplus_{i=1}^r \mathcal{O}(t_i H)$ then it satisfies all the conditions in (1). \square

Remark 3.2. If $c = 2$ the above theorem is the Horrocks criterion on $\mathbb{P}^1 \times \mathbb{P}^1$ (see [3,4]).

Corollary 3.3. *Let E be a vector bundle on X with $\text{Reg}(E) = 0$ and $H^2(E(-2H + (c - 2)f)) \neq 0$ or $H^1(E(-2H + (c - 1)f)) = H^1(E(-H - f)) = 0$, then \mathcal{O}_X is direct summand of E .*

Proof. Since $E(-H)$ is not regular, if $H^1(E(-2H + (c - 1)f)) = H^1(E(-H - f)) = 0$, we have $H^2(E(-2H + (c - 2)f)) \neq 0$ and \mathcal{O}_X is a direct summand of E by the proof of the above Theorem. \square

Theorem 3.4. *Let E be a vector bundle on X . Then following conditions are equivalent:*

1. *for any integer t ,*

$$\begin{aligned} H^1(E(tH)) &= H^1(E(tH + (c - 2)f)) = H^1(E(tH + (a_0 - 1)f)) \\ &= H^1(E(tH + (a_1 - 1)f)) = 0 \end{aligned}$$

2. *E is a direct sum of line bundles \mathcal{O}_X , $\mathcal{O}(f)$ and $\mathcal{O}(H - f)$ with a finite number of suitable twists $t_i H$.*

Proof. (1) \Rightarrow (2). Let assume that t is an integer such that $E(tH)$ is regular but $E((t - 1)H)$ not. Up to a twist we may assume $t = 0$. By the definition of regularity and (1) we can say that $E(-H)$ is not regular if and only if one of the following conditions is satisfied:

- (i) $h^2(E(-2H + (c - 2)f)) \neq 0$,
- (ii) $h^1(E(-2H + (c - 1)f)) \neq 0$.
- (iii) $h^1(E(-H - f)) \neq 0$.

Let us consider one by one the conditions: (i) Let $h^2(E(-2H + (c - 2)f)) \neq 0$, we can conclude that \mathcal{O}_X is a direct summand as in the above theorem. (ii) Let $h^1(E(-2H + (c - 1)f)) \neq 0$. Let us consider the exact sequence:

$$0 \rightarrow E(-2H + (c - 1)f) \rightarrow E(-H + (a_0 - 1)f) \oplus E(-H + (a_1 - 1)f) \rightarrow E(-f) \rightarrow 0$$

Since

$$H^1(E(-H + (a_0 - 1)f)) = H^1(E(-H + (a_1 - 1)f)) = 0,$$

we have a surjective map

$$H^0(E(-f)) \rightarrow H^1(E(-2H + (c - 1)f)).$$

Therefore $H^0(E \otimes \mathcal{O}_X(-f)) \neq 0$ and there exists a non zero map

$$g : \mathcal{O}_X(f) \rightarrow E.$$

On the other hand

$$H^1(E(-2H + (c - 1)f)) \cong H^1(E^\vee(-f))$$

so let us consider the exact sequence

$$0 \rightarrow E^\vee(-f) \rightarrow (E^\vee)^2 \rightarrow E^\vee(f) \rightarrow 0.$$

Since

$$H^1(E^\vee) = H^1(E(-2H + (c - 2)f)) = 0,$$

we have a surjective map

$$H^0(E^\vee(f)) \rightarrow H^1(E(-2H + (c - 1)f)).$$

Therefore $H^0(E^\vee \otimes \mathcal{O}_X(f)) \neq 0$ and there exists a non zero map

$$h : E \rightarrow \mathcal{O}_X(f).$$

Let us consider the following commutative diagram:

$$\begin{array}{ccc} H^1(E(-2H + (c - 1)f)) \otimes H^1(E^\vee(-f)) & \xrightarrow{\sigma} & H^2(E(-2H + (c - 2)f)) \cong \mathbb{C} \\ \downarrow & & \downarrow \\ H^0(E(-f)) \otimes H^1(E^\vee(-f)) & \xrightarrow{\mu} & H^1(\mathcal{O}_X(-f) \otimes \mathcal{O}_X(-f)) \cong \mathbb{C} \\ \downarrow & & \downarrow \\ H^0(E(-f)) \otimes H^0(E^\vee(f)) & \xrightarrow{\tau} & H^0(\mathcal{O}_X(-f) \otimes \mathcal{O}_X(f)) \cong \mathbb{C} \\ \uparrow \cong & & \uparrow \cong \\ \text{Hom}(E, \mathcal{O}_X(f)) \otimes \text{Hom}(\mathcal{O}_X(f), E) & \xrightarrow{\gamma} & \text{Hom}(\mathcal{O}_X(f), \mathcal{O}_X(f)). \end{array}$$

The map σ comes from Serre duality and it is not zero, the right vertical maps are isomorphisms and the left vertical maps are surjective so also the map τ is not zero. This means that the map

$$h \circ g : \mathcal{O}(f) \rightarrow \mathcal{O}(f)$$

is non-zero and hence it is an isomorphism.

This isomorphism shows that $\mathcal{O}(f)$ is a direct summand of E . (iii) Let $h^1(E(-H - f)) \neq 0$. Let us consider the exact sequence:

$$0 \rightarrow E(-H - f) \rightarrow E(-H)^2 \rightarrow E(-H + f) \rightarrow 0.$$

Since $h^1(E(-H)) = 0$ we get $h^0(E(-H + f)) \neq 0$. By Serre duality $h^1(E(-H - f)) = h^1(E^\vee(-H + (c - 1)f))$. By the exact sequence

$$0 \rightarrow E^\vee(-H + (c - 1)f) \rightarrow E^\vee((a_0 - 1)f) \oplus E^\vee((a_1 - 1)f) \rightarrow E^\vee(H - f) \rightarrow 0.$$

Since $h^1(E^\vee((a_0 - 1)f)) = h^1(E(-2H + (a_1 - 1)f)) = 0$ and $h^1(E^\vee((a_1 - 1)f)) = h^1(E(-2H + (a_0 - 1)f)) = 0$ we get also $h^0(E^\vee(H - f)) \neq 0$. By arguing as above we can conclude that $\mathcal{O}_X(H - f)$ is a direct summand of E . (2) \Rightarrow (1). As in [Theorem 3.1](#). \square

These results are useful to investigate ACM and Ulrich bundle on $S(a_0, a_1)$. To this aim, we recall that a bundle E on a surface with hyperplane section H is called ACM if its intermediate cohomology vanishes, i.e. $H^1(E(tH)) = 0$ for any $t \in \mathbb{Z}$ and it is called Ulrich if $H^i(E(-H)) = 0 = H^i(E(-2H))$, for any $i \geq 0$. In particular every Ulrich bundle is ACM.

Remark 3.5. If $c = 2$ we get $a_0 = a_1 = 1$ so $H^1(E(tH + (c - 2)f)) = H^1(E(tH + (a_0 - 1)f)) = H^1(E(tH + (a_1 - 1)f))$ coincide with $H^1(E(tH))$ and we have exactly the classification of the ACM bundles on $\mathbb{P}^1 \times \mathbb{P}^1$ (see [\[16\]](#)). The proof in this case coincides with [\[3\]](#) Theorem 1.4.

Remark 3.6. If $c = 3$ we get $a_0 = 1, a_1 = 2$ so $H^1(E(tH + (c - 2)f)) = H^1(E(tH + (a_1 - 1)f)) = H^1(E(tH + f))$ and $H^1(E(tH + (a_0 - 1)f)) = H^1(E(tH))$. It is well known (see [\[11\]](#)) that the only indecomposable ACM bundles on $X = S(1, 2)$ are $\mathcal{O}_X, \mathcal{O}_X(f), \mathcal{O}_X(2f), \mathcal{O}_X(H - f)$ and the rank two vector bundle obtained as the extension among $\mathcal{O}_X(H - f)$ and $\mathcal{O}_X(2f)$. Since the condition $H^1(E(tH + f)) = 0$ for any integer t is not satisfied by $\mathcal{O}_X(2f)$ we obtain again [Theorem 3.4](#).

Remark 3.7. If $c = 4$ we get $a_0 = 2, a_1 = 2$ or $a_0 = 1, a_1 = 3$ so in the first case $H^1(E(tH + (c - 2)f)) = H^1(E(tH + 2f))$ and $H^1(E(tH + (a_0 - 1)f)) = H^1(E(tH + (a_1 - 1)f)) = H^1(E(tH + f))$ and in the second case $H^1(E(tH + (c - 2)f)) = H^1(E(tH + (a_1 - 1)f)) = H^1(E(tH + 2f))$ and $H^1(E(tH + (a_0 - 1)f)) = H^1(E(tH))$. It is well known (see [\[12\]](#)) the classification of indecomposable ACM bundles on $X = S(2, 2)$ or $X = S(1, 3)$ and it is easy to obtain again [Theorem 3.4](#).

Corollary 3.8. *Let E be an indecomposable vector bundle on X with $\text{Reg}(E) = 0$ and $H^2(E(-2H + (c - 2)f)) = 0$.*

1. *If $h^1(E(-2H + (c - 1)f)) \neq 0$ and $H^1(E(-H)) = H^1(E(-2H + (a_0 - 1)f)) = H^1(E(-2H + (a_1 - 1)f)) = 0$ then $E \cong \mathcal{O}_X(H - f)$.*
2. *$h^1(E(-H - f)) \neq 0$ and $H^1(E(-2H + (c - 2)f)) = H^1(E(-H + (a_0 - 1)f)) = H^1(E(-H + (a_1 - 1)f)) = 0$ then $E \cong \mathcal{O}_X(f)$.*

Proof. Since $E(-H)$ is not regular and $H^2(E(-2H + (c - 2)f)) = 0$, if $H^1(E(-2H + (c - 1)f)) = 0$ then $H^1(E(-H - f)) \neq 0$ and viceversa. So, thanks the other vanishings, by the proof of the above Theorem, we obtain (i) and (ii). \square

For $c > 4$ the family of ACM bundles are too complicated (see [\[12\]](#)) but we can use our notion of regularity to study Ulrich bundles. We need the following Lemmas:

Lemma 3.9. *If E is a globally generated ACM bundle on X , then*

$$H^1(E((a-1)H + bf)) = H^1(E(aH + (b-1)f)) = 0$$

for any $a, b \geq 0$.

Proof. Since E is globally generated we have a surjective map

$$\mathcal{O}_X \rightarrow E \rightarrow 0.$$

Since $h^2(\mathcal{O}_X(-H - f)) = 0$ we obtain $h^2(E(-H - f)) = 0$. Let us consider this exact cohomology sequence:

$$\dots \rightarrow H^1(E(-H)) \rightarrow H^1(E|_f(-H)) \rightarrow H^2(E(-H - f)) \rightarrow \dots$$

Since the first and the third groups vanish by hypothesis, then also the middle group vanishes. As in [Lemma 2.5](#) $H^1(E|_f((a-1)H + bf)) = 0$ for any $a \geq 0$ and for any integer b . This implies $H^1(E((a-1)H + bf)) = 0$ for any $a, b \geq 0$. By sequence (1) tensored by $E(-H - f)$ we get

$$\begin{aligned} 0 \rightarrow E(-2H + (c-1)f) \rightarrow E(-H + (a_0-1)f) \\ \oplus E(-H + (a_1-1)f) \rightarrow E(-f) \rightarrow 0. \end{aligned}$$

Since E is globally generated $h^2(\mathcal{O}_X(-2H + (c-1)f)) = 0$ we obtain $h^2(E(-2H + (c-1)f)) = 0$. Now in the sequence in cohomology

$$\begin{aligned} H^1(E(-H + (a_0-1)f)) \oplus H^1(E(-H + (a_1-1)f)) \\ \rightarrow H^1(E(-f)) \rightarrow H^2(E(-2H + (c-1)f)) = 0 \end{aligned}$$

the first (notice that $a_0 - 1, a_1 - 1 \geq 0$) group vanishes (notice that $a_0 - 1, a_1 - 1 \geq 0$), then also the middle group vanishes. We obtain $H^1(E(aH + (b-1)f)) = 0$ for any $a, b \geq 0$. \square

Lemma 3.10. *If E is an Ulrich bundle on X , then*

- (i) $H^2(E((a-2)H + bf)) = 0$ for any $a, b \geq 0$.
- (ii) $H^1(E((a-1)H + bf)) = H^1(E(aH + (b-1)f)) = 0$ for any $a, b \geq 0$.
- (iii) E is regular.

Proof. Since E is Ulrich we have

$$h^i(E(-H)) = h^i(E(-2H)) = 0$$

for any i . So we obtain (i) as in [Lemma 2.4](#). Since an Ulrich bundle is ACM and globally generated, by the above Lemma (ii) is proved. By (i) and (ii) we obtain the vanishing of [Definition 2.2](#) and hence (iii). \square

Thanks to our notion of regularity and the above we can give a simpler proof of [12] Theorem B without a Beilinson type spectral sequence:

Theorem 3.11 ([12] Theorem B). *An indecomposable E on X is Ulrich if and only if E fits into:*

$$0 \rightarrow \mathcal{O}_X(H - f)^a \rightarrow E \rightarrow \mathcal{O}_X((c-1)f)^b \rightarrow 0, \quad \text{for some } a, b \geq 0. \quad (4)$$

Proof. By (iii) of [Lemma 3.10](#), E is regular and since $h^0(E(-H)) = 0$, $E(-H)$ must be not regular. By (i) and (ii) of [Lemma 3.10](#), $h^2(E(-2H + (c - 2)f)) = 0$ and we may conclude that one of the following conditions is satisfied:

- (α) $h^1(E(-H - f)) \neq 0$.
- (β) $h^1(E(-2H + (c - 1)f)) \neq 0$.

(α) Let $h^1(E(-H - f)) = a \neq 0$. Let us consider the exact sequence:

$$0 \rightarrow E(-H - f) \rightarrow E(-H) \rightarrow E(-H + f) \rightarrow 0.$$

Since $h^1(E(-H)) = h^0(E(-H)) = 0$ we get $h^0(E(-H + f)) = a$.

So there exists a map

$$h : \mathcal{O}(H - f)^a \rightarrow E.$$

Let $b = h^1(E(-2H + f))$. We distinguish two cases: $b = 0$ and $b \neq 0$.

Let assume first $b = 0$. By Serre duality $h^1(E(-H - f)) = h^1(E^\vee(-H + (c - 1)f)) = a$. From

$$\dots \rightarrow H^1(E(-2H + f)) \rightarrow H^1(E|_f(-2H + f)) \rightarrow H^2(E(-H)) \rightarrow \dots$$

since the first ($b = 0$) and the third groups vanish by hypothesis, then also the middle group vanishes. As in [Lemma 2.5](#) $H^1(E|_f((a - 2)H + bf)) = 0$ for any $a \geq 0$ and for any integer b . This implies $H^1(E((a - 1)H + bf)) = 0$ for any $a, b \geq 0$. In particular $h^1(E^\vee((a_0 - 1)f)) = h^1(E(-2H + (a_1 - 1)f)) = 0$ and $h^1(E^\vee((a_1 - 1)f)) = h^1(E(-2H + (a_0 - 1)f)) = 0$ so, from the exact sequence

$$0 \rightarrow E^\vee(-H + (c - 1)f) \rightarrow E^\vee((a_0 - 1)f) \oplus E^\vee((a_1 - 1)f) \rightarrow E^\vee(H - f) \rightarrow 0,$$

we get also $h^0(E^\vee(H - f)) \neq 0$. Hence as in [Theorem 3.4](#) we obtain $E \cong \mathcal{O}_X(H - f)$.

Let assume now $b > 0$. By [\[8\]](#) we may assume that the kernel K and the cokernel G of h are also Ulrich. So we obtain two exact sequences with also U Ulrich:

$$0 \rightarrow U \rightarrow E \rightarrow G \rightarrow 0,$$

and

$$0 \rightarrow K \rightarrow \mathcal{O}_X(H - f)^a \rightarrow U \rightarrow 0. \tag{5}$$

Notice that by [Lemma 3.10](#) and sequence (1) tensored by $U(-2H - f)$ we get $h^2(U(-H - f)) = 0$. So if we twist the sequence (5) by $-H - f$, since $h^0(U(-H - f)) = h^2(U(-H - f)) = 0$ and $h^1(\mathcal{O}_X(-2f)^a) = a$, we have $h^1(K(-H - f)) \neq 0$. Now let us consider the sequence in cohomology:

$$H^0(U(-2H + f)) \rightarrow H^1(K(-2H + f)) \rightarrow H^1(\mathcal{O}_X(-H)^a)$$

Since $H^0(U(-2H + f)) = H^1(\mathcal{O}_X(-H)^a) = 0$, we get $H^1(K(-2H + f)) = 0$ and by the argument above for $b = 0$ we may conclude that $\mathcal{O}_X(H - f)$ is a direct summand of K . By iterating this argument we get the sequence (5) simply becomes

$$0 \rightarrow \mathcal{O}_X(H - f)^{a'} \rightarrow \mathcal{O}_X(H - f)^a \rightarrow \mathcal{O}_X(H - f)^{a-a'} \rightarrow 0$$

for a suitable positive integer a' . Hence we may assume that h is injective. Let us denote by G the cokernel of h :

$$0 \rightarrow \mathcal{O}_X(H - f)^a \rightarrow E \rightarrow G \rightarrow 0.$$

Notice that $h^1(G(-H - f)) = 0$. If we twist the above exact sequence by $\mathcal{O}_X(-2H + tf)$, since $h^i(\mathcal{O}_X(-H + tf)) = 0$ for any i, t we get $b = h^1(G(-2H + f))$ and $h^1(G(-2H + tf)) = h^1(E(-2H + tf))$ for any integer t .

From

$$0 \rightarrow G(-H - 2f) \rightarrow G(-H - f)^2 \rightarrow G(-H) \rightarrow 0,$$

we obtain $h^1(G(-H - tf)) = 0$ for any $t \geq 0$.

Let us consider the exact sequence:

$$\begin{aligned} 0 \rightarrow G(-2H + f) \rightarrow G(-H + (-a_0 + 1)f) \\ \oplus G(-H + (-a_1 + 1)f) \rightarrow G(-(c - 1)f) \rightarrow 0 \end{aligned}$$

Since

$$H^i(G(-H + (-a_0 + 1)f)) = H^i(G(-H + (-a_1 + 1)f)) = 0,$$

for any i , we have $H^0(G \otimes \mathcal{O}_X(-(c - 1)f)) = b \neq 0$. On the other hand

$$H^1(G(-2H + f)) \cong H^1(G^\vee((c - 3)f))$$

so let us consider the exact sequence

$$0 \rightarrow G^\vee((c - 3)f) \rightarrow 2G^\vee((c - 2)f) \rightarrow G^\vee((c - 1)f) \rightarrow 0.$$

Since

$$H^1(G^\vee((c - 2)f)) = H^1(G(-2H)) = 0,$$

we get $H^0(G^\vee \otimes \mathcal{O}_X((c - 1)f)) = b \neq 0$. So by arguing as in [Theorem 3.4](#) we obtain $G \cong \mathcal{O}_X((c - 1)f)^b$ and E fits in (4).

(β) Notice that if E is Ulrich also $E' = E^\vee(H + (c - 2)f)$ is Ulrich. The condition

$$h^1(E(-2H + (c - 1)f)) = b \neq 0$$

by Serre duality corresponds to

$$\begin{aligned} h^1(E^\vee(-f)) &= h^1(E^\vee(H + (c - 2)f) \otimes \mathcal{O}_X(-H - (c - 1)f)) \\ &= h^1(E'(-H - (c - 1)f)) \neq 0. \end{aligned}$$

Let us consider for any integer t the exact sequence

$$0 \rightarrow E'(-H - (t + 2)f) \rightarrow E'(-H - (t + 1)f) \rightarrow E'(-H - tf) \rightarrow 0.$$

If $t \geq 0$ the map

$$H^1(E'(-H - (t + 2)f)) \rightarrow H^1(E'(-H - (t + 1)f))$$

is injective. Since $h^1(E'(-H - (c - 1)f)) \neq 0$, we have $h^1(E'(-H - (c - 2)f)) \neq 0$ and by a recursive argument $h^1(E'(-H - tf)) \neq 0$ for $-(c - 1) \leq t \leq 1$. In particular $h^1(E'(-H - f)) \neq 0$, so we may repeat the argument of the case (α) for the Ulrich bundle E' and we obtain $E' \cong \mathcal{O}_X(H - f)$ (hence $E \cong \mathcal{O}_X((c - 1)f)$) or E' fits in the extension

$$0 \rightarrow \mathcal{O}_X(H - F)^b \rightarrow E' \rightarrow \mathcal{O}_X((c - 1)f)^a \rightarrow 0$$

that dualized and tensored by $\mathcal{O}_X(H + (c - 2)f)$ becomes (4). \square

Remark 3.12. If $c = 2$, $\dim(\text{Ext}^1(\mathcal{O}_X((c-1)f), \mathcal{O}_X(H-f))) = 0$ so (4) splits.

If $c = 3$, $\dim(\text{Ext}^1(\mathcal{O}_X((c-1)f), \mathcal{O}_X(H-f))) = 1$ so from (4) we only obtain a unique rank two indecomposable Ulrich bundle.

If $c = 4$, $\dim(\text{Ext}^1(\mathcal{O}_X((c-1)f), \mathcal{O}_X(H-f))) = 2$ so from (4) we only obtain families of dimension at most one of indecomposable Ulrich bundles (see [12]).

If $c > 4$, $\dim(\text{Ext}^1(\mathcal{O}_X((c-1)f), \mathcal{O}_X(H-f))) > 2$ so from (4) we may obtain arbitrary large families of indecomposable Ulrich bundles (see [12]).

4. Logarithmic bundle on $S(a_0, a_1)$

In this section we will show how the notion of regularity introduced here can be useful also in the study of logarithmic bundles of some families of divisors on rational normal scrolls.

Definition 4.1. A divisor D on a non-singular variety X is said to have normal crossings if $\mathcal{O}_{D,x}$ is formally isomorphic to the quotient of $\mathcal{O}_{X,x}$ by an ideal generated by t_1, \dots, t_k , where t_1, \dots, t_k is a subset of the set of local parameters in $\mathcal{O}_{X,x}$ for all $x \in D$. D is also said to have simple normal crossings if it is the union of smooth divisors $D_i, i = 1, \dots, m$, which intersect transversely at each point.

Definition 4.2. An arrangement on X is defined to be a set $D = \{D_1, \dots, D_m\}$ of smooth irreducible divisors of X such that $D_i \neq D_j$ for $i \neq j$. To an arrangement D on X , we can associate the logarithmic sheaf $\Omega_X^1(\log D)$, the sheaf of differential 1-forms with logarithmic poles along D .

If D has simple normal crossings, its logarithmic sheaf is known to be locally free and so it can be called to be the logarithmic bundle. It admits the residue exact sequence

$$0 \rightarrow \Omega_X^1 \rightarrow \Omega_X^1(\log D) \xrightarrow{res} \bigoplus \epsilon_i^* \mathcal{O}_{D_i} \rightarrow 0. \tag{6}$$

From now on, let X be $S(a_0, a_1)$ and let $e = a_1 - a_0 = -C_0^2$. Let us consider the lines $L_i \in |\mathcal{O}_X(f)|$. Recall (see [14] II 8.11.) that the cotangent bundle of X is given in

$$0 \rightarrow \mathcal{O}_X(-2f) \rightarrow \Omega_X^1 \rightarrow \mathcal{O}_X(-2H + cf) \rightarrow 0,$$

and this extension splits only if $e = 0$:

Proposition 4.3. Let $D = \{L_1, \dots, L_a\}$ be an arrangement of a lines on X with $L_i \in |\mathcal{O}_X(f)|$. Then we have

$$\Omega_X^1(\log D) \cong \mathcal{O}_X((a-2)f) \oplus \mathcal{O}_X(-2H + cf),$$

if $a \geq e + 1$.

Proof. Let $D = \{L_1\}$. We apply the covariant functor $\text{Hom}(\mathcal{O}_{L_1}, -)$ to vertical column of the diagram

$$\begin{array}{ccccccc}
 & & 0 & & & & \\
 & & \downarrow & & & & \\
 & & \mathcal{O}_X(-2f) & & & & \\
 & & \downarrow & & & & \\
 0 \rightarrow & & \Omega_X^1 & \rightarrow & \Omega_X^1(\log D) & \rightarrow & \mathcal{O}_{L_1} \rightarrow 0 \\
 & & \downarrow & & & & \\
 & & \mathcal{O}_X(-2H + cf) & & & & \\
 & & \downarrow & & & & \\
 & & 0 & & & &
 \end{array}$$

As the dimension of $\text{Ext}^1(\mathcal{O}_{L_1}, \mathcal{O}_X(-2H + cf))$ is $h^1(\mathcal{O}_{L_1} \otimes \mathcal{O}_X(-2f)) = h^1(\mathcal{O}_{\mathbb{P}^1}) = 0$ and the dimension of $\text{Ext}^0(\mathcal{O}_{L_1}, \mathcal{O}_X(-2H + cf))$ is $h^2(\mathcal{O}_{L_1} \otimes \mathcal{O}_X(-2f)) = h^2(\mathcal{O}_{\mathbb{P}^1}) = 0$, we get

$$\text{Ext}^1(\mathcal{O}_{L_1}, \Omega_X^1) \cong \text{Ext}^1(\mathcal{O}_{L_1}, \mathcal{O}_X(-2f))$$

and their dimension is $h^1(\mathcal{O}_{L_1} \otimes \mathcal{O}_X(-2H + cf)) = h^1(\mathcal{O}_{L_1}(-2)) = h^1(\mathcal{O}_{\mathbb{P}^1}(-2)) = 1$. So we get the unique extension (note that $\mathcal{O}_{L_1} \otimes \mathcal{O}_X(-f) \cong \mathcal{O}_{L_1}$) $0 \rightarrow \mathcal{O}_X(-2f) \rightarrow \mathcal{O}_X(-f) \rightarrow \mathcal{O}_{L_1} \rightarrow 0$ to close the following diagram

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 0 \rightarrow & & \mathcal{O}_X(-2f) & \rightarrow & \mathcal{O}_X(-f) & \rightarrow & \mathcal{O}_{L_1} \rightarrow 0 \\
 & & \downarrow & & \downarrow & & \parallel \\
 0 \rightarrow & & \Omega_X^1 & \rightarrow & \Omega_X^1(\log D) & \rightarrow & \mathcal{O}_{L_1} \rightarrow 0 \\
 & & \downarrow & & \downarrow & & \\
 & & \mathcal{O}(-2H + cf) & = & \mathcal{O}(-2H + cf) & & \\
 & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & &
 \end{array}$$

Now, let $a \geq 2$, $D = L_1, \dots, L_a$, $D' = L_1, \dots, L_{a-1}$ and the assertion true for $a - 1$ to argue by induction. Similarly, we get the diagram

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 0 \rightarrow & & \mathcal{O}_X((a-3)f) & \rightarrow & \mathcal{O}_X((a-2)f) & \rightarrow & \mathcal{O}_{L_1} \rightarrow 0 \\
 & & \downarrow & & \downarrow & & \parallel \\
 0 \rightarrow & & \Omega_X^1(\log((a-1)f)) & \rightarrow & \Omega_X^1(\log(af)) & \rightarrow & \mathcal{O}_{L_1} \rightarrow 0 \\
 & & \downarrow & & \downarrow & & \\
 & & \mathcal{O}_X(-2H + cf) & = & \mathcal{O}_X(-2H + cf) & & \\
 & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & &
 \end{array}$$

in fact, the dimension of $\text{Ext}^1(\mathcal{O}_{L_1}, \mathcal{O}_X((a-3)f))$ is $h^1(\mathcal{O}_{L_1} \otimes \mathcal{O}_X(-2H + (c-e)f)) = h^1(\mathcal{O}_{L_1}(-2)) = h^1(\mathcal{O}_{\mathbb{P}^1}(-2)) = 1$ and the dimension of $\text{Ext}^1(\mathcal{O}_{L_1}, \mathcal{O}_X(-2H + cf))$ is $h^1(\mathcal{O}_{L_1} \otimes \mathcal{O}_X(-2f)) = 0$. In order to have a splitting sequence in the second column, it is enough that $\text{Ext}^1(\mathcal{O}_X(-2H + cf), \mathcal{O}_X((a-2)f)) = 0$. We have that the dimension of $\text{Ext}^1(\mathcal{O}_X(-2H + cf), \mathcal{O}_X((a-2)f))$ is $h^1(\mathcal{O}_X(2H + (a-2-c)f))$ which is zero if $h^1(\mathcal{O}_{\mathbb{P}^1}(2a_0 + a - 2 - c)) = 0$. This implies $2a_0 + a - c \geq 1$, hence $a \geq e + 1$. \square

Proposition 4.4. *Let $D = \{L_1, \dots, L_a, C_1\}$ be an arrangement of $a \geq e + 1$ lines and one rational curve on X with $L_i \in |\mathcal{O}_X(f)|$ and $C_1 \in |\mathcal{O}_X(H - a_1f)|$. Then we have*

$$\Omega_X^1(\log D) \cong \mathcal{O}_X((a - 2)f) \oplus \mathcal{O}_X(-H + a_0f).$$

Proof. In the proof we will use that the dimension of $Ext^1(\mathcal{O}_{C_1}, \mathcal{O}_X((a - 2)f))$ is $h^1(\mathcal{O}_{C_1} \otimes \mathcal{O}_X(-2H + (c - a)f)) = h^1(\mathcal{O}_{\mathbb{P}^1}(-2c + (c - a + 2a_1))) = h^1(\mathcal{O}_{\mathbb{P}^1}(-c + 2a_1 - a)) = h^1(\mathcal{O}_{\mathbb{P}^1}(e - a))$, for any a . Let us first consider the case of $D = \{L_1, \dots, L_{e+1}, C_1\}$. Let $D' = \{L_1, \dots, L_{e+1}\}$. Then we have the sequence

$$0 \rightarrow \Omega_X^1(\log D') \rightarrow \Omega_X^1(\log D) \rightarrow \mathcal{O}_{C_1} \rightarrow 0.$$

Thanks to Proposition 4.3

$$Ext^1(\mathcal{O}_{C_1}, \mathcal{O}_X(-2H + cf))\Omega_X^1(\log D') \cong \mathcal{O}_X((e - 1)f) \oplus \mathcal{O}_X(-2H + cf).$$

Now, as noted before, $Ext^1(\mathcal{O}_{C_1}, \mathcal{O}_X((a - 2)f))$ vanishes if $a \leq e + 1$ and the dimension of $Ext^1(\mathcal{O}_{C_1}, \mathcal{O}_X(-2H + cf))$ is $h^1(\mathcal{O}_{C_1} \otimes \mathcal{O}_X(-2f)) = h^1(\mathcal{O}_{\mathbb{P}^1}(-2)) = 1$ and we get the unique extension

$$0 \rightarrow \mathcal{O}_X(-2H + cf) \rightarrow \mathcal{O}_X(-H + (c - a_1)f) \rightarrow \mathcal{O}_{C_1} \rightarrow 0.$$

Thus there exists a uniquely determined extension of \mathcal{O}_{C_1} by $\Omega_X^1(\log D')$ and it must be $\mathcal{O}_X((e - 1)f) \oplus \mathcal{O}_X(-H + a_0f)$.

Now assume that the assertion is true for $a \geq e + 1$ to use induction. For $D = \{L_1, \dots, L_{a+1}, C_1\}$, if $D' = \{L_1, \dots, L_a, C_1\}$ we have the sequence

$$0 \rightarrow \Omega_X^1(\log D') \rightarrow \Omega_X^1(\log D) \rightarrow \mathcal{O}_{L_{a+1}} \rightarrow 0.$$

Thanks to the above argument

$$\Omega_X^1(\log D') \cong \mathcal{O}_X((a - 2)f) \oplus \mathcal{O}_X(-H + a_0f),$$

As before, the dimension of $Ext^1(\mathcal{O}_{L_{a+1}}, \mathcal{O}_X((a - 2)f))$ is $h^1(\mathcal{O}_{L_{a+1}} \otimes \mathcal{O}_X(-2H + (c - a)f)) = h^1(\mathcal{O}_{\mathbb{P}^1}(-2)) = 1$ and the dimension of $Ext^1(\mathcal{O}_{L_{a+1}}, \mathcal{O}_X(-H + a_0f))$ is $h^1(\mathcal{O}_{L_{a+1}} \otimes \mathcal{O}_X(-H + (a_1 - 2)f)) = h^1(\mathcal{O}_{\mathbb{P}^1}(-1)) = 0$ and we get the unique extension

$$0 \rightarrow \mathcal{O}_X((a - 2)f) \rightarrow \mathcal{O}_X((a - 1)f) \rightarrow \mathcal{O}_{L_{a+1}} \rightarrow 0.$$

Thus there exists a uniquely determined extension of $\mathcal{O}_{L_{a+1}}$ by $\Omega_X^1(\log D')$ and it must be $\mathcal{O}_X((a - 1)f) \oplus \mathcal{O}_X(-H + a_0f)$. \square

If $e > 0$, $h^0(\mathcal{O}_X(H - a_1f)) = 1$ and we cannot consider an arrangement with more than a curve $C_j \in |\mathcal{O}_X(H - a_1f)|$. When $e = 0$, $h^0(\mathcal{O}_X(H - a_1f)) = 2$ and a curve $C \in |\mathcal{O}_X(H - a_1f)|$ is rational of degree a_0 . Moreover $(H - a_1f)^2 = 0$ so we have a one dimensional family of disjoint lines in $|\mathcal{O}_X(f)|$ and a one dimensional family of disjoint rational curve of degree a_0 . For this reason we consider now in more detail the case $e = 0$.

Theorem 4.5. *Let $e = 0$. Let $D = \{L_1, \dots, L_a, C_1, \dots, C_b\}$ be an arrangement of $a \geq 1$ lines and $b \geq 1$ rational curves on X with $L_i \in |\mathcal{O}_X(f)|$ and $C_j \in |\mathcal{O}_X(H - a_1f)|$. Then we have*

$$\Omega_X^1(\log D) \cong \mathcal{O}_X((a - 2)f) \oplus \mathcal{O}_X((b - 2)H + (c - ba_1)f).$$

Proof. Let us first consider the case of $D = \{L_1, C_1, C_2\}$. Let $D' = \{L_1, C_1\}$. Then we have the sequence

$$0 \rightarrow \Omega_X^1(\log D') \rightarrow \Omega_X^1(\log D) \rightarrow \mathcal{O}_{C_2} \rightarrow 0.$$

Thanks to [Proposition 4.4](#)

$$\Omega_X^1(\log D') \cong \mathcal{O}_X(-f) \oplus \mathcal{O}_X(-H + a_0f).$$

Note that the dimension of $Ext^1(\mathcal{O}_{C_2}, \mathcal{O}_X(-f))$ is $h^1(\mathcal{O}_{C_2} \otimes \mathcal{O}_X(-2H + (c - 1)f)) = h^1(\mathcal{O}_{\mathbb{P}^1}(-2c + c - 1 + 2a_1)) = h^1(\mathcal{O}_{\mathbb{P}^1}(-c - 1 + 2a_1)) = 0$ and the dimension of $Ext^1(\mathcal{O}_{C_2}, \mathcal{O}_X(-H + a_0f))$ is $h^1(\mathcal{O}_{C_2} \otimes \mathcal{O}_X(-H + (c - 2 - a_0)f)) = h^1(\mathcal{O}_{\mathbb{P}^1}(-c + c - 2 + a_1 - a_0)) = 1$ and we get the unique extension

$$0 \rightarrow \mathcal{O}_X(-H + a_1f) \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_{C_2} \rightarrow 0.$$

Thus there exists a uniquely determined extension of \mathcal{O}_{C_2} by $\Omega_X^1(\log D')$ and it must be $\mathcal{O}_X(-f) \oplus \mathcal{O}_X$.

Now assume that the assertion is true for $(1, b)$ with $b \geq 2$ to use induction. For the case of $(1, b + 1)$ $D = \{L_1, C_1, \dots, C_{b+1}\}$. Let $D' = \{L_1, C_1, \dots, C_b\}$, then we have the sequence

$$0 \rightarrow \Omega_X^1(\log D') \rightarrow \Omega_X^1(\log D) \rightarrow \mathcal{O}_{C_{b+1}} \rightarrow 0.$$

Thanks to above argument and the inductive hypothesis,

$$\Omega_X^1(\log D') \cong \mathcal{O}_X(-f) \oplus \mathcal{O}_X((b - 2)H + (c - ba_1)f).$$

Let us tensor the above sequence by $\mathcal{O}_X(f)$ and we obtain

$$0 \rightarrow \mathcal{O}_X \oplus \mathcal{O}_X((b - 2)H + (c - ba_1 + 1)f) \rightarrow \Omega_X^1(\log D) \otimes \mathcal{O}_X(f) \rightarrow \mathcal{O}_{C_{b+1}} \otimes \mathcal{O}_X(f) \rightarrow 0.$$

We call $E = \Omega_X^1(\log D) \otimes \mathcal{O}_X(f)$ and we want to show that E is regular. Notice that $\mathcal{O}_X \oplus \mathcal{O}_X((b - 2)H + (c - ba_1 + 1)f)$ is regular. So $h^1(E(-f)) = h^1(\mathcal{O}_{C_{b+1}} \otimes \mathcal{O}_X) = h^1(\mathcal{O}_{\mathbb{P}^1}) = 0$. Moreover $h^1(E(-H + (c - 1)f)) = h^1(\mathcal{O}_{C_{b+1}} \otimes \mathcal{O}_X(-H + cf)) = h^1(\mathcal{O}_{\mathbb{P}^1}(-c + c + a_1 + c)) = h^1(\mathcal{O}_{\mathbb{P}^1}(a_1 + c)) = 0$ and $h^2(E(-H + (c - 2)f)) = h^2(\mathcal{O}_{C_{b+1}} \otimes \mathcal{O}_X(-H + (c - 1)f)) = 0$. Thus we have that E is regular and, since $h^1(\mathcal{O}_{C_{b+1}} \otimes \mathcal{O}_X(-2H + (c - 1)f)) = h^1(\mathcal{O}_{\mathbb{P}^1}(-2c + 2a_1 + c - 1)) = 0$, we get $h^2(E(-2H + (c - 2)f)) = h^0(E) \geq 1$ so, by [Corollary 3.3](#), we can conclude that \mathcal{O}_X is a direct summand of E . Hence, if E is a vector bundle, $E \cong \mathcal{O}_X \oplus \mathcal{O}_X((b - 1)H + (c - (b + 1)a_1 + 1)f)$ and

$$\Omega_X^1(\log D) \cong \mathcal{O}_X(-f) \oplus \mathcal{O}_X((b - 1)H + (c - (b + 1)a_1)f).$$

Finally let us deal with the case when a and b are at least 1 and b at least 2. The logarithmic bundle $\Omega_X^1(\log D)$ is an extension of $(\oplus \mathcal{O}_{L_i}) \oplus (\oplus \mathcal{O}_{C_j})$ by $\mathcal{O}_X(-2f) \oplus \mathcal{O}_X(-H + a_0f)$. Note that we have

$$Ext^1(\oplus \mathcal{O}_{L_i}, \mathcal{O}_X(-H + a_0f)) = 0$$

by [Proposition 4.4](#) and

$$Ext^1(\oplus \mathcal{O}_{C_j}, \mathcal{O}_X(-2f)) = 0$$

by the above argument.

Thus $\Omega_X^1(\log D)$ corresponds to an element ϵ ;

$$\epsilon \in Ext^1(\oplus \mathcal{O}_{L_i}, \mathcal{O}_X(-2f)) \oplus Ext^1(\oplus \mathcal{O}_{C_j}, \mathcal{O}_X(-H + a_0f)).$$

From the argument in Proposition 4.4, we observe that the first factor of ϵ with

$$\text{Ext}^1(\oplus \mathcal{O}_{L_i}, \mathcal{O}_X(-H + a_0f)) = 0$$

generates $\mathcal{O}_X((a - 2)f) \oplus \mathcal{O}_X(-H + a_0f)$ and similarly, by the argument for the case $(0, b)$ the second factor generates $\mathcal{O}_X(-2f) \oplus \mathcal{O}_X((b - 2)H + (c - ba_1)f)$. Thus ϵ corresponds to the bundle $\mathcal{O}_X((a - 2)f) \oplus \mathcal{O}_X((b - 2)H + (c - ba_1)f)$. \square

Remark 4.6. When $c = 2$ the above Theorem coincides with [2] Proposition 6.3.

We are able finally to classify regular ACM logarithmic bundles:

Corollary 4.7. *Let $e = 0$ and $c > 2$. Let D be an arrangement of smooth curves on X with simple normal crossings. If $\Omega_X^1(\log D)$ is a regular ACM bundle, then D consists of a lines in $|\mathcal{O}_X(f)|$ with $2 \leq a \leq c + 1$ and 2 rational curves in $|\mathcal{O}_X(H - a_1f)|$. In particular we have that $\Omega_X^1(\log D)$ has always regularity 0:*

$$\Omega_X^1(\log D) \cong \mathcal{O}_X((a - 2)f) \oplus \mathcal{O}_X.$$

Proof. If $D := \{D_1, \dots, D_m\}$ consists of m smooth curves, then it admits the sequence

$$0 \rightarrow \mathcal{O}_X(-2f) \oplus \mathcal{O}_X(-2H + cf) \rightarrow \Omega_X^1(\log D) \rightarrow \oplus_{i=1}^m \mathcal{O}_{D_i} \rightarrow 0. \quad (7)$$

If $E = \Omega_X^1(\log D)$ is regular, then we have $h^1(E(-f)) = 0$. From the sequence (7) twisted by $\mathcal{O}_X(-f)$ and the fact that $h^2(\mathcal{O}_X(-3f)) = h^2(\mathcal{O}_X(-2H + (c - 1)f)) = 0$, we deduce that $h^1(\mathcal{O}_{D_i} \otimes \mathcal{O}_X(-f)) = 0$ for any $i = 1, \dots, m$. Let $D_i \in |\mathcal{O}_X(s_iH + t_i f)|$ with $s_i \geq 0$ and $t_i \geq -a_1$, we get $h^1(\mathcal{O}_{D_i} \otimes \mathcal{O}_X(-f)) = h^1(\mathcal{O}_{\mathbb{P}^1}(-s_i)) = 0$, so $s_i = 0$ or $s_i = 1$.

Moreover by Lemma 3.9 $H^1(E((a - 1)H + bf)) = H^1(E(aH + (b - 1)f)) = 0$ for any $a, b \geq 0$, in particular we have $h^1(E(-H + (a_1 - 1)f)) = 0$. From the sequence (7) twisted by $\mathcal{O}_X(-H + (a_1 - 1)f)$ and the fact that $h^2(\mathcal{O}_X(-H + (a_1 - 3)f)) = 0$ and $h^2(\mathcal{O}_X(-3H + (c + a_1 - 1)f)) = h^0(\mathcal{O}_X(H + (-1 - a_1)f)) = 0$, we deduce that $h^1(\mathcal{O}_{D_i} \otimes \mathcal{O}_X(-H + (a_1 - 1)f)) = 0$ for any $i = 1, \dots, m$. We get $h^1(\mathcal{O}_{D_i} \otimes \mathcal{O}_X(-H + (a_1 - 1)f)) = h^1(\mathcal{O}_{\mathbb{P}^1}(-cs_i + (a_1 - 1)s_i - t_i)) = 0$. If $s_i = 0$ we obtain $t_i = 1$ so $D_i \in |\mathcal{O}_X(f)|$. If $s_i = 1$ we must have $-c + a_1 - 1 - t_i \geq -1$, hence $t_i \leq -a_0$. Since $a_0 = a_1$ and $t_i \geq -a_1$ we may conclude that $t_i = -a_1$. So we have only two cases: $D_i \in |\mathcal{O}_X(f)|$ or $D_i \in |\mathcal{O}_X(H - a_1f)|$. By Theorem 4.5

$$\Omega_X^1(\log D) \cong \mathcal{O}_X((a - 2)f) \oplus \mathcal{O}_X((b - 2)H + (c - ba_1)f).$$

We recall that $\mathcal{O}_X(sH + tf)$ is ACM if and only if $-1 \leq t \leq c - 1$ so we must have $2 \leq a \leq c + 1$ and (since $c > 2$) $b = 2$. \square

Acknowledgments

The second author is partially supported by PRIN project ‘‘Multilinear Algebraic Geometry’’ of MUR (2022ZRRL4C).

References

- [1] E. Arrondo, F. Malaspina, Cohomological characterization of vector bundles on grassmannians of lines, J. Algebra 323 (2010) 1098–1106.
- [2] E. Ballico, S. Huh, F. Malaspina, A Torelli-type problem for logarithmic bundles over projective varieties, Q. J. Math. (2015) 417–436.

- [3] E. Ballico, F. Malaspina, Q-regularity and an extension of Evans-Griffiths criterion to vector bundles on quadrics, *J. Pure Appl. Algebra* 213 (2009) 194–202.
- [4] E. Ballico, F. Malaspina, Regularity and cohomological splitting conditions for vector bundles on multiprojective spaces, *J. Algebra* 345 (2011) 137–149.
- [5] N. Botbol, M. Chardin, Castelnuovo mumford regularity with respect to multigraded ideals, *J. Algebra* 474 (2017) 361–392.
- [6] M.K. Brown, D. Erman, Tate resolutions on toric varieties, *J. Eur. Math. Soc. (JEMS)* <http://dx.doi.org/10.4171/JEMS/1474>.
- [7] M.K. Brown, M. Sayrafi, A short resolution of the diagonal for smooth projective toric varieties of picard rank 2, *Algebra Number Theory* 18-10 (2024) 1923–1943.
- [8] M. Casanellas, R. Hartshorne, F. Geiss, F.O. Schreyer, Stable Ulrich bundles, *Int. J. Math.* 23 (2012) 1250083.
- [9] L. Costa, R.M. Miró-Roig, m - blocks collections and Castelnuovo–Mumford regularity in multiprojective spaces, *Nagoya Math. J.* 186 (2007) 119–155.
- [10] D. Eisenbud, J. Harris, On varieties of minimal degree (a centennial account), in: *Algebraic geometry, Bowdoin, 1985* (Brunswick, Maine, 1985). *Proceedings of Symposia in Pure Mathematics*, Vol. 46, No. 1, 1987, pp. 3–13.
- [11] D. Eisenbud, J. Herzog, The classification of homogeneous Cohen–Macaulay rings of finite representation type, *Math. Ann.* 280 (1988) 347–352.
- [12] D. Faenzi, F. Malaspina, Surfaces of minimal degree of tame representation type and mutations of Cohen–Macaulay modules, *Adv. Math.* 310 (2017) 663–695.
- [13] M. Fulger, M. Marchitan, Some splitting criteria on Hirzebruch surfaces, *Bull. math. de la Société des Sciences Math. Roumanie* 4 (2011) 313–323, <http://www.jstor.org/stable/43679229>.
- [14] R. Hartshorne, Algebraic geometry, in: *G.T.M.*, vol. 52, Springer, 1977.
- [15] J.W. Hoffman, H.H. Wang, Castelnuovo–Mumford regularity in biprojective spaces, *Adv. Geom.* 4 (4) (2004) 513–536.
- [16] H. Knörrer, Cohen–Macaulay modules of hypersurface singularities I, *Invent. Math.* 88 (1987) 153–164; *Positivity in Algebraic Geometry*, vol. 2, Springer, Berlin, 2004.
- [17] D. Maclagan, G. Smith, Multigraded Castelnuovo–Mumford regularity, *J. Reine Angew. Math.* 571 (2004) 179–212.
- [18] D. Mumford, Lectures on Curves on an Algebraic Surface, in: *Annals of Mathematics Studies*, vol. 59, Princeton University Press, Princeton, N.J, 1966.
- [19] G. Pareschi, M. Popa, Regularity on abelian varieties I, *J. Amer. Math. Soc.* 16 (2003) 285–302.
- [20] Deligne Pierre, Équations différentielles à points singuliers réguliers, in: *Lecture Notes in Mathematics*, vol. 163, Springer-Verlag, 1970.
- [21] K. Saito, Theory of logarithmic differential forms and logarithmic vector fields, *J. Fac. Sci. Univ. Tokyo* 27 (1980) 265–291.