

Higher order Schauder estimates for degenerate or singular parabolic equations

Original

Higher order Schauder estimates for degenerate or singular parabolic equations / Audrito, A., Fioravanti, G., Vita, S.. - In: REVISTA MATEMATICA IBEROAMERICANA. - ISSN 0213-2230. - 41:4(2025), pp. 1513-1554. [10.4171/rmi/1540]

Availability:

This version is available at: 11583/2999694 since: 2025-04-30T08:00:50Z

Publisher:

EMS press

Published


DOI:10.4171/rmi/1540

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)



Higher order Schauder estimates for degenerate or singular parabolic equations

Alessandro Audrito, Gabriele Fioravanti and Stefano Vita

Abstract. In this paper, we complete the analysis initiated in [Calc. Var. Partial Differential Equations 63 (2024), article no. 204] establishing some higher order $C^{k+2,\alpha}$ Schauder estimates ($k \in \mathbb{N}$) for a class of parabolic equations with weights that are degenerate/singular on a characteristic hyperplane. The $C^{2,\alpha}$ -estimates are obtained through a blow-up argument and a Liouville theorem, while the higher order estimates are obtained by a fine iteration procedure. As a byproduct, we present two applications. First, we prove similar Schauder estimates when the degeneracy/singularity of the weight occurs on a regular hypersurface of cylindrical type. Second, we provide an alternative proof of the higher order boundary Harnack principles established in [J. Differential Equations 260 (2016), 1801–1829] and [Discrete Contin. Dyn. Syst. 42 (2022), 2667–2698].

1. Introduction

In this paper we complete the study started in [4], establishing some higher order Schauder regularity estimates for solutions to a special class of parabolic equations having weights which degenerate or explode on a characteristic hyperplane Σ as $\text{dist}(\cdot, \Sigma)^a$, where $a > -1$ is a fixed parameter. More precisely, for every $k \in \mathbb{N}$, we prove local regularity estimates in $C_p^{k+2,\alpha}$ (parabolic Hölder) spaces “up to” Σ for weak solutions to

$$(1.1) \quad \begin{cases} y^a \partial_t u - \text{div}(y^a A \nabla u) = y^a f + \text{div}(y^a F) & \text{in } Q_1^+, \\ \lim_{y \rightarrow 0^+} y^a (A \nabla u + F) \cdot e_{N+1} = 0 & \text{on } \partial^0 Q_1^+. \end{cases}$$

Here $N \geq 1$, $(z, t) = (x, y, t) \in \mathbb{R}^N \times \mathbb{R} \times \mathbb{R}$, $\Sigma = \{y = 0\}$ and $\text{dist}(P, \Sigma)^a = y^a$. Further, $Q_1^+ := B_1^+ \times I_1$ is the unit upper-half cylinder and $\partial^0 Q_1^+ = Q_1 \cap \{y = 0\}$, where $B_1^+ := B_1 \cap \{y > 0\}$ ($B_1 \subset \mathbb{R}^{N+1}$ is the unit ball centered at 0) and $I_1 := (-1, 1)$, while the symbols ∇ and div denote the gradient and the divergence with respect to the spatial variable z , respectively.

The function $A: Q_1^+ \rightarrow \mathbb{R}^{N+1, N+1}$ is assumed to be symmetric and to satisfy the following ellipticity condition: There exist $0 < \lambda \leq \Lambda < +\infty$ such that

$$(1.2) \quad \lambda |\xi|^2 \leq A(z, t) \xi \cdot \xi \leq \Lambda |\xi|^2,$$

Mathematics Subject Classification 2020: 35B65 (primary); 58J35, 35B44, 35B53 (secondary).

Keywords: weighted parabolic equations, degenerate/singular weights, uniform regularity estimates.

for all $\xi \in \mathbb{R}^{N+1}$ and a.e. $(z, t) \in Q_1^+$, while $f: Q_1^+ \rightarrow \mathbb{R}$ and $F: Q_1^+ \rightarrow \mathbb{R}^{N+1}$ are given functions belonging to some suitable functional spaces. The notion of weak solution is given in Definition 2.2.

Our theory fits into the context of the regularity theory for linear non-uniformly parabolic equations; in particular, second order linear parabolic equations where the lack of uniform parabolicity is entailed by a weight term. Among all the papers on this topic, we quote the pioneering works [15, 25], where Harnack estimates and local Hölder continuity of solutions have been established when the weight ω either comes from quasiconformal mappings or belongs to the A_2 -Muckenhoupt class, that is,

$$\sup_B \left(\frac{1}{|B|} \int_B \omega \right) \left(\frac{1}{|B|} \int_B \omega^{-1} \right) \leq C,$$

where the supremum is taken over every ball $B \subset \mathbb{R}^{N+1}$.

The weight term $|y|^a$ we are considering here is A_2 -Muckenhoupt in the range $a \in (-1, 1)$. However, the peculiar geometry of the degeneracy/singularity set of our weight – the characteristic hyperplane Σ – allows us to get more information compared to the general theory quoted above and to deal with the full range $a > -1$.

In the spirit of the elliptic framework, see [46, 47, 49], one can build a complete Schauder theory in $C_p^{k,\alpha}$ spaces for weak solutions to (1.1): this is the main issue of the present paper, together with its first part [4]. Let us remark here that the regularity we obtain strongly relies on the *natural conormal boundary condition*

$$\lim_{y \rightarrow 0^+} y^a (A \nabla u + F) \cdot e_{N+1} = 0$$

we impose on the characteristic hyperplane Σ . The reader should keep in mind that the function y^{1-a} is a solution to the homogeneous equation $\operatorname{div}(y^a \nabla (y^{1-a})) = 0$ when $a < 1$ with homogeneous Dirichlet boundary condition on Σ but, if $a \in (0, 1)$, it is no more than $(1 - a)$ -Hölder continuous up to Σ .

We also mention [20, 29], where Schauder estimates are obtained when data are of Dini type (elliptic framework), and [23, 24], where the authors established some regularity estimates of Sobolev type for a wide class of parabolic equations including (1.1) (see also [36–39]).

Moreover, the study of weighted problems like (1.1) is strongly related to the theory of *edge operators* [34, 35] and *nonlocal operators*. The latter relies in the connection between a class of fractional heat operators like $(\partial_t - \Delta)^{(1-a)/2}$ – possibly with variable coefficients – and their extension theories [8, 40, 48], which represent the parabolic counterpart of [13]. Within this context, Schauder estimates for solutions to fractional parabolic equations involving $(\partial_t - \operatorname{div}_x(A(x)\nabla_x))^{(1-a)/2}$ have been established in [10]. With respect to our notation, this corresponds to regularity estimates in the (x, t) -variables on Σ and $a \in (-1, 1)$ (see also [11, 14, 22, 42]). Let us also mention that space analyticity (in the full z variable) and smoothness in (z, t) of solutions to equation (1.1) were already available by [9] when $a \in (-1, 1)$ and coefficients are analytic and satisfy suitable extra assumptions.

It is worth mentioning that the study of such operators is central in numerous papers of recent years; we quote [3, 12] (reaction-diffusion equations), [2, 6, 16] (obstacle problems), [5, 45] (nodal set analysis), [28] (nonlocal harmonic maps flow) and the references therein.

According to [49] (elliptic setting), the Schauder estimates for equations with degenerate weights have a remarkable application in the context of the *boundary Harnack principles*. Such boundary Harnack principles allow to “compare the regularity” of two solutions u, v of the same equation ($u > 0$) which vanish on the same portion of a fixed boundary. In particular, in rough domains such as Lipschitz, NTA or Hölder domains, the ratio $w = v/u$ is bounded up to the boundary where u and v vanish (in the first two cases w is even Hölder continuous). The literature on the topic is extensive; we refer to [18, 19] for a unified approach (equations in divergence and non-divergence form) and an interesting review of the topic. Then, when the boundary is $C^{k,\alpha}$, the *higher order boundary Harnack principle* improves the regularity of the quotient w up to $C^{k,\alpha}$, see [17] for the elliptic case and [7, 30] for its parabolic counterpart. We will see that our Schauder estimates for weighted equations provides an alternative proof of some of the results contained in the last two references.

Notably, the weighted elliptic Schauder theory developed in [46, 49] was used in the recent papers [1] and [41] to derive higher regularity of free interfaces for some semilinear free boundary problems (Alt–Phillips type). We wonder if the parabolic Schauder theory we develop here, together with [4], may help to address similar results for semilinear free boundary problems of parabolic type as well.

Main results

This paper is devoted to the higher order Schauder estimates for weak solutions to (1.1). The statement of our main result follows.

Theorem 1.1. *Let $N \geq 1$, $a > -1$, $r \in (0, 1)$, $\alpha \in (0, 1)$ and $k \in \mathbb{N}$. Let $A \in C_p^{k+1,\alpha}(Q_1^+)$ satisfy (1.2), let $f \in C_p^{k,\alpha}(Q_1^+)$ and $F \in C_p^{k+1,\alpha}(Q_1^+)$, and let u be a weak solution to (1.1). Then there exists $C > 0$, depending only on $N, a, \lambda, \Lambda, r, \alpha$ and $\|A\|_{C_p^{k+1,\alpha}(Q_1^+)}$, such that*

$$(1.3) \quad \|u\|_{C_p^{k+2,\alpha}(Q_r^+)} \leq C(\|u\|_{L^2(Q_1^+, y^a)} + \|f\|_{C_p^{k,\alpha}(Q_1)} + \|F\|_{C_p^{k+1,\alpha}(Q_1)}).$$

In our previous work [4], we established $C_p^{0,\alpha}$ and $C_p^{1,\alpha}$ estimates for solutions to (1.1), under suitable assumptions on coefficients and data, see Theorem 1.1 in [4]. These are obtained through a regularization of the weight and approximation, that is, by proving uniform-in- ε regularity estimates for solutions u_ε of the equation with the regularized weight $(\varepsilon^2 + y^2)^{a/2}$ and then passing to the limit as $\varepsilon \rightarrow 0^+$. The strategy to prove $C_p^{2,\alpha}$ (or higher order) estimates cannot rely on such ε -regularization scheme, since the ε -stability of the $C_p^{2,\alpha}$ estimate is false in general, even in the elliptic framework, see Remark 5.4 in [46].

Before sketching the main steps of the proof of Theorem 1.1, it is important to highlight the following facts, which substantially differ our strategy from the existing literature:

- In the *weighted elliptic framework* (see [46, 49]), as soon as the $C^{1,\alpha}$ regularity is available, one can iterate it on derivatives. This is obtained in two steps: first, one notices that, since the weighted elliptic operator commutes with all but one derivatives, $\partial_{x_i} u$ is also a solution for any $i = 1, \dots, N$ (and so $\partial_{x_i} u$ gains regularity); then, the

operator itself gives the regularity of the last derivative $\partial_y u$. Formally, this is because, in the special case $A = I$, one can re-write the equation as

$$-\partial_{yy}u - \partial_y F \cdot e_{N+1} - \frac{a}{y}(\partial_y u + F \cdot e_{N+1}) = f + \operatorname{div}_x F + \Delta_x u,$$

and thus if $\Delta_x u$ is smooth, then $\partial_y u$ is smooth by ODE methods (of course, provided that the data are smooth as well).

- In the *non-weighted parabolic framework* (see [32]), the idea is roughly the same. If $\Delta_x u$ is smooth, then the equation

$$\partial_t u = f + \operatorname{div} F + \Delta_x u$$

yields smoothness of $\partial_t u$.

- In the present *degenerate parabolic setting*, the “degenerate” variables are two, y and t , and the above strategies do not apply. In particular, the induction argument requires, as starting point, the $C_p^{2,\alpha}$ regularity of weak solutions (see Proposition 4.2).

Given the above remarks, our approach relies on a priori estimates and a regularization procedure by convolution with standard mollifiers. More precisely, for the $C_p^{2,\alpha}$ regularity:

- We establish some *a priori* $C_p^{2,\alpha}$ estimates in Proposition 4.2 using a blow-up argument combined with a Liouville theorem (see Theorem 1.2 below), in the spirit of [44] (see also [46] in the weighted elliptic setting).
- We prove $C_p^{2,\alpha}$ regularity of weak solutions when the data are C^∞ (see Lemma 4.3). In this step, the $C_p^{1,\alpha}$ regularity of weak solutions (see Theorem 2.4) is crucial.
- We use an approximation scheme to regularize (1.1) by convolution of the data with a family of standard mollifiers. Along the approximating sequence, the $C_p^{2,\alpha}$ regularity estimate extends to weak solutions with $f \in C_p^{0,\alpha}$ and $A, F \in C_p^{1,\alpha}$. In other words, we prove the *a posteriori* regularity estimate in Theorem 1.1 when $k = 0$.

For the $C^{k+2,\alpha}$ regularity, for every $k \geq 1$:

- When the forcing term is zero, i.e., when $f = 0$, we iterate the regularity estimates previously obtained – i.e., the $C_p^{1,\alpha}$ and $C_p^{2,\alpha}$ regularity – on partial derivatives of solutions, by using the same scheme as in the proof of Lemma 4.3 and Theorem 1.1 follows quite easily.
- In the case of general forcing terms $f \in C_p^{k,\alpha}$, the argument of (iv) does not apply (at least for $k = 1$), and hence we proceed as follows: we use the procedure described at points (i), (ii) and (iii) at any order k . To be more precise, the $C_p^{k+2,\alpha}$ *a priori* estimates are obtained inductively on k , starting from the $C_p^{2,\alpha}$ *a priori* estimates proved at point (i). This part crucially uses a delicate analysis of a second order weighted-type derivative of solutions in y (see Lemma 5.3). The $C_p^{k+2,\alpha}$ regularity when the data are smooth (the analogous of point (ii)) is also proved by induction in Lemma 5.2. Finally, with the same regularization argument in (iii), we finally obtain Theorem 1.1.

As anticipated above, the proof of the *a priori* $C_p^{2,\alpha}$ estimates strongly relies on the following Liouville theorem.

Theorem 1.2. *Let $a > -1$, $m \in \mathbb{N}$, $\gamma \in [0, m + 1)$, and let u be an entire solution to*

$$(1.4) \quad \begin{cases} y^a \partial_t u - \operatorname{div}(y^a \nabla u) = 0 & \text{in } \mathbb{R}_+^{N+1} \times \mathbb{R}, \\ \lim_{y \rightarrow 0^+} y^a \partial_y u = 0 & \text{on } \partial \mathbb{R}_+^{N+1} \times \mathbb{R}. \end{cases}$$

Assume that

$$(1.5) \quad |u(z, t)| \leq C(1 + (|z|^2 + |t|)^\gamma)^{1/2} \quad \text{for a.e. } (z, t) \in \mathbb{R}_+^{N+1} \times \mathbb{R}.$$

Then u is a polynomial with degree at most m in z and at most $\lfloor m/2 \rfloor$ in t .

As a consequence of our main theorem, we can treat more general equations with weights behaving as *distance functions* to a $C^{k+2,\alpha}$ ($k \in \mathbb{N}$) hypersurface $\Gamma \subset \mathbb{R}^{N+1}$ (curved characteristic manifolds) that we introduce below. The case of weights behaving as *distance functions* to a $C^{1,\alpha}$ hypersurface is treated in Corollary 1.3 of [4].

Such equations are set in cylindrical domains $\Omega^+ \times (-1, 1)$ of \mathbb{R}^{N+2} which “live” on one side of $\Gamma \times (-1, 1)$. Specifically, up to rotations and dilations, $0 \in \Gamma$ and there exist a spacial direction y and a function $\varphi \in C^{k+2,\alpha}(B_1 \cap \{y = 0\})$, with $\varphi(0) = 0$ and $\nabla_x \varphi(0) = 0$, such that

$$(1.6) \quad \Omega^+ \cap B_1 = \{y > \varphi(x)\} \cap B_1 \quad \text{and} \quad \Gamma \cap B_1 = \{y = \varphi(x)\} \cap B_1.$$

Then the family of weights $\delta = \delta(z)$ we consider behave as a distance function to Γ in the sense that $\delta \in C^{k+2,\alpha}(\Omega^+ \cap B_1)$, and

$$(1.7) \quad \begin{cases} \delta > 0 & \text{in } \Omega^+ \cap B_1, \\ |\nabla \delta| \geq c_0 > 0 & \text{in } \Omega^+ \cap B_1, \\ \delta = 0 & \text{on } \Gamma \cap B_1, \end{cases}$$

and we consider weighted equations of the form

$$(1.8) \quad \begin{cases} \delta^a \partial_t u - \operatorname{div}(\delta^a A \nabla u) = \delta^a f + \operatorname{div}(\delta^a F) & \text{in } (\Omega^+ \cap B_1) \times (-1, 1), \\ \delta^a (A \nabla u + F) \cdot \nu = 0 & \text{on } (\Gamma \cap B_1) \times (-1, 1), \end{cases}$$

where ν is the unit outward normal vector to Ω^+ on Γ . See Definition 7.2 in [4] for the definition of solutions to (1.8).

Corollary 1.3. *Let $a > -1$, $k \in \mathbb{N}$, $\alpha \in (0, 1)$ and let u be a weak solution to (1.8). Let $\varphi \in C^{k+2,\alpha}(B_1 \cap \{y = 0\})$ be the parametrization defined in (1.6) and $\delta \in C^{k+2,\alpha}(\Omega^+ \cap B_1)$ satisfying (1.7). Let also $A, F \in C_p^{k+1,\alpha}((\Omega^+ \cap B_1) \times (-1, 1))$, with A satisfying (1.2), and let $f \in C_p^{k,\alpha}((\Omega^+ \cap B_1) \times (-1, 1))$. Then there exists a constant $C > 0$, depending on $N, a, \lambda, \Lambda, \alpha, c_0, \|A\|_{C_p^{k+1,\alpha}((\Omega^+ \cap B_1) \times (-1, 1))}, \|\varphi\|_{C^{k+2,\alpha}(B_1 \cap \{y=0\})}$ and $\|\delta\|_{C^{k+2,\alpha}(\Omega^+ \cap B_1)}$, such that*

$$\begin{aligned} \|u\|_{C_p^{k+2,\alpha}((\Omega^+ \cap B_{1/2}) \times (-1/2, 1/2))} &\leq C \left(\|u\|_{L^2((\Omega^+ \cap B_1) \times (-1, 1), \delta^a)} \right. \\ &\quad + \|f\|_{C_p^{k,\alpha}((\Omega^+ \cap B_1) \times (-1, 1))} \\ &\quad \left. + \|F\|_{C_p^{k+1,\alpha}((\Omega^+ \cap B_1) \times (-1, 1))} \right). \end{aligned}$$

Finally, following the program of the elliptic setting (see [49]), we provide an alternative proof of some *parabolic higher order boundary Harnack principles* as in [7, 30]. Such kind of “regularity comparison principle” between two caloric functions u, v (or solutions to more general parabolic equations), vanishing on the same fixed boundary, can be viewed as the Schauder regularity of their quotient $w = v/u$, which in turn satisfies a parabolic equation with degenerate weight u^2 , see (7.1). After proper diffeomorphic transformations of the domain, the Schauder theory for the ratio w follows as a byproduct of our main Theorem 1.1.

The “regularity comparison principle” is localized at boundary points which lie on the *lateral parabolic boundary* of a space-time domain. In other words, let us consider u, v solutions of

$$(1.9) \quad \begin{cases} \partial_t u - \operatorname{div}(A \nabla u) = g + Vu + b \cdot \nabla u & \text{in } \Omega \cap Q_1, \\ \partial_t v - \operatorname{div}(A \nabla v) = f + Vv + b \cdot \nabla v & \text{in } \Omega \cap Q_1, \\ u(z, t) \geq c_0 d_p((z, t), \partial\Omega \cap Q_1) & \text{in } \Omega \cap Q_1, \\ u = v = 0 & \text{on } \partial\Omega \cap Q_1, \end{cases}$$

where A, V, b, g and f are suitable data (see Theorem 1.4 below). Here, up to rotations, dilations and translations, 0 belongs to the parabolic lateral boundary of Ω ; that is, there exists a parametrization φ such that

$$(1.10) \quad \Omega \cap Q_1 = \{y > \varphi(x, t)\} \quad \text{and} \quad \partial\Omega \cap Q_1 = \{y = \varphi(x, t)\},$$

with $\varphi(0) = 0$ and $\nabla_x \varphi(0) = 0$. Moreover, the parabolic distance to the boundary is defined as

$$d_p((z, t), \partial\Omega \cap Q_1) = \inf_{(\xi, \tau) \in \partial\Omega \cap Q_1} d_p((z, t), (\xi, \tau)),$$

and the parabolic distance between points is defined in (2.1).

We will present here the parabolic higher order boundary Harnack principle for equations in divergence form in $C_p^{k+2, \alpha}$ -domains, $k \in \mathbb{N}$. However, let us stress the fact that the regularity assumptions we make on boundaries, coefficients and right-hand sides, always allows to pass from non-divergence to divergence form equations and vice versa, interchangeably. So, we are considering the same conditions set in [7], which are slightly more general compared to [30], where the assumptions on the drift terms are suboptimal. Actually, our approach allows us to treat equations with nontrivial forcing terms g in the right-hand side of the equation of u .

Theorem 1.4. *Let $k \in \mathbb{N}$, $\alpha \in (0, 1)$, and let u and v be solutions to (1.9). Let also $\varphi \in C_p^{k+2, \alpha}(Q_1 \cap \{y = 0\})$ be the parametrization defined in (1.10). Finally, let $A, f, g \in C_p^{k+1, \alpha}(\Omega \cap Q_1)$, with A satisfying (1.2), and $V, b \in C_p^{k, \alpha}(\Omega \cap Q_1)$. Then there exists a constant $C > 0$, depending on $N, \lambda, \Lambda, c_0, \alpha, \|A\|_{C_p^{k+1, \alpha}(\Omega \cap Q_1)}, \|g\|_{C_p^{k+1, \alpha}(\Omega \cap Q_1)}, \|V\|_{C_p^{k, \alpha}(\Omega \cap Q_1)}, \|b\|_{C_p^{k, \alpha}(\Omega \cap Q_1)}, \|\varphi\|_{C_p^{k+2, \alpha}(Q_1 \cap \{y=0\})}$ and $\|u\|_{L^2(\Omega \cap Q_1)}$, such that*

$$\left\| \frac{v}{u} \right\|_{C_p^{k+2, \alpha}(\Omega \cap Q_{1/2})} \leq C(\|v\|_{L^2(\Omega \cap Q_1)} + \|f\|_{C_p^{k+1, \alpha}(\Omega \cap Q_1)}).$$

2. Preliminaries

In this section, we introduce some preliminary notions from [4] (parabolic Hölder spaces, weak solutions, and so on). Further, we prove some auxiliary/technical results we will repeatedly use throughout the paper. We begin with the definitions of the parabolic Hölder spaces, see Chapter 4 of [32] and Chapter 1 of [31].

2.1. Parabolic Hölder spaces

Let $\Omega \subset \mathbb{R}^{N+1} \times \mathbb{R}$ be an open subset and $u: \Omega \rightarrow \mathbb{R}$. The parabolic distance $d_p: \Omega \times \Omega \rightarrow \mathbb{R}$ is defined by

$$(2.1) \quad d_p((z, t), (\zeta, \tau)) := (|z - \zeta|^2 + |t - \tau|)^{1/2},$$

for all $(z, t), (\zeta, \tau) \in \Omega$, where $z, \zeta \in \mathbb{R}^{N+1}$, $t, \tau \in \mathbb{R}$. For $\alpha \in (0, 1]$, we define the seminorms

$$[u]_{C_p^{0,\alpha}(\Omega)} := \sup_{\substack{(z,t),(\zeta,\tau) \in \Omega \\ (z,t) \neq (\zeta,\tau)}} \frac{|u(z, t) - u(\zeta, \tau)|}{(|z - \zeta|^2 + |t - \tau|)^{\alpha/2}},$$

$$[u]_{C_t^{0,\alpha}(\Omega)} := \sup_{\substack{(z,t),(\zeta,\tau) \in \Omega \\ t \neq \tau}} \frac{|u(z, t) - u(z, \tau)|}{|t - \tau|^\alpha},$$

and the norm

$$\|u\|_{C_p^{0,\alpha}(\Omega)} := \|u\|_{L^\infty(\Omega)} + [u]_{C_p^{0,\alpha}(\Omega)}.$$

If $\beta \in \mathbb{N}^{N+1}$ is a multi-index and $k \geq 1$, we define the seminorms

$$[u]_{C_p^{k,\alpha}(\Omega)} := \sum_{|\beta|+2j=k} [\partial_x^\beta \partial_t^j u]_{C_p^{0,\alpha}(\Omega)} + [u]_{C_t^{k-1,(1+\alpha)/2}(\Omega)},$$

$$[u]_{C_t^{k,(1+\alpha)/2}(\Omega)} := \sum_{|\beta|+2j=k} [\partial_x^\beta \partial_t^j u]_{C_t^{0,(1+\alpha)/2}(\Omega)},$$

and the norm

$$\|u\|_{C_p^{k,\alpha}(\Omega)} = \sum_{|\beta|+2j \leq k} \sup_{\Omega} |\partial_x^\beta \partial_t^j u| + [u]_{C_p^{k,\alpha}(\Omega)}.$$

We set

$$C_p^{k,\alpha}(\Omega) := \{u: \Omega \rightarrow \mathbb{R} : \|u\|_{C_p^{k,\alpha}(\Omega)} < +\infty\}.$$

Finally, we recall some interpolation inequalities in parabolic Hölder spaces.

Lemma 2.1 (Proposition 4.2 in [32]). *Let $N \geq 1$, and $0 < \beta < \alpha \leq 1$. Then, for every $\varepsilon > 0$, there exists $C > 0$ depending on N and ε such that*

$$(2.2) \quad \begin{cases} \|u\|_{C_p^{0,\beta}(\Omega)} \leq C \|u\|_{L^\infty(\Omega)} + \varepsilon \|u\|_{C_p^{0,\alpha}(\Omega)}, \\ \|\nabla u\|_{L^\infty(\Omega)} \leq C \|u\|_{L^\infty(\Omega)} + \varepsilon [u]_{C_p^{1,\alpha}(\Omega)}, \\ \|D^2 u\|_{L^\infty(\Omega)} + \|\partial_t u\|_{L^\infty(\Omega)} \leq C \|u\|_{L^\infty(\Omega)} + \varepsilon [u]_{C_p^{2,\alpha}(\Omega)}. \end{cases}$$

2.2. Weak solutions, energy estimates and $C_p^{1,\alpha}$ regularity

Let $r > 0$. In what follows, $B_r \subset \mathbb{R}^{N+1}$ denotes the ball of radius r centered at the origin, $I_r := (-r^2, r^2) \subset \mathbb{R}$, $Q_r := B_r \times I_r \subset \mathbb{R}^{N+2}$ is the parabolic cylinder of radius r centered at the origin and $Q_r^+ := Q_r \cap \{y > 0\}$, while $\partial^0 Q_r^+ := Q_r \cap \{y = 0\}$ is the flat boundary of the half cylinder.

We first recall the definition of weak solutions to problem (1.1), see Definition 2.15 in [4]. The weighted energy spaces appearing below, $L^2(Q_r^+, y^a)$, $L^2(Q_r^+, y^a)^{N+1}$, $H^1(B_r^+, y^a)$, $L^2(I_r; H^1(B_r^+, y^a))$, $L^\infty(I_r; L^2(B_r^+, y^a))$, are defined in Section 2.1 of [4].

Definition 2.2. Let $a > -1$, $N \geq 1$, $r > 0$, $f \in L^2(Q_r^+, y^a)$, $F \in L^2(Q_r^+, y^a)^{N+1}$. We say that u is a weak solution to (1.1) if $u \in L^2(I_r; H^1(B_r^+, y^a)) \cap L^\infty(I_r; L^2(B_r^+, y^a))$ and satisfies

$$-\int_{Q_r^+} y^a u \partial_t \phi \, dz \, dt + \int_{Q_r^+} y^a A \nabla u \cdot \nabla \phi \, dz \, dt = \int_{Q_r^+} y^a (f \phi - F \cdot \nabla \phi) \, dz \, dt,$$

for every $\phi \in C_c^\infty(Q_r)$. We say that u is an entire solution to

$$\begin{cases} y^a \partial_t u - \operatorname{div}(y^a A \nabla u) = y^a f + \operatorname{div}(y^a F) & \text{in } \mathbb{R}_+^{N+1} \times \mathbb{R}, \\ \lim_{y \rightarrow 0^+} y^a (A \nabla u + F) \cdot e_{N+1} = 0 & \text{on } \partial \mathbb{R}_+^{N+1} \times \mathbb{R}, \end{cases}$$

if, for every $r > 0$, u is a weak solution to (1.1).

Weak solutions satisfy the following local energy inequality. We state the version we obtained in [4] in the spirit of [8].

Lemma 2.3 (Lemma 3.2 in [4]). *Let $N \geq 1$, $a \in \mathbb{R}$, and let the function A satisfy (1.2). Let $f \in L^2(Q_1^+, y^a)$, $F \in L^2(Q_1, y^a)^{N+1}$, and let u be a weak solution to (1.1). Then there exists $C > 0$ depending only on N , a , λ and Λ such that, for every $1/2 \leq r' < r < 1$,*

$$(2.3) \quad \begin{aligned} & \operatorname{ess\,sup}_{t \in (-r'^2, r'^2)} \int_{B_{r'}^+} y^a u^2 + \int_{Q_{r'}^+} y^a |\nabla u|^2 \\ & \leq C \left[\frac{1}{(r - r')^2} \int_{Q_r^+} y^a u^2 + \|f\|_{L^2(Q_1^+, y^a)}^2 + \|F\|_{L^2(Q_1^+, y^a)}^2 \right]. \end{aligned}$$

Finally, we state the main theorem in [4].

Theorem 2.4 (Theorem 1.1 in [4]). *Let $N \geq 1$, $a > -1$, $r \in (0, 1)$, $p > N + 3 + a^+$ and $\alpha \in (0, 1) \cap (0, 1 - (N + 3 + a^+)/p]$. Let $A \in C_p^{0,\alpha}(Q_1^+)$ satisfy (1.2), $f \in L^p(Q_1^+, y^a)$, $F \in C_p^{0,\alpha}(Q_1^+)$, and let u be a weak solution to (1.1). Then there exists $C > 0$, depending only on N , a , λ , Λ , r , p , α and $\|A\|_{C_p^{0,\alpha}(Q_1^+)}$, such that*

$$\|u\|_{C_p^{1,\alpha}(Q_r^+)} \leq C (\|u\|_{L^2(Q_1^+, y^a)} + \|f\|_{L^p(Q_1^+, y^a)} + \|F\|_{C_p^{0,\alpha}(Q_1)}).$$

Moreover, u satisfies the conormal boundary condition

$$(A \nabla u + F) \cdot e_{N+1} = 0 \quad \text{on } \partial^0 Q_r^+.$$

2.3. Technical results

In what follows, we prove some auxiliary results that we will use throughout the paper. We begin with a local L^2 bound for difference quotients of weak solutions with respect to the time variable.

Lemma 2.5. *Let $N \geq 1, a > -1$ and let A satisfying (1.2) be such that $\partial_t A \in L^\infty(Q_1^+)$. Let also $f \in L^2(Q_1^+, y^a)$ and $F \in L^2(Q_1^+, y^a)^{N+1}$ be such that $\partial_t f \in L^2(Q_1^+, y^a)$ and $\partial_t F \in L^2(Q_1^+, y^a)^{N+1}$, and let u be a weak solution to (1.1). Consider the difference quotient of u with respect to t :*

$$(2.4) \quad u^h(z, t) := \frac{u(z, t + h) - u(z, t)}{h}, \quad h > 0.$$

Then there exists $C > 0$, depending only on N, a, λ and Λ , such that, for every $r', r \in \mathbb{R}$ satisfying $1/2 \leq r' < r < 1$ and $h > 0$,

$$(2.5) \quad \int_{Q_r^+} y^a (u^h)^2 \leq C \left(\frac{1}{(r - r')^2} \int_{Q_r^+} y^a |\nabla u|^2 + \|f\|_{L^2(Q_1^+, y^a)}^2 + \|F\|_{L^2(Q_1^+, y^a)}^2 + \|\partial_t A\|_{L^\infty(Q_1^+)}^2 \int_{Q_r^+} y^a |\nabla u|^2 + \|\partial_t f\|_{L^2(Q_1^+, y^a)}^2 + \|\partial_t F\|_{L^2(Q_1^+, y^a)}^2 \right).$$

Proof. Fix r, r' such that $1/2 \leq r' < r < 1$. For $h > 0$, with $r < 1 - h$, let us consider the Steklov average of u :

$$u_h(z, t) = \frac{1}{h} \int_t^{t+h} u(z, s) dz,$$

which, by definition, satisfies $\partial_t u_h = u^h$ a.e. in Q_1 and the equation

$$(2.6) \quad \int_{Q_r^+} y^a (\partial_t u_h \phi + (A \nabla u)_h \cdot \nabla \phi) = \int_{Q_r^+} y^a (f_h \phi - F_h \cdot \nabla \phi) dz dt,$$

for all $\phi \in C_c^\infty(Q_1^+)$. Now, for simplicity of the exposition, we assume $f = 0, F = 0$, and we discuss how to treat the general case in a second step.

Let us take $\phi = \eta^2 u^h$ as test function in (2.6), where η is a smooth cut-off function which will define later. Using the Hölder and Young inequalities, the properties of Steklov averages and (1.2), we obtain

$$(2.7) \quad \begin{aligned} \int_{Q_1^+} y^a \eta^2 (u^h)^2 &= \int_{Q_1^+} y^a (\eta^2 (A \nabla u)_h \cdot \nabla u^h + 2\eta u^h (A \nabla u)_h \cdot \nabla \eta) \\ &\leq \left(\int_{Q_1^+} y^a \eta^2 |(A \nabla u)_h|^2 \right)^{1/2} \left(\int_{Q_1^+} y^a \eta^2 |\nabla u^h|^2 \right)^{1/2} \\ &\quad + 2 \left(\int_{Q_1^+} y^a \eta^2 (u^h)^2 \right)^{1/2} \left(\int_{Q_1^+} y^a |(A \nabla u)_h|^2 |\nabla \eta|^2 \right)^{1/2} \\ &\leq \frac{C}{\delta} \int_{Q_1^+} y^a |\nabla u|^2 + \delta \int_{Q_1^+} y^a \eta^2 |\nabla u^h|^2 \\ &\quad + \frac{1}{2} \int_{Q_1^+} y^a \eta^2 (u^h)^2 + C \int_{Q_1^+} y^a |\nabla \eta|^2 |\nabla u|^2, \end{aligned}$$

for any fixed $\delta > 0$ and $C > 0$ depending only on N, a, λ and Λ .

In the spirit of Lemma 3.3 in [21], we set

$$r_0 = r', \quad r_n = r' + \sum_{k=1}^n \frac{r - r'}{2^k} \quad \text{and} \quad s_n = \frac{r_n + r_{n+1}}{2}, \quad n \in \mathbb{N},$$

and notice that r_n and s_n are increasing sequences satisfying $r_n < s_n < r_{n+1}$, $r_n \rightarrow r$ and $s_n \rightarrow r$.

For a given $n \in \mathbb{N}$, taking a cut-off function $\eta_n \in C_c^\infty(Q_1^+)$ in (2.7) such that

$$\text{spt } \eta_n \subset Q_{s_n}^+, \quad \eta_n \equiv 1 \quad \text{in } Q_{r_n}^+, \quad 0 \leq \eta_n \leq 1 \quad \text{and} \quad |\nabla \eta_n| \leq C \frac{2^n}{r - r'},$$

we deduce

$$(2.8) \quad \frac{1}{2} \int_{Q_{r_n}^+} y^a (u^h)^2 \leq \delta \int_{Q_{s_n}^+} y^a |\nabla u^h|^2 + C \left(\frac{2^{2n}}{(r - r')^2} + \frac{1}{\delta} \right) \int_{Q_r^+} y^a |\nabla u|^2.$$

Now, noticing that u^h is a weak solution to

$$y^a \partial_t u^h - \text{div}(y^a A \nabla u^h) = \text{div}(y^a A^h \nabla u) \quad \text{in } Q_r^+,$$

we may apply the Caccioppoli inequality (2.3) to u^h to obtain

$$(2.9) \quad \int_{Q_{s_n}^+} y^a |\nabla u^h|^2 \leq \frac{C' 2^{2n}}{(r - r')^2} \int_{Q_{r_{n+1}}^+} y^a (u^h)^2 + C' \int_{Q_r^+} y^a |A^h \nabla u|^2,$$

for some $C' > 0$ independent of h, r, r' . Then, setting $\delta = \frac{1}{9} \frac{(r - r')^2}{C' 2^{2n}}$ in (2.8) and using (2.9), we have

$$\begin{aligned} \int_{Q_{r_n}^+} y^a (u^h)^2 &\leq \frac{1}{9} \int_{Q_{r_{n+1}}^+} y^a (u^h)^2 \\ &\quad + \frac{C 2^{2n}}{(r - r')^2} \int_{Q_r^+} y^a |\nabla u|^2 + \frac{C \|\partial_t A\|_{L^\infty(Q_1^+)}^2}{2^{2n}} \int_{Q_r^+} y^a |\nabla u|^2. \end{aligned}$$

Now, multiplying both sides by 3^{-2n} and summing over n , we see that

$$\begin{aligned} \sum_{n=0}^\infty 3^{-2n} \int_{Q_{r_n}^+} y^a (u^h)^2 &\leq \sum_{n=0}^\infty 3^{-2n-2} \int_{Q_{r_{n+1}}^+} y^a (u^h)^2 \\ &\quad + \frac{C}{(r - r')^2} \sum_{n=0}^\infty \left(\frac{2}{3}\right)^{2n} \int_{Q_r^+} y^a |\nabla u|^2 \\ &\quad + \sum_{n=0}^\infty \frac{C \|\partial_t A\|_{L^\infty(Q_1^+)}^2}{6^{2n}} \int_{Q_r^+} y^a |\nabla u|^2, \end{aligned}$$

which implies that

$$\int_{Q_r^+} y^a (u^h)^2 \leq \frac{C}{(r - r')^2} \int_{Q_r^+} y^a |\nabla u|^2 + C \|\partial_t A\|_{L^\infty(Q_1^+)}^2 \int_{Q_r^+} y^a |\nabla u|^2,$$

for some new $C > 0$, which is exactly (2.5) in the case $f = 0$ and $F = 0$.

For non-trivial f and F in the right-hand side, we have two additional terms: one in (2.7) and one in (2.9). Both of them can be estimated using the arguments above, namely,

$$\int_{Q_1^+} y^a (f_h \eta^2 u^h + F_h \cdot \nabla(\eta^2 u^h)) \leq C \|f\|_{L^2(Q_1^+, y^a)}^2 + C_\delta \|F\|_{L^2(Q_1^+, y^a)}^2 + \frac{1}{4} \int_{Q_1^+} y^a \eta^2 (u^h)^2 + \delta \int_{Q_1^+} y^a \eta^2 |\nabla u^h|^2,$$

for every $\delta > 0$, where we have implicitly used that

$$\int_{Q_1^+} y^a ((f^h)^2 + |F^h|^2) \leq \int_{Q_1^+} y^a ((\partial_t f)^2 + |\partial_t F|^2),$$

for every $h \in (0, 1)$. With such estimate at hand, the argument above can be slightly adapted to obtain (2.5) in the general case. ■

An immediate consequence of the above estimates is that, under suitable regularity assumptions on the data, derivatives (with respect to t and x) of weak solutions to (1.1) are still weak solutions (of a suitable problem of the class (1.1)).

Lemma 2.6. *Let $a > -1$, $N \geq 1$, $r \in (0, 1)$ and let A satisfying (1.2) be such that $\partial_t A \in L^\infty(Q_1^+)$. Let $f \in L^2(Q_1^+, y^a)$ and $F \in L^2(Q_1^+, y^a)$ be such that $\partial_t f, \partial_t F \in L^2(Q_1^+, y^a)$, and let u be a weak solution to (1.1). Then $v := \partial_t u$ is a weak solution to*

$$(2.10) \quad \begin{cases} y^a \partial_t v - \operatorname{div}(y^a A \nabla v) = y^a \partial_t f + \operatorname{div}(y^a (\partial_t A \nabla u + \partial_t F)) & \text{in } Q_r^+, \\ \lim_{y \rightarrow 0^+} y^a (A \nabla v + \partial_t A \nabla u + \partial_t F) \cdot e_{N+1} = 0 & \text{on } \partial^0 Q_r^+. \end{cases}$$

Proof. Let us fix $0 < r < r' < r'' < 1$ and $h > 0$ such that $r'' < 1 - h$. Let u^h be the difference quotient of u with respect to t defined in (2.4). By Lemma 2.5, $\|u^h\|_{L^2(Q_{r''}^+, y^a)}$ is bounded independently of $h > 0$. Further, since u^h is a weak solution to

$$(2.11) \quad y^a \partial_t u^h - \operatorname{div}(y^a A \nabla u^h) = y^a f^h + \operatorname{div}(y^a (F^h + A^h \nabla u)) \quad \text{in } Q_{r''}^+,$$

we may use Lemma 2.3 to deduce that $\|u^h\|_{L^\infty(I_{r'}, L^2(B_{r'}^+, y^a))}$ and $\|u^h\|_{L^2(I_{r'}, H^1(B_{r'}^+, y^a))}$ are bounded independently of $h > 0$ as well.

Now, let $\xi \in C_c^\infty(B_{r'})$ be a cut-off function such that $0 \leq \xi \leq 1$ and $\xi \equiv 1$ in B_r and set $v^h := \xi u^h \in L^2(I_{r'}, H_0^1(B_{r'}^+, y^a))$. Arguing as in Lemma 4.2 and Remark 2.16 of [4], we obtain that v^h is a weak solution to

$$y^a \partial_t v^h - \operatorname{div}(y^a A \nabla v^h) = y^a \tilde{f} + \operatorname{div}(y^a \tilde{F}) \quad \text{in } Q_{r'}^+,$$

where

$$\tilde{f} := f^h \xi - (F^h + A^h \nabla u) \cdot \nabla \xi - A \nabla u^h \cdot \nabla \xi \quad \text{and} \quad \tilde{F} := (F^h + A^h \nabla u) \xi - u^h A \nabla \xi,$$

satisfying also $\|\partial_t v^h\|_{L^2(I_{r'}, H^{-1}(B_{r'}^+, y^a))} \leq C$, for some $C > 0$ independent of $h > 0$. Consequently,

$$\|v^h\|_{L^2(I_{r'}, H_0^1(B_{r'}^+, y^a))} + \|\partial_t v^h\|_{L^2(I_{r'}, H^{-1}(B_{r'}^+, y^a))} \leq C,$$

for some $C > 0$ independent of $h > 0$. Consequently, the Aubin–Lion lemma (see, e.g., Corollary 8 in [43]) yields the existence of $v \in L^2(I_{r'}, H_0^1(B_{r'}^+, y^a))$ such that $v^h \rightarrow v$ in $L^2(Q_{r'}^+, y^a)$ and $\nabla v^h \rightharpoonup \nabla v$ in $L^2(Q_{r'}^+, y^a)$. Since $\xi \equiv 1$ in Q_r^+ , one has that $u^h \rightarrow \partial_t u$ in $L^2(Q_r^+, y^a)$ and $\nabla u^h \rightharpoonup \nabla(\partial_t u)$ in $L^2(Q_r^+, y^a)$. Furthermore, by the (H=W) property (see [50, 51]), one has $\partial_t u \in L^2(I_r, H^1(B_r^+, y^a))$ and $\partial_t u \in L^\infty(I_r, L^2(B_r^+, y^a))$ by Fatou’s lemma.

Finally, let us fix a test function $\phi \in C_c^\infty(Q_r)$ if $a \in (-1, 1)$ or $\phi \in C_c^\infty(Q_r^+)$ if $a \geq 1$. By the same argument of Lemma 4.2 in [4], we can take the limit as $h \rightarrow 0^+$ in the weak formulation of (2.11), to deduce

$$\begin{aligned} 0 &= \int_{Q_r^+} y^a (-u^h \phi_t + A \nabla u^h \cdot \nabla \phi - f^h \phi + (F^h + A^h \nabla u) \cdot \nabla \phi) \\ &\rightarrow \int_{Q_r^+} y^a (-\partial_t u \phi_t + A \nabla \partial_t u \cdot \nabla \phi - \partial_t f \phi + (\partial_t F + \partial_t A \nabla u) \cdot \nabla \phi) \end{aligned}$$

as $h \rightarrow 0^+$, that is, $\partial_t u$ is a weak solution to (2.10). ■

Analogously, we obtain the equations of the partial derivatives with respect to x .

Lemma 2.7. *Let $a > -1$, $N \geq 1$, $r \in (0, 1)$, $i \in \{1, \dots, N\}$ and let A satisfying (1.2) be such that $\partial_{x_i} A \in L^\infty(Q_1^+)$. Let $f \in L^2(Q_1^+, y^a)$ and $F \in L^2(Q_1^+, y^a)$ be such that $\partial_{x_i} f, \partial_{x_i} F \in L^2(Q_1^+, y^a)$, and let u be a weak solution to (1.1). Then $v_i := \partial_{x_i} u$ is a weak solution to*

$$(12) \quad \begin{cases} y^a \partial_t v_i - \operatorname{div}(y^a A \nabla v_i) = y^a \partial_{x_i} f + \operatorname{div}(y^a (\partial_{x_i} A \nabla u + \partial_{x_i} F)) & \text{in } Q_r^+, \\ \lim_{y \rightarrow 0^+} y^a (A \nabla v_i + \partial_{x_i} A \nabla u + \partial_{x_i} F) \cdot e_{N+1} = 0 & \text{on } \partial^0 Q_r^+. \end{cases}$$

The proof closely follows the above one and we skip it.

The following two auxiliary results are in the spirit of Lemma 2.3 in [49] and Theorem 7.5 in [46] (see also Lemma 2.4 and Remark 2.5 in [49]) in the elliptic setting, and turn out to be crucial in rest of the paper.

Lemma 2.8. *Let $k \in \mathbb{N}$ and $v \in C_p^{k+1, \alpha}(Q_1^+)$ be such that $v(x, 0, t) \equiv 0$. Then $v/y \in C_p^{k, \alpha}(Q_1^+)$ and $[v/y]_{C_p^{k, \alpha}(Q_1^+)} \leq [v]_{C_p^{k+1, \alpha}(Q_1^+)}$.*

The proof follows its elliptic counterpart and we skip it.

Lemma 2.9. *Let $a > -1$, $k \in \mathbb{N}$, $\alpha \in (0, 1)$ and let $g \in C_p^{k, \alpha}(Q_1^+)$. Then the function*

$$\varphi(x, y, t) = \frac{1}{y^{1+a}} \int_0^y s^a g(x, s, t) ds$$

belongs to $C_p^{k, \alpha}(Q_1^+)$ and $[\varphi]_{C_p^{k, \alpha}(Q_1^+)} \leq C [g]_{C_p^{k+1, \alpha}(Q_1^+)}$, for some $C > 0$ depending only on a . Moreover, the function

$$\psi(x, y, t) = \frac{1}{y^a} \int_0^y s^a g(x, s, t) ds$$

satisfies $\partial_y \psi \in C_p^{k, \alpha}(Q_1^+)$ and $[\partial_y \psi]_{C_p^{k, \alpha}(Q_1^+)} \leq C [g]_{C_p^{k+1, \alpha}(Q_1^+)}$, for some $C > 0$ depending only on a .

Proof. First, notice that the second statement follows immediately from the first since $\partial_y \psi = -a\varphi + g$.

We prove the first statement by induction. Let $k = 0$ and $g \in C_p^{0,\alpha}(Q_1^+)$. The parabolic Hölder continuity in x and t is trivially verified. Indeed, let $P_1 = (x_1, y, t_1)$ and $P_2 = (x_2, y, t_2)$. Then

$$|\varphi(P_2) - \varphi(P_1)| \leq \frac{1}{y^{1+\alpha}} \int_0^y s^\alpha |g(x_1, s, t_1) - g(x_2, s, t_2)| ds \leq \frac{[g]_{C_p^{0,\alpha}(Q_1^+)}}{1+\alpha} d_p(P_2, P_1)^\alpha.$$

For $\delta > 0$, let us consider

$$S_1 := \{(y_1, y_2) : 0 < y_1 < y_2 \leq 1 \text{ and } y_2 - y_1 \geq \delta y_2\},$$

$$S_2 := \{(y_1, y_2) : 0 < y_1 < y_2 \leq 1 \text{ and } y_2 - y_1 < \delta y_2\}.$$

Taking $y_1, y_2 \in S_1$, one has

$$\begin{aligned} |\varphi(x, y_2, t) - \varphi(x, y_1, t)| &= \left| \frac{1}{y_2^{a+1}} \int_0^{y_2} s^a g(x, s, t) ds - \frac{1}{y_1^{a+1}} \int_0^{y_1} s^a g(x, s, t) ds \right| \\ &= \left| \frac{1}{y_2^{a+1}} \int_0^{y_2} s^a (g(x, s, t) - g(x, 0, t)) ds \right. \\ &\quad \left. - \frac{1}{y_1^{a+1}} \int_0^{y_1} s^a (g(x, s, t) - g(x, 0, t)) ds \right| \\ &\leq \frac{[g]_{C_p^{0,\alpha}(Q_1^+)}}{y_2^{a+1}} \int_0^{y_2} s^{a+\alpha} ds + \frac{[g]_{C_p^{0,\alpha}(Q_1^+)}}{y_1^{a+1}} \int_0^{y_1} s^{a+\alpha} ds \\ &= \frac{[g]_{C_p^{0,\alpha}(Q_1^+)}}{a+\alpha+1} (y_2^\alpha + y_1^\alpha) \leq \frac{2[g]_{C_p^{0,\alpha}(Q_1^+)}}{a+\alpha+1} y_2^\alpha \\ &\leq \frac{2[g]_{C_p^{0,\alpha}(Q_1^+)}}{\delta^\alpha(a+\alpha+1)} (y_2 - y_1)^\alpha. \end{aligned}$$

Let now $y_1, y_2 \in S_2$. Then

$$\begin{aligned} |\varphi(x, y_2, t) - \varphi(x, y_1, t)| &= \left| \frac{1}{y_2^{a+1}} \int_0^{y_2} s^a (g(x, s, t) - g(x, 0, t)) ds \right. \\ &\quad \left. - \frac{1}{y_1^{a+1}} \int_0^{y_1} s^a (g(x, s, t) - g(x, 0, t)) ds \right| \\ &\leq \frac{1}{y_2^{a+1}} \int_{y_1}^{y_2} s^a |g(x, s, t) - g(x, 0, t)| ds \\ &\quad + \left(\frac{1}{y_1^{a+1}} - \frac{1}{y_2^{a+1}} \right) \int_0^{y_1} s^a |g(x, s, t) - g(x, 0, t)| ds \\ &\leq \frac{[g]_{C_p^{0,\alpha}(Q_1^+)}}{a+\alpha+1} \left(\frac{y_2^{a+\alpha+1} - y_1^{a+\alpha+1}}{y_2^{a+1}} + \left(\frac{1}{y_1^{a+1}} - \frac{1}{y_2^{a+1}} \right) y_1^{a+\alpha+1} \right) \end{aligned}$$

$$\begin{aligned}
 &= \frac{[g]C_p^{0,\alpha}(Q_1^+)}{a + \alpha + 1} \left(y_2^\alpha - y_1^\alpha \left(\frac{y_1}{y_2} \right)^{a+1} + y_1^\alpha \left(1 - \left(\frac{y_1}{y_2} \right)^{a+1} \right) \right) \\
 &= \frac{[g]C_p^{0,\alpha}(Q_1^+)}{a + \alpha + 1} \left(y_2^\alpha + y_1^\alpha - 2y_1^\alpha \left(\frac{y_1}{y_2} \right)^{a+1} \right) \\
 &= \frac{[g]C_p^{0,\alpha}(Q_1^+)}{a + \alpha + 1} \left(y_2^\alpha - y_1^\alpha + 2y_1^\alpha \left(1 - \left(\frac{y_1}{y_2} \right)^{a+1} \right) \right) \\
 &\leq \frac{[g]C_p^{0,\alpha}(Q_1^+)}{a + \alpha + 1} \left(y_2^\alpha - y_1^\alpha + C_a y_1^\alpha \left(1 - \frac{y_1}{y_2} \right) \right),
 \end{aligned}$$

where $C_a > 0$ is a constant which depends only on a . Consequently, since by definition $y_1/y_2 > 1 - \delta$, we have

$$\begin{aligned}
 \frac{|\varphi(x, y_2, t) - \varphi(x, y_1, t)|}{(y_2 - y_1)^\alpha} &\leq \frac{[g]C_p^{0,\alpha}(Q_1^+)}{a + \alpha + 1} \left(\frac{y_2^\alpha - y_1^\alpha}{(y_2 - y_1)^\alpha} + C_a \frac{y_1^\alpha (y_2 - y_1)}{y_2 (y_2 - y_1)^\alpha} \right) \\
 &\leq \frac{[g]C_p^{0,\alpha}(Q_1^+)}{a + \alpha + 1} (1 + C_a \delta^{1-\alpha}),
 \end{aligned}$$

and hence, the case $k = 0$ follows.

Next, let us assume that the our claim is true for some $k \in \mathbb{N}$ and let us prove it for $k + 1$. We assume $g \in C_p^{k+1,\alpha}(Q_1^+)$ and show that $\varphi \in C_p^{k+1,\alpha}(Q_1^+)$.

Since

$$\partial_{x_i} \varphi = \frac{1}{y^{1+a}} \int_0^y s^a \partial_{x_i} g(x, s, t) ds, \quad i = 1, \dots, N, \quad \partial_t \varphi = \frac{1}{y^{1+a}} \int_0^y s^a \partial_t g(x, s, t) ds,$$

we immediately have that φ is C^{k+1} in x and t . Moreover, the boundedness of the $C_t^{(1+\alpha)/2}$ -seminorm of the mixed-derivatives follows as the case $k = 0$.

We are left to prove that $\partial_y \varphi \in C_p^{k,\alpha}(Q_1^+)$. To do this, we can rewrite φ as

$$\varphi(x, y, t) = \frac{1}{y^{1+a}} \int_0^y s^a (g(x, s, t) - g(x, 0, t)) ds + \frac{g(x, 0, t)}{a + 1},$$

and observe that

$$\partial_y \varphi(x, y, t) = -\frac{a + 1}{y^{2+a}} \int_0^y s^{a+1} \frac{g(x, s, t) - g(x, 0, t)}{s} ds + \frac{g(x, y, t) - g(x, 0, t)}{y}.$$

By Lemma 2.8, one has that

$$\frac{g(x, y, t) - g(x, 0, t)}{y} \in C_p^{k,\alpha}(Q_1^+),$$

and our claim follows by the inductive assumption. ■

3. Liouville theorem

This section is devoted to the proof of the Liouville-type Theorem 1.2. We remark that, in the case $a \in (-1, 1)$, entire solutions to (1.4) satisfy the smoothness estimates in Theorem 1.1 of [9], hence, the proof of the Liouville theorem follows by a standard rescaling argument (for example, see Proposition 1.19 in [26]).

Proof of Theorem 1.2. Let us fix $R > 1$ and define

$$\tilde{\gamma} := a^+ + 2\gamma + N + 3.$$

Step 1. Choosing $r' = R$ and $r = 2R$ in (2.3) and using (1.5), we get

$$(3.1) \quad \int_{Q_R^+} y^a |\nabla u|^2 \leq \frac{C}{R^2} \int_{Q_{2R}^+} y^a u^2 \leq CR^{\tilde{\gamma}-2},$$

for some $C > 0$ depending only on N and a . On the other hand, choosing $r' = R$ and $r = 2R$ in (2.5) and combining (3.1) and (1.5), we obtain

$$(3.2) \quad \int_{Q_R^+} y^a (\partial_t u)^2 \leq \frac{C}{R^4} \int_{Q_{4R}^+} y^a u^2 \leq CR^{\tilde{\gamma}-4},$$

for some new $C > 0$.

Step 2. In this step, we prove that u is a polynomial in x . By Lemma 2.7, for every multi-index $\beta \in \mathbb{N}^N$, $\partial_x^\beta u$ is a weak solution to (1.4). Then, by iterating (3.1), one has

$$\int_{Q_R} y^a (\partial_x^\beta u)^2 \leq \int_{Q_R} y^a |\nabla u|^2 \leq CR^{\tilde{\gamma}-2|\beta|}.$$

Consequently, taking β such that $\tilde{\gamma} - 2|\beta| < 0$ and passing to the limit as $R \rightarrow +\infty$, it follows that $\partial_x^\beta u = 0$ and therefore u is a polynomial in the variable x , with degree less than or equal to m (the bound on the degree immediately follows by (1.5)).

Step 3. A slight modification of the above argument, which uses (3.2) instead of (3.1), shows that u is a polynomial in the variable t , with degree less or equal than $\lfloor m/2 \rfloor$.

Step 4. The last step is to prove that u is polynomial in y . By Remark 4.4 in [4], we notice that the even extension of u with respect to y is an entire solution to

$$(3.3) \quad |y|^a \partial_t u - \operatorname{div}(|y|^a \nabla u) = 0 \quad \text{in } \mathbb{R}^{N+1} \times \mathbb{R}.$$

Further, by Lemma 5.2 in [4], $v := |y|^a \partial_y u$ is an entire solution to

$$|y|^{-a} \partial_t u - \operatorname{div}(|y|^{-a} \nabla u) = 0 \quad \text{in } \mathbb{R}^{N+1} \times \mathbb{R},$$

while

$$(3.4) \quad w_1 := |y|^{-a} \partial_y v = \partial_{yy} u - a \frac{\partial_y u}{y},$$

is an entire solution to (3.3). Now, applying (2.3) twice, we deduce

$$\begin{aligned} \int_{Q_R} |y|^a w_1^2 &\leq \int_{Q_R} |y|^{-a} |\nabla v|^2 \leq \frac{C}{R^2} \int_{Q_{2R}} |y|^{-a} v^2 \\ &\leq \frac{C}{R^2} \int_{Q_{2R}} |y|^a |\nabla u|^2 \leq \frac{C}{R^4} \int_{Q_{4R}} |y|^a u^2 \leq CR^{\tilde{\gamma}-4}. \end{aligned}$$

Setting

$$(3.5) \quad w_{j+1} := \partial_{yy} w_j + a \frac{\partial_y w_j}{y},$$

and noticing that w_{j+1} is an entire solution to (3.3) for $j \in \mathbb{N}_+$, we may iterate the argument above to show the existence of $k \in \mathbb{N}$ such that $\tilde{\gamma} - 4k < 0$ and

$$\int_{Q_R} |y|^a w_k \leq CR^{\tilde{\gamma}-4k}.$$

Hence, taking the limit as $R \rightarrow +\infty$, we obtain $w_k = 0$, that is,

$$\partial_{yy} w_{k-1} + a \frac{\partial_y w_{k-1}}{y} = 0.$$

The above ODE can be explicitly solved:

$$w_{k-1} = c_{2k-1}(x, t) y |y|^{-a} + c_{2k-2}(x, t),$$

where $c_{2k-1}(x, t)$ and $c_{2k-2}(x, t)$ are polynomials. Now, iteratively solving the ODEs in (3.5) and (3.4), we obtain an explicit formula for u :

$$(3.6) \quad u = c_0(x, t) + \sum_{i \geq 1} y^{2i} c_{2i}(x, t) + \sum_{i \geq 1} y^{2i-1} |y|^{-a} c_{2i-1}(x, t),$$

where $c_i(x, t)$ are polynomial. All solutions to (3.3) satisfying a polynomial growth condition (without imposing any symmetry condition) have the form (3.6). Since u is an even solution (which comes from the conormal condition at the hyperplane), $c_{2i-1} \equiv 0$ for every $i \geq 1$. Therefore, our statement follows from the growth assumption (1.5). ■

In the following remark, we also provide a classification of the entire solutions to (1.4) satisfying the growth condition (1.5). Such classification was already obtained in Lemma 3.2 of [6] in the range $a \in (-1, 1)$ (see also Lemma 5.2 in [27] in the elliptic setting). We present the proof for completeness.

Remark 3.1. Let $a > -1$ and let $q_\kappa = q_\kappa(x, t)$ be a polynomial of parabolic degree κ in $\mathbb{R}^N \times \mathbb{R}$. Then there exists a unique polynomial $\tilde{q}_\kappa = \tilde{q}_\kappa(x, y, t)$ of parabolic degree κ in $\mathbb{R}^N \times \mathbb{R}_+ \times \mathbb{R}$ such that \tilde{q}_κ satisfies (1.4) and $\tilde{q}_\kappa(x, 0, t) = q_\kappa(x, t)$ for every $(x, t) \in \mathbb{R}^N \times \mathbb{R}$. Moreover,

$$(3.7) \quad \tilde{q}_\kappa(x, y, t) = q_\kappa(x, t) + \sum_{i=1}^{\lfloor \kappa/2 \rfloor} \frac{y^{2i}}{2i!} c_{2i} (\partial_t - \Delta_x)^i q_\kappa(x, t), \quad \text{where} \quad c_{2i} = \prod_{j=1}^i \frac{2j-1}{2j-1+a}.$$

Proof of (3.7). We denote with $\Delta_{(x,y)}$ the Laplacian in the variables (x, y) , Δ_x the Laplacian in the variable x and $(\partial_t - \Delta_x)^i$ the heat operator applied i times. Let $M := \lfloor \kappa/2 \rfloor$.

If such a polynomial \tilde{q}_κ exists and satisfies the Neumann boundary condition, that is, $y^a \partial_y \tilde{q}_\kappa \rightarrow 0$ as $y \rightarrow 0^+$, then

$$\tilde{q}_\kappa(x, y, t) = q_\kappa(x, t) + \sum_{i=1}^M y^{2i} q_i(x, t),$$

where $q_i(x, t)$ are polynomials such that $y^{2i}q_i(x, t)$ have parabolic degree at most κ . Indeed, according to Theorem 2.4, \tilde{q}_κ satisfies the stronger Neumann boundary condition $\partial_y \tilde{q}_\kappa = 0$ on $y = 0$. This implies that \tilde{q}_κ cannot contain a nontrivial term $yq_1(x, t)$. As in the proof of Theorem 1.2, we may iterate this argument to show that any term of the form $y^{2i+1}q_i(x, t)$ is identically zero.

Notice that

$$\begin{aligned}
 (3.8) \quad 0 &= (\partial_t - \Delta_{(x,y)})\tilde{q}_\kappa - \frac{a}{y} \partial_y \tilde{q}_\kappa \\
 &= (\partial_t - \Delta_x)q_\kappa + \sum_{i=1}^M y^{2i}(\partial_t - \Delta_x)q_i - \sum_{i=1}^M 2i(2i-1)y^{2i-2}q_i - a \sum_{i=1}^M 2iy^{2i-2}q_i \\
 &= (\partial_t - \Delta_x)q_\kappa - (2+2a)q_1 + \sum_{i=1}^{M-1} y^{2i}((\partial_t - \Delta_x)q_i - (2i+2)(2i+1+a)q_{i+1}) \\
 &\quad + y^{2M}(\partial_t - \Delta_x)q_M.
 \end{aligned}$$

Now, by iteratively solving the equation in (3.8), we obtain

$$(3.9) \quad \begin{cases} q_1 = \frac{(\partial_t - \Delta_x)q_\kappa}{2(1+a)}, \\ q_2 = \frac{(\partial_t - \Delta_x)q_1}{4(3+a)} = \frac{(\partial_t - \Delta_x)^2 q_\kappa}{4(3+a)2(1+a)}, \\ q_i = (\partial_t - \Delta_x)^i q_\kappa \prod_{j=1}^i \frac{1}{2j(2j-1+a)} = \frac{(\partial_t - \Delta_x)^i q_\kappa}{2i!} \prod_{j=1}^i \frac{2j-1}{2j-1+a}, \end{cases}$$

for $i \in \{1, \dots, M\}$. By construction, the function \tilde{q}_κ defined in (3.7) satisfies our statement. The uniqueness of \tilde{q}_κ immediately follows by the explicit formula (3.9) and the linearity of the differential operator. ■

4. $C_p^{2,\alpha}$ regularity

The goal of this section is to prove Theorem 1.1 when $k = 0$.

Theorem 4.1. *Let $N \geq 1, a > -1, r \in (0, 1), \alpha \in (0, 1)$. Let $A \in C_p^{1,\alpha}(Q_1^+)$ satisfy (1.2), let $f \in C_p^{0,\alpha}(Q_1^+)$ and $F \in C_p^{1,\alpha}(Q_1^+)$, and let u be a weak solution to (1.1). Then there exists $C > 0$, depending only on $N, a, \lambda, \Lambda, r, \alpha$ and $\|A\|_{C_p^{1,\alpha}(Q_1^+)}$, such that*

$$(4.1) \quad \|u\|_{C_p^{2,\alpha}(Q_r^+)} \leq C(\|u\|_{L^2(Q_1^+, y^\alpha)} + \|f\|_{C_p^{0,\alpha}(Q_1)} + \|F\|_{C_p^{1,\alpha}(Q_1)}).$$

The proof is based on some a priori estimates and an approximation argument we present below.

4.1. A priori $C_p^{2,\alpha}$ estimates

We begin by showing the a priori $C_p^{2,\alpha}$ estimates, stated in the following.

Proposition 4.2. *Let $N \geq 1$, $a > -1$, $\alpha \in (0, 1)$ and $r \in (0, 1)$. Let $A \in C_p^{1,\alpha}(Q_1^+)$ satisfy (1.2), let $f \in C_p^{0,\alpha}(Q_1^+)$, $F \in C_p^{1,\alpha}(Q_1^+)$ and let $u \in C_p^{2,\alpha}(Q_1^+)$ be a weak solution to (1.1). Then there exists $C > 0$, depending only on $N, a, \lambda, \Lambda, r, \alpha, \|A\|_{C_p^{1,\alpha}(Q_1^+)}$, such that (4.1) holds.*

Proof. The proof is divided in several steps as follows.

Step 1. Without loss of generality, we prove the statement for $r = 1/2$. To simplify the notation, let $x_{N+1} = y$, $\partial_i := \partial_{x_i}$ for $i = 1, \dots, N + 1$, and $\partial_{ij} := \partial_i \partial_j$ for $i, j = 1, \dots, N + 1$. In the following, we will refer to the variable y as either y or x_{N+1} depending on what seems more convenient. We begin with some preliminary observations.

By the regularity assumptions and Theorem 2.4, one has that u satisfies the equation pointwise in Q_1^+ , and so

$$(4.2) \quad \left(\partial_t u - \sum_{i,j=1}^{N+1} A_{i,j} \partial_{ij} u - \frac{a}{y} \sum_{j=1}^{N+1} A_{N+1,j} \partial_j u \right) = \left(g + \frac{a}{y} F_{N+1} \right) \quad \text{in } Q_1^+,$$

where

$$g := \sum_{i,j=1}^{N+1} \partial_i A_{i,j} \partial_j u + f + \sum_{i=1}^{N+1} \partial_i F \in C_p^{0,\alpha}(Q_1^+)$$

satisfies

$$(4.3) \quad \begin{aligned} \|g\|_{C_p^{0,\alpha}(Q_1^+)} &\leq 3\|A\|_{C_p^{1,\alpha}(Q_1^+)} \|u\|_{C_p^{1,\alpha}(Q_1^+)} + \|f\|_{C_p^{0,\alpha}(Q_1^+)} + \|F\|_{C_p^{1,\alpha}(Q_1^+)} \\ &\leq C(\|u\|_{L^\infty(Q_1^+)} + \|D^2 u\|_{L^\infty(Q_1^+)}) + \|f\|_{C_p^{0,\alpha}(Q_1^+)} + \|F\|_{C_p^{1,\alpha}(Q_1^+)}, \end{aligned}$$

for some $C > 0$ depending on $\|A\|_{C_p^{1,\alpha}(Q_1^+)}$, thanks to the interpolation inequality (2.2).

By the regularity assumptions on the data and u , and using the conormal boundary condition in (1.1) (which is satisfied pointwise by Theorem 2.4), we can take the limit as $y \rightarrow 0^+$ in (4.2) to get

$$(4.4) \quad \begin{aligned} \lim_{y \rightarrow 0^+} \frac{a \left(\sum_{j=1}^{N+1} A_{N+1,j} \partial_j u + F_{N+1} \right)(x, y, t)}{y} \\ = a \partial_y \left(\sum_{j=1}^{N+1} A_{N+1,j} \partial_j u + F_{N+1} \right)(x, 0, t) = \left(\partial_t u - \sum_{i,j} A_{i,j} \partial_{ij} u - g \right)(x, 0, t), \end{aligned}$$

for every $(x, 0, t) \in \partial^0 Q_1^+$.

It is enough to prove that for every $\delta > 0$ sufficiently small,

$$(4.5) \quad \begin{aligned} [u]_{C_p^{2,\alpha}(Q_{1/2}^+)} &\leq \delta [u]_{C_p^{2,\alpha}(Q_1^+)} + C_\delta (\|D^2 u\|_{L^\infty(Q_1^+)} + \|u\|_{L^\infty(Q_1^+)}) \\ &\quad + \|f\|_{C_p^{0,\alpha}(Q_1^+)} + \|F\|_{C_p^{1,\alpha}(Q_1^+)}, \end{aligned}$$

for some $C_\delta > 0$ depending only on $\delta, N, a, \lambda, \Lambda, \alpha, \|A\|_{C_p^{1,\alpha}(Q_1^+)}$. We will show later how (4.1) follows by (4.5).

Step 2. *Contradiction argument and blow-up sequences.*

By contradiction, we assume that there exist $\alpha \in (0, 1)$, $A^{(k)}, F^{(k)} \in C_p^{1,\alpha}(Q_1^+)$, $f_k \in C_p^{0,\alpha}(Q_1^+)$ with $\|A^{(k)}\|_{C_p^{1,\alpha}(Q_1^+)} \leq C$ and $u_k \in C_p^{2,\alpha}(Q_1^+)$ such that

$$\begin{cases} y^\alpha \partial_t u_k - \operatorname{div}(y^\alpha A^{(k)} \nabla u_k) = y^\alpha f_k + \operatorname{div}(y^\alpha F^{(k)}) & \text{in } Q_1^+, \\ \lim_{y \rightarrow 0^+} y^\alpha (A^{(k)} \nabla u_k + F^{(k)}) \cdot e_{N+1} = 0 & \text{on } \partial^0 Q_1^+, \end{cases}$$

and there exists a small $\delta_0 > 0$ such that

$$(4.6) \quad [u_k]_{C_p^{2,\alpha}(Q_{1/2}^+)} > \delta_0 [u_k]_{C_p^{2,\alpha}(Q_1^+)} + k (\|D^2 u_k\|_{L^\infty(Q_1^+)} + \|u_k\|_{L^\infty(Q_1^+)} + \|f_k\|_{C_p^{0,\alpha}(Q_1^+)} + \|F^{(k)}\|_{C_p^{1,\alpha}(Q_1^+)}).$$

Let us define

$$L_k := \max \left\{ \{ [\partial_{ij} u_k]_{C_p^{0,\alpha}(Q_{1/2}^+)} : i, j = 1, \dots, N + 1 \}, [\partial_t u_k]_{C_p^{0,\alpha}(Q_{1/2}^+)}, \{ [\partial_i u_k]_{C_t^{(1+\alpha)/2}(Q_{1/2}^+)} : i = 1, \dots, N + 1 \} \right\},$$

and distinguish two cases: first, we assume that there exist $i, j \in \{1, \dots, N + 1\}$ such that

$$(4.7) \quad L_k = [\partial_{ij} u_k]_{C_p^{0,\alpha}(Q_{1/2}^+)}.$$

Later we will deal with the second case, when $L_k = [\partial_t u_k]_{C_t^{(1+\alpha)/2}(Q_{1/2}^+)}$. The case $L_k = [\partial_i u_k]_{C_p^{0,\alpha}(Q_{1/2}^+)}$ is very similar to (4.7) and we skip it.

Now, we consider two sequences of points $P_k(z_k, t_k), \bar{P}_k(\xi_k, \tau_k) \in Q_{1/2}^+$ such that

$$\frac{|\partial_{ij} u_k(P_k) - \partial_{ij} u_k(\bar{P}_k)|}{d_p(P_k, \bar{P}_k)^\alpha} \geq \frac{L_k}{2},$$

and define $r_k := d_p(P_k, \bar{P}_k)$. Notice that it must be $r_k \rightarrow 0$ as $k \rightarrow +\infty$, since

$$\frac{L_k}{2} \leq \frac{|\partial_{ij} u_k(P_k) - \partial_{ij} u_k(\bar{P}_k)|}{d_p(P_k, \bar{P}_k)^\alpha} \leq 2 \frac{\|\partial_{ij} u_k\|_{L^\infty(Q_{1/2}^+)}}{r_k^\alpha} \leq 2 \frac{[u_k]_{C_p^{2,\alpha}(Q_{1/2}^+)}}{r_k^\alpha k} \leq 2 \frac{L_k}{r_k^\alpha k},$$

where we have used (4.6) and the definition of L_k .

Let $\hat{z}_k = (\hat{x}_k, \hat{y}_k) \in B_{1/2}^+$ to be specified below. For k large, let us define

$$Q(k) := \frac{B_1^+ - \hat{z}_k}{r_k} \times \frac{(-1 - t_k, 1 - t_k)}{r_k^2},$$

and set $Q^\infty := \lim_{k \rightarrow +\infty} Q(k)$, along an appropriate subsequence. For $(z, t) \in Q(k)$, consider the blow-up sequence

$$(4.8) \quad w_k(z, t) := \frac{u_k(r_k z + \hat{z}_k, r_k^2 t + t_k) - T_k(z, t)}{[u_k]_{C_p^{2,\alpha}(Q_1^+)} r_k^{2+\alpha}},$$

where T_k is the quadratic parabolic polynomial

$$T_k(z, t) = u_k(\hat{z}_k, t_k) + r_k \sum_{i=1}^{N+1} \partial_i u_k(\hat{z}_k, t_k) x_i + \frac{r_k^2}{2} \sum_{i,j=1}^{N+1} \partial_{ij} u_k(\hat{z}_k, t_k) x_i x_j + r_k^2 \partial_t u_k(\hat{z}_k, t_k) t.$$

Notice that w_k satisfies

$$(4.9) \quad w_k(0) = |\nabla w_k(0)| = |D^2 w_k(0)| = \partial_t w_k(0) = 0.$$

At this point we distinguish two cases:

- *Case 1:*

$$\frac{y_k}{r_k} = \frac{d_p(P_k, \Sigma)}{r_k} \rightarrow +\infty \quad \text{as } k \rightarrow \infty.$$

In this case, we set $\hat{z}_k = z_k$, and we have $Q^\infty = \mathbb{R}^{N+2}$.

- *Case 2:*

$$\frac{y_k}{r_k} = \frac{d_p(P_k, \Sigma)}{r_k} \leq C,$$

for some $C > 0$ independent of k . In this case, we set $\hat{z}_k = (x_k, 0)$, and we have $Q^\infty = \mathbb{R}_+^{N+1} \times \mathbb{R}$.

Step 3. Hölder estimates and convergence of the blow-up sequences.

Let us fix a compact set $K \subset Q^\infty$. Then $K \subset Q(k)$ for any k large enough. By definition of the $C_p^{0,\alpha}$ seminorm and the parabolic scaling, for every $P = (z, t)$, $Q = (\xi, \tau) \in K$ and $i, j \in \{1, \dots, N + 1\}$, we have

$$|\partial_{ij} w_k(P) - \partial_{ij} w_k(Q)| \leq \frac{|\partial_{ij} u_k(r_k z + \hat{z}_k, r_k^2 t + t_k) - \partial_{ij} u_k(r_k \xi + \hat{z}_k, r_k^2 \tau + t_k)|}{[u_k]_{C_p^{2,\alpha}(Q_1^+) } r_k^\alpha} \leq d_p(P, Q)^\alpha,$$

and thus

$$(4.10) \quad \sup_{\substack{P, Q \in K \\ P \neq Q}} \frac{|\partial_{ij} w_k(P) - \partial_{ij} w_k(Q)|}{d_p(P, Q)^\alpha} \leq 1.$$

In a similar way, it is not difficult to obtain

$$(4.11) \quad \sup_{\substack{P, Q \in K \\ P \neq Q}} \frac{|\partial_t w_k(P) - \partial_t w_k(Q)|}{d_p(P, Q)^\alpha} \leq 1.$$

Further, for every $(z, t), (z, \tau) \in K$ and $i \in \{1, \dots, N + 1\}$,

$$|\partial_i w_k(z, t) - \partial_i w_k(z, \tau)| \leq \frac{|\partial_i u_k(r_k z + \hat{z}_k, r_k^2 t + t_k) - \partial_i u_k(r_k z + \hat{z}_k, r_k^2 \tau + t_k)|}{[u_k]_{C_p^{2,\alpha}(Q_1^+) } r_k^{1+\alpha}} \leq |t - \tau|^{(1+\alpha)/2},$$

which implies

$$(4.12) \quad \sup_{\substack{(z,t),(z,\tau) \in K \\ t \neq \tau}} \frac{|\partial_i w_k(z,t) - \partial_i w_k(z,\tau)|}{|t - \tau|^{(1+\alpha)/2}} \leq 1.$$

Combining (4.10), (4.11) and (4.12), we deduce that $[w_k]_{C_p^{2,\alpha}(K)}$ is uniformly bounded in k , for every compact set $K \subset Q^\infty$ (notice that the estimates above are valid in both case 1 and case 2, by the definition of Q^∞). Consequently, in light of (4.9), $\|w_k\|_{C_p^{2,\alpha}(K)}$ is uniformly bounded as well, and so we may apply the Arzelà–Ascoli theorem to conclude that $w_k \rightarrow \bar{w}$ in $C_p^{2,\gamma}(K)$, for every $\gamma \in (0, \alpha)$. Finally, a standard diagonal argument, combined with (4.10), (4.11) and (4.12), shows that

$$w_k \rightarrow \bar{w} \quad \text{in } C_p^{2,\gamma}(K), \text{ for every } K \subset\subset Q^\infty,$$

up to passing to a suitable subsequence, and

$$(4.13) \quad [\bar{w}]_{C_p^{2,\alpha}(Q^\infty)} \leq C_N.$$

for some $C_N > 0$ which depends only on N , by the definition of the $C_p^{2,\alpha}$ seminorm.

Step 4. The next step is to prove that $\partial_{ij} \bar{w}$ is not constant, where i, j are the indexes fixed in (4.7). To do this, we consider two sequences of points in $Q(k)$, defined as

$$S_k = \left(\frac{\xi_k - \hat{z}_k}{r_k}, \frac{\tau_k - t_k}{r_k^2} \right) \quad \text{and} \quad \bar{S}_k := \left(\frac{z_k - \hat{z}_k}{r_k}, 0 \right), \quad k \in \mathbb{N}.$$

In case 1, one has $\hat{z}_k = z_k$, then $S_k \rightarrow S \in Q^\infty$, up to passing to a subsequence and $\bar{S}_k = 0$ for every k . Then, using the definition of L_k and (4.6), it follows that

$$|\partial_{ij} w_k(S_k) - \partial_{ij} w_k(\bar{S}_k)| = |\partial_{ij} u_k(\bar{P}_k) - \partial_{ij} u_k(P_k)| \geq \frac{L_k}{2[u_k]_{C_p^{2,\alpha}(Q_1^+)}} \geq C \delta_0,$$

for some $C > 0$ independent on k and thus, passing to the limit as $k \rightarrow +\infty$, we obtain $|\partial_{ij} \bar{w}(S) - \partial_{ij} \bar{w}(0)| \geq C \delta_0$, that is, $\partial_{ij} \bar{w}$ is not constant.

In case 2, we can argue in a similar way. We have $\hat{z}_k = (x_k, 0)$ and therefore, $\bar{S}_k = \frac{y_k}{r_k} e_{n+1}$. Recalling that y_k/r_k is uniformly bounded by definition, $S_k \rightarrow \bar{S}$, for some \bar{S} , up to passing to a subsequence. On the other hand, the sequence S_k can be written as

$$S_k = \left(\frac{\xi_k - z_k}{r_k}, \frac{\tau_k - t_k}{r_k^2} \right) + \frac{y_k}{r_k} e_{N+1}.$$

Therefore, $S_k \rightarrow S$ as $k \rightarrow +\infty$, for some $S \in Q^\infty$, up to passing to a subsequence and so, as above, we have $|\partial_{ij} \bar{w}(S) - \partial_{ij} \bar{w}(\bar{S})| \geq C \delta_0$, which shows our claim.

Step 5. The equation of the limit \bar{w} .

In this step, we derive the equation of \bar{w} : as in the steps above, we divide the proof in two additional steps (case 1 and case 2).

In case 1, we have $r_k/y_k \rightarrow 0$ as $k \rightarrow +\infty$ and $\hat{z}_k = z_k$. Further, if

$$\begin{aligned} \bar{A}^{(k)}(z,t) &:= A^{(k)}(r_k z + \hat{z}_k, r_k^2 t + t_k), \\ \bar{\rho}_k(y) &:= r_k y + y_k, \\ (\bar{z}, \bar{t}) &:= \lim_{k \rightarrow +\infty} (\hat{z}_k, t_k), \end{aligned}$$

then, by the regularity assumptions on $A^{(k)}$, one has $\bar{A}^{(k)} \rightarrow \bar{A}$ as $k \rightarrow +\infty$, where $\bar{A} := \lim_{k \rightarrow +\infty} A^{(k)}(\bar{z}, \bar{t})$ is a symmetric matrix with constant coefficients satisfying (1.2).

We claim that \bar{w} is an entire solution to

$$(4.14) \quad \partial_t \bar{w} - \operatorname{div}(\bar{A} \nabla \bar{w}) = 0 \quad \text{in } \mathbb{R}^{N+2}.$$

Let us fix a compact set $K \subset Q^\infty$. By (4.2) and using estimates in parabolic Hölder spaces, w_k satisfies

$$\begin{aligned} & \left| \partial_t w_k - \sum_{i,j=1}^{N+1} (\bar{A}_{i,j}^{(k)}) \partial_{ij} w_k \right| \\ &= \frac{1}{[u_k]_{C_p^{2,\alpha}(Q_1^+) } r_k^\alpha} \left| \partial_t u_k(r_k z + z_k, r_k^2 t + t_k) - \sum_{i,j=1}^{N+1} (A_{i,j}^{(k)}) \partial_{ij} u_k(r_k z + z_k, r_k^2 t + t_k) \right. \\ & \quad \left. - \partial_t u_k(z_k, t_k) + \sum_{i,j=1}^{N+1} A_{i,j}^{(k)}(r_k z + z_k, r_k^2 t + t_k) \partial_{ij} u_k(z_k, t_k) \right| \\ &= \frac{1}{[u_k]_{C_p^{2,\alpha}(Q_1^+) } r_k^\alpha} \left| g_k(r_k z + z_k, r_k^2 t + t_k) \right. \\ & \quad \left. + \frac{a(\sum_{j=1}^{N+1} A_{N+1,j}^{(k)}) \partial_j u_k + F_{N+1}^{(k)}(r_k z + z_k, r_k^2 t + t_k)}{r_k y + y_k} - \partial_t u_k(z_k, t_k) \right. \\ & \quad \left. + \sum_{i,j=1}^{N+1} A_{i,j}^{(k)} \partial_{ij} u_k(z_k, t_k) + \sum_{i,j=1}^{N+1} (A_{i,j}^{(k)}(r_k z + z_k, r_k^2 t + t_k) - A_{i,j}^{(k)}(z_k, t_k)) \partial_{ij} u_k(z_k, t_k) \right| \\ &\leq \frac{1}{[u_k]_{C_p^{2,\alpha}(Q_1^+) } r_k^\alpha} \left| g_k(r_k z + z_k, r_k^2 t + t_k) \right. \\ & \quad \left. + \frac{a(\sum_{j=1}^{N+1} A_{N+1,j}^{(k)}) \partial_j u_k + F_{N+1}^{(k)}(r_k z + z_k, r_k^2 t + t_k)}{r_k y + y_k} - g_k(z_k, t_k) \right. \\ & \quad \left. - \frac{a(\sum_{j=1}^{N+1} A_{N+1,j}^{(k)}) \partial_j u_k + F_{N+1}^{(k)}(z_k, t_k)}{y_k} \right| + C \frac{\|D^2 u_k\|_{L^\infty(Q_1^+)}}{[u_k]_{C_p^{2,\alpha}(Q_1^+)}} \\ &= \frac{|g_k(r_k z + z_k, r_k^2 t + t_k) - g_k(z_k, t_k)|}{[u_k]_{C_p^{2,\alpha}(Q_1^+) } r_k^\alpha} + \frac{C \|D^2 u_k\|_{L^\infty(Q_1^+)}}{[u_k]_{C_p^{2,\alpha}(Q_1^+)}} \\ & \quad + \frac{a}{[u_k]_{C_p^{2,\alpha}(Q_1^+) } r_k^\alpha} \left| \frac{H_k(r_k z + z_k, r_k^2 t + t_k)}{r_k y + y_k} - \frac{H_k(z_k, t_k)}{y_k} \right| = \text{I} + \text{II} + \text{III}, \end{aligned}$$

where $C > 0$ is a new constant independent of k (here and below the constant $C > 0$ depends on K ; we omit this dependence to simplify the exposition) and

$$H_k(z, t) = \sum_{j=1}^{N+1} (A_{N+1,j}^{(k)}) \partial_j u_k(z, t) + F_{N+1}^{(k)}(z, t),$$

which satisfies $H_k(x_k, 0, t_k) = 0, \nabla H_k(x_k, 0, t_k) = \partial_y H_k(x_k, 0, t_k)e_{N+1}, H_k(z, t)/y \in C^{0,\alpha}(Q_1^+)$ by Lemma 2.8 and $[H_k/y]_{C^{0,\alpha}(Q_1^+)} \leq C[\nabla H_k]_{C^{1,\alpha}(Q_1^+)} \leq C[u_k]_{C_p^{2,\alpha}(Q_1^+)}$, by the assumption (4.6).

Now, by (4.6) and (4.3), we can estimate I as follows:

$$I := \frac{|g_k(r_k z + z_k, r_k^2 t + t_k) - g_k(z_k, t_k)|}{[u_k]_{C_p^{2,\alpha}(Q_1^+)} r_k^\alpha} \leq \frac{[g_k]_{C_p^{0,\alpha}(Q_1^+)}}{[u_k]_{C_p^{2,\alpha}(Q_1^+)}} \leq \frac{1}{k} \rightarrow 0,$$

as $k \rightarrow +\infty$. The term II vanishes as well as $k \rightarrow +\infty$, by similar considerations. Finally, let us prove that III vanishes as $k \rightarrow +\infty$. First,

$$\begin{aligned} & \left| \frac{H_k(r_k z + z_k, r_k^2 t + t_k)}{r_k y + y_k} - \frac{H_k(z_k, t_k)}{y_k} \right| \\ &= \left| \frac{H_k(r_k z + z_k, r_k^2 t + t_k)}{r_k y + y_k} - \frac{H_k(z_k, t_k)}{r_k y + y_k} - \frac{r_k y}{y_k} \frac{H_k(z_k, t_k)}{r_k y + y_k} \right| \\ &\leq \left| \frac{H_k(r_k z + z_k, r_k^2 t + t_k)}{r_k y + y_k} - \frac{H_k(z_k, t_k)}{r_k y + y_k} - \frac{\nabla H_k(z_k, t_k) \cdot r_k z}{r_k y + y_k} \right| \\ &\quad + \left| \frac{\nabla H_k(z_k, t_k) \cdot r_k z}{r_k y + y_k} - \frac{r_k y}{y_k} \frac{H_k(z_k, t_k)}{r_k y + y_k} \right| = III_i + III_{ii}. \end{aligned}$$

By using the parabolic first order expansion of H_k , (4.6) and $r_k y + y_k \geq y_k/2$, one has that

$$(4.15) \quad |III_i| \leq C \frac{[H_k]_{C_p^{1,\alpha}(Q_1^+)} r_k^{1+\alpha}}{r_k y + y_k} \leq C [u_k]_{C_p^{2,\alpha}(Q_1^+)} r_k^{1+\alpha} y_k^{-1}.$$

Instead, we estimate the term III_{ii} in the following way:

$$\begin{aligned} (4.16) \quad |III_{ii}| &\leq \left| \frac{\nabla H_k(z_k, t_k) \cdot r_k z}{r_k y + y_k} - \frac{\nabla H_k(x_k, 0, t_k) \cdot r_k z}{r_k y + y_k} \right| \\ &\quad + \left| \frac{\nabla H_k(x_k, 0, t_k) \cdot r_k z}{r_k y + y_k} - \frac{r_k y}{y_k} \frac{H_k(z_k, t_k)}{r_k y + y_k} \right| \\ &\leq C [H_k]_{C_p^{1,\alpha}(Q_1^+)} r_k y_k^{\alpha-1} \leq C [u_k]_{C_p^{2,\alpha}(Q_1^+)} r_k y_k^{\alpha-1}, \end{aligned}$$

where, in order to estimate the second term in the previous inequality, we have used the properties of H_k stated above. Hence, combining (4.15) and (4.16), we have that

$$|III| \leq C \frac{r_k}{y_k} + C \left(\frac{r_k}{y_k} \right)^{1-\alpha} \rightarrow 0 \quad \text{as } k \rightarrow +\infty,$$

since in case 1, $r_k/y_k \rightarrow 0$,

$$\partial_t w_k - \sum_{i,j=1}^{N+1} (\bar{A}_{i,j}^{(k)} \partial_{ij} w_k) \rightarrow \partial_t \bar{w} - \sum_{i,j=1}^{N+1} \bar{A}_{i,j} \partial_{ij} \bar{w} \quad \text{locally uniformly in } \mathbb{R}^{N+2},$$

as $k \rightarrow +\infty$, and hence, passing to the limit as $k \rightarrow +\infty$ into the equation of w_k above, (4.14) follows.

In case 2, we have $\hat{z}_k = (x_k, 0)$ and $r_k/y_k \leq C$ for some $C > 0$ independent of k . We claim that \bar{w} is an entire solution to

$$(4.17) \quad \begin{cases} y^\alpha \partial_t \bar{w} - \operatorname{div}(y^\alpha \bar{A} \nabla \bar{w}) = 0 & \text{in } \mathbb{R}_+^{N+1} \times \mathbb{R}, \\ \lim_{y \rightarrow 0^+} y^\alpha \bar{A} \nabla \bar{w} \cdot e_{N+1} = 0 & \text{on } \partial \mathbb{R}_+^{N+1} \times \mathbb{R}. \end{cases}$$

Let us fix a compact set $K \subset Q^\infty$. By using (4.2), (4.4) and the fact that (\hat{z}_k, t_k) belongs to $\partial^0 Q_1^+$, w_k satisfies

$$\begin{aligned} \mathcal{L}w_k &:= \partial_t w_k - \sum_{i,j=1}^{N+1} (\bar{A}_{i,j}^{(k)}) \partial_{ij} w_k - \frac{a}{y} \sum_{j=1}^{N+1} (\bar{A}_{N+1,j}^{(k)}) \partial_j w_k \\ &= \frac{1}{[u_k]_{C_p^{2,\alpha}(Q_1^+) r_k^\alpha}} \left[\partial_t u_k(r_k z + \hat{z}_k, r_k^2 t + t_k) - \sum_{i,j=1}^{N+1} (A_{i,j}^{(k)}) \partial_{ij} u_k(r_k z + \hat{z}_k, r_k^2 t + t_k) \right. \\ &\quad - \frac{a}{r_k y} \sum_{j=1}^{N+1} (A_{N+1,j}^{(k)}) \partial_j u_k(r_k z + \hat{z}_k, r_k^2 t + t_k) - \partial_t u_k(\hat{z}_k, t_k) \\ &\quad - \sum_{i,j=1}^{N+1} A_{i,j}^{(k)}(r_k z + \hat{z}_k, r_k^2 t + t_k) \partial_{ij} u_k(\hat{z}_k, t_k) \\ &\quad + \frac{a}{r_k y} \sum_{j=1}^{N+1} A_{N+1,j}^{(k)}(r_k z + \hat{z}_k, r_k^2 t + t_k) \partial_j u_k(\hat{z}_k, t_k) \\ &\quad \left. + \frac{a}{r_k y} \sum_{i,j=1}^{N+1} A_{N+1,j}^{(k)}(r_k z + \hat{z}_k, r_k^2 t + t_k) \partial_{ij} u_k(\hat{z}_k, t_k) r_k x_i \right] \\ &= \left\{ \frac{g_k(r_k z + z_k, r_k^2 t + t_k) - g_k(\hat{z}_k, t_k)}{[u_k]_{C_p^{2,\alpha}(Q_1^+) r_k^\alpha}} \right. \\ &\quad \left. - \frac{\sum_{i,j=1}^{N+1} (A_{i,j}^{(k)}(r_k z + \hat{z}_k, r_k^2 t + t_k) - A_{i,j}^{(k)}(\hat{z}_k, t_k)) \partial_{ij} u_k(\hat{z}_k, t_k)}{[u_k]_{C_p^{2,\alpha}(Q_1^+) r_k^\alpha}} \right\} \\ &\quad + \frac{a}{[u_k]_{C_p^{2,\alpha}(Q_1^+) r_k^\alpha}} \left\{ \frac{F_{N+1}^{(k)}(r_k z + \hat{z}_k, r_k^2 t + t_k)}{r_k y} \right. \\ &\quad \left. - \partial_y \left(\sum_{j=1}^{N+1} A_{N+1,j} \partial_j u + F_{N+1} \right) (\hat{z}_k, t_k) \right. \\ &\quad + \frac{1}{r_k y} \sum_{j=1}^{N+1} A_{N+1,j}^{(k)}(r_k z + \hat{z}_k, r_k^2 t + t_k) \partial_j u_k(\hat{z}_k, t_k) \\ &\quad \left. + \frac{1}{r_k y} \sum_{i,j=1}^{N+1} A_{N+1,j}^{(k)}(r_k z + \hat{z}_k, r_k^2 t + t_k) \partial_{ij} u_k(\hat{z}_k, t_k) r_k x_i \right\} =: J + JJ. \end{aligned}$$

Similar to case 1, J vanishes as $k \rightarrow +\infty$ (see the proof for I and II above). We are left to treat JJ . By Lemma 2.7 and Theorem 2.4, we may differentiate u_k with respect to x_i

($i = 1, \dots, N$) and $\partial_i u$ satisfies the following conormal boundary condition:

$$(4.18) \quad \lim_{y \rightarrow 0^+} \partial_i \left(\sum_{j=1}^{N+1} A_{N+1,j}^{(k)} \partial_j u_k + F_{N+1}^{(k)} \right) = 0,$$

and thus, recalling the conormal boundary condition of u_k , we deduce

$$\begin{aligned} \text{JJJ} := & -\frac{1}{r_k y} \left[\sum_{j=1}^{N+1} (A_{N+1,j}^{(k)} \partial_j u_k)(\hat{z}_k, t_k) + F_{N+1}^{(k)}(\hat{z}_k, t_k) \right. \\ & \left. + \sum_{i=1}^N \partial_i \left(\sum_{j=1}^{N+1} A_{N+1,j}^{(k)} \partial_j u_k + F_{N+1}^{(k)} \right)(\hat{z}_k, t_k) r_k x_i \right] = 0. \end{aligned}$$

Adding JJ and JJJ, expanding $F^{(k)}$ and $A^{(k)}$ at order one and using the estimates in parabolic Hölder spaces, we obtain

$$\begin{aligned} & \left| \frac{F_{N+1}^{(k)}(r_k z + \hat{z}_k, r_k^2 t + t_k)}{r_k y} - \partial_y \left(\sum_{j=1}^{N+1} A_{N+1,j} \partial_j u + F_{N+1} \right)(\hat{z}_k, t_k) \right. \\ & \quad + \frac{1}{r_k y} \sum_{j=1}^{N+1} A_{N+1,j}^{(k)}(r_k z + \hat{z}_k, r_k^2 t + t_k) \partial_j u_k(\hat{z}_k, t_k) \\ & \quad \left. + \frac{1}{r_k y} \sum_{i,j=1}^{N+1} A_{N+1,j}^{(k)}(r_k z + \hat{z}_k, r_k^2 t + t_k) \partial_{ij} u_k(\hat{z}_k, t_k) r_k x_i \right| \\ & = \left| \frac{F_{N+1}^{(k)}(r_k z + \hat{z}_k, r_k^2 t + t_k) - F_{N+1}^{(k)}(\hat{z}_k, t_k) - \sum_{i=1}^{N+1} \partial_i (F_{N+1}^{(k)})(\hat{z}_k, t_k) r_k x_i}{r_k y} \right. \\ & \quad + \frac{1}{r_k y} \sum_{j=1}^{N+1} \partial_j u_k(\hat{z}_k, t_k) (A_{N+1,j}^{(k)}(r_k z + \hat{z}_k, r_k^2 t + t_k) - A_{N+1,j}^{(k)}(\hat{z}_k, t_k)) \\ & \quad - \sum_{i=1}^{N+1} \partial_i A_{N+1,j}^{(k)}(\hat{z}_k, t_k) r_k x_i \\ & \quad \left. + \frac{1}{r_k y} \sum_{i,j=1}^{N+1} (A_{N+1,j}^{(k)}(r_k z + \hat{z}_k, r_k^2 t + t_k) - A_{N+1,j}^{(k)}(\hat{z}_k, t_k)) \partial_{i,j} u_k(\hat{z}_k, t_k) r_k x_i \right| \\ & \leq C r_k^\alpha ([F^{(k)}]_{C_p^{1,\alpha}(Q_1^+)} + [A^{(k)}]_{C_p^{1,\alpha}(Q_1^+)}) \|\nabla u_k\|_{L^\infty(Q_1^+)} + [A^{(k)}]_{C_p^{0,1}(Q_1^+)} \|D^2 u_k\|_{L^\infty(Q_1^+)} \\ & \leq \frac{C r_k^\alpha [u_k]_{C_p^{2,\alpha}(Q_{1/2}^+)}}{k}. \end{aligned}$$

Consequently,

$$|\mathcal{L} w_k| = o(1) \quad \text{as } k \rightarrow +\infty.$$

As in case 1, by step 3, one has

$$\mathcal{L}w_k \rightarrow \partial_t \bar{w} - \sum_{i,j=1}^{N+1} \bar{A}_{i,j} \partial_{ij} \bar{w} - \frac{a}{y} \sum_{j=1}^{N+1} \bar{A}_{N+1,j} \partial_j \bar{w} \quad \text{locally uniformly in } \mathbb{R}_+^{N+1},$$

as $k \rightarrow +\infty$, and so \bar{w} satisfies the equation in (4.17) in the classical sense. It remains to prove that \bar{w} satisfies the conormal boundary condition in (4.17). Since u_k satisfies (4.18) and following the arguments above, we find

$$\begin{aligned} \left| \sum_{j=1}^{N+1} (\bar{A}_{N+1,j}^{(k)} \partial_j w_k) \right| &= \frac{1}{[u_k]_{C_p^{2,\alpha}(Q_1^+)} r_k^{1+\alpha}} \left| \sum_{j=1}^{N+1} (A_{N+1,j}^{(k)} \partial_j u_k)(r_k z + \hat{z}_k, r_k^2 t + t_k) \right. \\ &\quad - \sum_{j=1}^{N+1} A_{N+1,j}^{(k)}(r_k z + \hat{z}_k, r_k^2 t + t_k) \partial_j u_k(\hat{z}_k, t_k) \\ &\quad \left. - \sum_{j=1}^{N+1} A_{N+1,j}^{(k)}(r_k z + \hat{z}_k, r_k^2 t + t_k) \partial_{ij} u_k(\hat{z}_k, t_k) r_k x_i \right| \\ &= \frac{1}{[u_k]_{C_p^{2,\alpha}(Q_1^+)} r_k^{1+\alpha}} \left| -F_{N+1}^{(k)}(r_k z + \hat{z}_k, r_k^2 t + t_k) \right. \\ &\quad - \sum_{j=1}^{N+1} \partial_j u_k(\hat{z}_k, t_k) (A_{N+1,j}^{(k)}(r_k z + \hat{z}_k, r_k^2 t + t_k) - A_{N+1,j}^{(k)}(\hat{z}_k, t_k)) \\ &\quad - \sum_{i=1}^{N+1} \partial_i (A_{N+1,j}^{(k)}(\hat{z}_k, t_k) r_k x_i) - \sum_{j=1}^{N+1} (A_{N+1,j}^{(k)} \partial_j u_k)(\hat{z}_k, t_k) \\ &\quad - \sum_{i,j=1}^{N+1} (\partial_i A_{N+1,j}^{(k)} \partial_j u_k)(\hat{z}_k, t_k) r_k x_i - \sum_{i,j=1}^{N+1} (A_{N+1,j}^{(k)} \partial_{ij} u_k)(\hat{z}_k, t_k) r_k x_i \\ &\quad \left. - \sum_{i,j=1}^{N+1} (A_{N+1,j}^{(k)}(r_k z + \hat{z}_k, r_k^2 t + t_k) - A_{N+1,j}^{(k)}(\hat{z}_k, t_k)) \partial_{ij} u_k(\hat{z}_k, t_k) r_k x_i \right| \\ &\leq \frac{1}{[u_k]_{C_p^{2,\alpha}(Q_1^+)} r_k^{1+\alpha}} \left| -F_{N+1}^{(k)}(r_k z + \hat{z}_k, r_k^2 t + t_k) + F_{N+1}^{(k)}(\hat{z}_k, t_k) \right. \\ &\quad + \sum_{i=1}^{N+1} \partial_i F_{N+1}^{(k)}(\hat{z}_k, t_k) r_k x_i - \sum_{i=1}^{N+1} \partial_i F_{N+1}^{(k)}(\hat{z}_k, t_k) r_k x_i \\ &\quad \left. - \sum_{i,j=1}^{N+1} \partial_i (A_{N+1,j}^{(k)} \partial_j u_k)(\hat{z}_k, t_k) r_k x_i \right| + o(1) \\ &\leq \frac{|\partial_y (\sum_{j=1}^{N+1} A_{N+1,j}^{(k)} \partial_j u_k + F_{N+1}^{(k)})(\hat{z}_k, t_k) r_k y|}{[u_k]_{C_p^{2,\alpha}(Q_1^+)} r_k^{1+\alpha}} + o(1) = o(1), \end{aligned}$$

as $k \rightarrow +\infty$. Thus, passing to the limit as $y \rightarrow 0^+$, we obtain

$$\lim_{y \rightarrow 0^+} \left| \sum_{j=1}^{N+1} (\tilde{A}_{N+1,j}^{(k)} \partial_j w_k) \right| \leq o(1),$$

and thus, taking the limit as $k \rightarrow +\infty$, it follows that

$$\lim_{y \rightarrow 0^+} \tilde{A} \nabla \bar{w} \cdot e_{N+1} = 0.$$

Combining this with the fact that $\bar{w} \in C_p^{2,\alpha}$ by (4.13) and recalling that $a > -1$, we have

$$\begin{aligned} \lim_{y \rightarrow 0^+} y^a \tilde{A} \nabla \bar{w} \cdot e_{N+1} &= \lim_{y \rightarrow 0^+} y^{1+a} \lim_{y \rightarrow 0^+} \frac{\tilde{A} \nabla \bar{w} \cdot e_{N+1}}{y} \\ &= \partial_y (\tilde{A} \nabla \bar{w} \cdot e_{N+1})|_{y=0} \cdot \lim_{y \rightarrow 0^+} y^{1+a} = 0, \end{aligned}$$

and so the proof of (4.17) is completed.

Step 6. Liouville theorems.

Since $\bar{w} \in C_p^{2,\alpha}(Q^\infty)$, see (4.13), it satisfies the growth condition

$$|\bar{w}(z, t)| \leq C(1 + (|z|^2 + |t|)^{2+\alpha})^{1/2}.$$

Moreover, \bar{w} has at least one non-constant second derivative and is an entire solution to (4.14) or (4.17). Then in case 1, we can invoke the Liouville theorem for the heat equation (see Remark 5.3 in [4]), and in case 2, we can invoke the Liouville Theorem 1.2 to reach the desired contradiction.

Step 7. We complete the analysis, considering the case when

$$L_k = [\partial_i u]_{C_i^{0,(1+\alpha)/2}(Q_{1/2}^+)},$$

for some $i \in \{1, \dots, N + 1\}$. We give a short sketch, pointing out the main differences in comparison to the preceding discussion.

We take two sequences of points $P_k = (z_k, t_k)$, $\bar{P}_k = (z_k, \tau_k) \in Q_{1/2}^+$ such that

$$\frac{|\partial_i u_k(z_k, t_k) - \partial_i u_k(z_k, \tau_k)|}{|t_k - \tau_k|^{(1+\alpha)/2}} \geq \frac{L_k}{2},$$

and set

$$r_k := d_p(P_k, \bar{P}_k) = |t_k - \tau_k|^{1/2}.$$

We define the blow-up sequence w_k as in (4.8), centered in P_k .

The steps 3, 5, 6 are the same as above. The only crucial difference is in step 4: In this case, one has that $\partial_i \bar{w}$ is non-constant in t . Indeed,

$$\left| \partial_i w_k \left(0, \frac{t_k - \tau_k}{r_k^2} \right) - \partial_i w_k(0, 0) \right| \geq \frac{L_k}{2 [u_k]_{C_p^{2,\alpha}(Q_1^+)}} \geq C_N \delta_0.$$

Taking the limit as $k \rightarrow +\infty$, we obtain that $|\partial_i \bar{w}(0, \bar{t}) - \partial_i \bar{w}(0, 0)| \geq C_N \delta_0$, where $\bar{t} = \lim_{k \rightarrow +\infty} (t_k - \tau_k) / r_k^2$. This allows us to conclude the proof of (4.5) by applying Theorem 1.2.

Step 8. Conclusion.

Finally, we briefly explain why (4.5) implies (4.1) (see Theorem 2.20 and Lemma 2.27 in [26] in the elliptic setting). First, by using a covering argument and the interpolation inequalities in (2.2), we have that (4.5) is satisfied in every $Q_\rho^+(P_0) \subset Q_1^+$, that is,

$$(4.19) \quad [u]_{C_p^{2,\alpha}(Q_{\rho/2}^+(P_0))} \leq \delta [u]_{C_p^{2,\alpha}(Q_\rho^+(P_0))} + C_\delta (\|u\|_{L^\infty(Q_1^+)} + \|f\|_{C_p^{0,\alpha}(Q_1^+)} + \|F\|_{C_p^{1,\alpha}(Q_1^+)}).$$

Now, let us define the seminorm

$$(4.20) \quad [u]_{2,\alpha,Q_1^+}^* := \sup_{Q_\rho^+(P_0) \subset Q_1^+} \rho^{2+\alpha} [u]_{C_p^{2,\alpha}(Q_{\rho/2}^+(P_0))}.$$

By using the sub-additivity of the Hölder seminorms with respect to unions of convex sets, one can prove that

$$(4.21) \quad [u]_{2,\alpha,Q_1^+}^* \leq C \sup_{Q_\rho^+(P_0) \subset Q_1^+} \rho^{2+\alpha} [u]_{C_p^{2,\alpha}(Q_{\rho/4}^+(P_0))},$$

for some constant $C > 0$ depending only on N and α . Then, by (4.19) and (4.20), we obtain

$$\rho^{2+\alpha} [u]_{C_p^{2,\alpha}(Q_{\rho/4}^+(P_0))} \leq \delta [u]_{2,\alpha,Q_1^+}^* + C_\delta (\|u\|_{L^\infty(Q_1^+)} + \|f\|_{C_p^{0,\alpha}(Q_1^+)} + \|F\|_{C_p^{1,\alpha}(Q_1^+)}).$$

Taking the supremum over $Q_\rho^+(P_0) \subset Q_1^+$ and recalling (4.21) yields

$$\frac{1}{C} [u]_{2,\alpha,Q_1^+}^* \leq \delta [u]_{2,\alpha,Q_1^+}^* + C_\delta (\|u\|_{L^\infty(Q_1^+)} + \|f\|_{C_p^{0,\alpha}(Q_1^+)} + \|F\|_{C_p^{1,\alpha}(Q_1^+)}).$$

Hence, our statement follows by taking $\delta > 0$ small enough and using the interpolation inequality (2.2). ■

4.2. A regularization scheme

In this second step, we proceed with a regularization argument. This allows to apply the a priori estimates above and prove Theorem 4.1.

Lemma 4.3. *Let $N \geq 1$, $a > -1$, $r \in (0, 1)$, $\alpha \in (0, 1)$. Let $A \in C^\infty(Q_1^+)$ satisfy (1.2), let $f, F \in C^\infty(Q_1^+)$, and let u be a weak solution to (1.1). Then $u \in C_p^{2,\alpha}(Q_r^+)$.*

Proof. We fix $0 < r < r' < 1$. For every $i = 1, \dots, N$, by the regularity assumption on A , f and F , and Lemma 2.7, we have that $\partial_{x_i} u$ solves (2.12) in Q_r^+ and, by Theorem 2.4, we deduce that $\partial_{x_i} u \in C_p^{1,\alpha}(Q_r^+)$. Analogously, by Lemma 2.6, $\partial_t u$ solves (2.10) in Q_r^+ and, by Theorem 2.4, we deduce that $\partial_t u \in C_p^{1,\alpha}(Q_r^+)$. To conclude, we need to prove that $\partial_y u \in C_p^{1,\alpha}(Q_r^+)$.

Using the regularity of ∇u and $\partial_t u$ obtained above, we may rewrite the equation of u as

$$(4.22) \quad \partial_y (y^a (A \nabla u + F)) \cdot e_{N+1} = y^a \left[\partial_t u - f - \sum_{i=1}^N \partial_{x_i} ((A \nabla u + F) \cdot e_i) \right] =: y^a g,$$

in the weak sense, where $g \in C_p^{0,\alpha}(Q_r^+)$. Then, integrating in y and using that

$$\lim_{y \rightarrow 0^+} (A \nabla u + F) \cdot e_{N+1} = 0,$$

see Theorem 2.4, one has

$$\psi(x, y, t) := (A \nabla u + F) \cdot e_{N+1}(x, y, t) = \frac{1}{y^\alpha} \int_0^y s^\alpha g(x, s, t) ds.$$

Since $\partial_{x_i} u, \partial_t u \in C_p^{1,\alpha}(Q_r^+)$, we have $\partial_{x_i} \psi \in C_p^{0,\alpha}(Q_r^+)$ by definition, for every $i = 1, \dots, N$, and $\partial_t \psi \in C_p^{0,\alpha}(Q_r^+)$. Consequently, $\psi \in C_t^{0,(1+\alpha)/2}(Q_r^+)$. Now, since $g \in C_p^{0,\alpha}(Q_r^+)$, Lemma 2.9 yields $\partial_y \psi \in C_p^{0,\alpha}(Q_r^+)$ and thus $\psi \in C_p^{1,\alpha}(Q_r^+)$. Noticing that, by (1.2), we have

$$(4.23) \quad \partial_y u = \frac{\psi - \sum_{j=1}^N A_{N+1,j} \partial_j u - F_{N+1}}{A_{N+1,N+1}},$$

and it follows that $\partial_y u \in C_p^{1,\alpha}(Q_r^+)$ and thus $u \in C_p^{2,\alpha}(Q_r^+)$. \blacksquare

We are now ready to show Theorem 4.1.

Proof of Theorem 4.1. Let us fix $0 < r < R < 1$ and let u be a weak solution to (1.1). Let us consider a smooth cut-off function $\xi \in C_c^\infty(B_R)$ such that $0 \leq \xi \leq 1$ and $\xi = 1$ in B_r . Then $v := \xi u$ is a weak solution to

$$\begin{cases} y^\alpha \partial_t v - \operatorname{div}(y^\alpha A \nabla v) = y^\alpha g + \operatorname{div}(y^\alpha G) & \text{in } Q_R^+, \\ \lim_{y \rightarrow 0^+} y^\alpha (A \nabla v + G) \cdot e_{N+1} = 0 & \text{on } \partial^0 Q_R^+, \\ v = 0 & \text{on } \partial B_R^+ \times I_R, \\ v = \eta u & \text{on } B_R^+ \times \{-R^2\}, \end{cases}$$

where

$$g := \xi f - F \cdot \nabla \xi - A \nabla u \cdot \nabla \xi \quad \text{and} \quad G := \xi F - u A \nabla \xi.$$

Let us denote with \bar{A} , \bar{f} and \bar{F} the even extensions of A , f and F with respect to y , respectively, and let

$$A_\varepsilon := \bar{A} * \rho_\varepsilon, \quad f_\varepsilon := \bar{f} * \rho_\varepsilon \quad \text{and} \quad F_\varepsilon := \bar{F} * \rho_\varepsilon,$$

where $\{\rho_\varepsilon\}_{\varepsilon>0}$ is a family of smooth mollifiers. Then, up to choosing ε small enough, $A_\varepsilon, f_\varepsilon, F_\varepsilon \in C_c^\infty(Q_R^+)$, and A_ε satisfies (1.2). For every $\varepsilon \in (0, 1)$, let v_ε be the weak solution to

$$\begin{cases} y^\alpha \partial_t v_\varepsilon - \operatorname{div}(y^\alpha A_\varepsilon \nabla v_\varepsilon) = y^\alpha g_\varepsilon + \operatorname{div}(y^\alpha G_\varepsilon) & \text{in } Q_R^+, \\ \lim_{y \rightarrow 0^+} y^\alpha (A_\varepsilon \nabla v_\varepsilon + G_\varepsilon) \cdot e_{N+1} = 0 & \text{on } \partial^0 Q_R^+, \\ v_\varepsilon = 0 & \text{on } \partial B_R^+ \times I_R, \\ v_\varepsilon = v & \text{on } B_R^+ \times \{-R^2\}, \end{cases}$$

where

$$g_\varepsilon := \xi f_\varepsilon - F_\varepsilon \cdot \nabla \xi - A_\varepsilon \nabla u \cdot \nabla \xi, \quad G_\varepsilon := \xi F_\varepsilon - u A_\varepsilon \nabla \xi.$$

By the same compactness argument of Lemma 2.6 (or, equivalently, Lemma 4.3 and Remark 4.4 in [4]), and by the classical theory of the Cauchy–Dirichlet problem in abstract Hilbert spaces, see [33], we have that $v_\varepsilon \rightarrow v$ in $L^2(Q_R^+, y^a)$, which implies that $v_\varepsilon \rightarrow u$ in $L^2(Q_r^+, y^a)$ by the definition of v . On the other hand, since $\xi \equiv 1$ in B_r , one has that v_ε is a weak solution to

$$\begin{cases} y^a \partial_t v_\varepsilon - \operatorname{div}(y^a D_\varepsilon \nabla v_\varepsilon) = y^a f_\varepsilon + \operatorname{div}(y^a F_\varepsilon) & \text{in } Q_r^+, \\ \lim_{y \rightarrow 0^+} y^a (D_\varepsilon \nabla v_\varepsilon + F_\varepsilon) \cdot e_{N+1} = 0 & \text{on } \partial^0 Q_r^+. \end{cases}$$

So, up to rescaling, Lemma 4.3 yields that $v_\varepsilon \in C_p^{2,\alpha}(Q_1^+)$. On the other hand, by Proposition 4.2, we deduce that v_ε satisfies the desired estimate (4.1) in Q_r^+ , uniformly in $\varepsilon > 0$. By the Arzelà–Ascoli theorem, we may thus take the limit as $\varepsilon \rightarrow 0^+$ and complete the proof of (4.1). ■

5. $C_p^{k+2,\alpha}$ regularity

In this section, we prove Theorem 1.1 for any $k \geq 1$ by combining some a priori estimates and an approximation argument. As anticipated in the introduction, we first deal with the case of a zero forcing term in the equation (1.1), i.e., $f = 0$. In this case, the main result follows by a simple iteration of the $C_p^{1,\alpha}$ and $C_p^{2,\alpha}$ estimates on partial derivatives. Secondly, we treat forcing terms $f \in C_p^{k,\alpha}$. In this case, the strategy is more involved and requires some additional and delicate steps (see Lemma 5.3).

5.1. Higher order Schauder estimates when $f = 0$

We begin by treating the simpler case $f = 0$.

Proof of Theorem 1.1 when $f = 0$. We proceed by induction. The initial step $k = 0$ follows by Theorem 4.1.

Let us fix $0 < r < r' < 1$ and assume that $A, F \in C_p^{j+2,\alpha}(Q_r^+)$ imply that (1.3) holds for $j = 0, \dots, k$ and prove it for $k + 1$. By Lemma 2.7 and the induction step we may differentiate the equation of u with respect to x_i to obtain $\partial_{x_i} u \in C_p^{k+2,\alpha}(Q_r^+)$ for every $i = 1, \dots, N$ and

$$\begin{aligned} (5.1) \quad \|\partial_{x_i} u\|_{C_p^{k+2,\alpha}(Q_r^+)} &\leq C(\|\partial_{x_i} u\|_{L^2(Q_{r'}^+, y^a)} + \|\partial_i F\|_{C_p^{k+1,\alpha}(Q_1^+)}) \\ &\leq C(\|u\|_{L^2(Q_1^+, y^a)} + \|F\|_{C_p^{k+2,\alpha}(Q_1^+)}) \end{aligned}$$

for some $C > 0$ which depends on $N, a, \lambda, \Lambda, r, \alpha$ and $\|A\|_{C_p^{k+2,\alpha}(Q_1^+)}$. On the other hand, By Lemma 2.6 and the induction step (noticing that in the case $k = 0$, we use Theorem 2.4), we may differentiate the equation of u with respect to t to obtain $\partial_t u \in C_p^{k+1,\alpha}(Q_r^+)$ and

$$\begin{aligned} (5.2) \quad \|\partial_t u\|_{C_p^{k+1,\alpha}(Q_r^+)} &\leq C(\|\partial_t u\|_{L^2(Q_{r'}^+, y^a)} + \|\partial_t F\|_{C_p^{k,\alpha}(Q_1^+)}) \\ &\leq C(\|u\|_{L^2(Q_1^+, y^a)} + \|F\|_{C_p^{k+2,\alpha}(Q_1^+)}) \end{aligned}$$

for some $C > 0$ which depends on $N, a, \lambda, \Lambda, r, \alpha$ and $\|A\|_{C_p^{k+2,\alpha}(Q_1^+)}$. Repeating exactly the same argument of Lemma 4.3, we obtain that the function g defined in (4.22) belongs to $C_p^{k+1,\alpha}(Q_r^+)$ and thus $\partial_y u \in C_p^{k+2,\alpha}(Q_r^+)$, which in turn implies $u \in C_p^{k+3,\alpha}(Q_r^+)$. Moreover, by using (4.23), (5.1) and (5.2), one has

$$(5.3) \quad \|\partial_y u\|_{C_p^{k+2,\alpha}(Q_r^+)} \leq C(\|u\|_{L^2(Q_1^+, y^a)} + \|F\|_{C_p^{k+2,\alpha}(Q_1^+)}),$$

for some $C > 0$ which depends on $N, a, \lambda, \Lambda, r, \alpha$ and $\|A\|_{C_p^{k+2,\alpha}(Q_1^+)}$. Then, combining (5.1), (5.2) and (5.3), our statement follows. ■

5.2. Higher order Schauder estimates

Now, we consider the case $f \in C_p^{k,\alpha}$. As remarked in the introduction, when $k = 1$, we cannot use the same argument of the case $f = 0$, since the function $\partial_t f$ is not well defined. In order to overcome this problem, we prove a priori $C_p^{3,\alpha}$ -estimates and combining these with Lemma 5.2 and Lemma 5.3, we obtain our statement in the case $k = 1$. For the general case $k \geq 2$, one could possibly iterate the estimates obtained to prove the main result, as done in the case $f = 0$. However, in order to keep the presentation uniform, we choose to iterate the full procedure (a priori estimates plus approximation) at any step.

Proposition 5.1. *Let $N \geq 1, a > -1, \alpha \in (0, 1), r \in (0, 1)$ and $k \in \mathbb{N}$. Let $A \in C_p^{k+1,\alpha}(Q_1^+)$ satisfy (1.2), let $f \in C_p^{k,\alpha}(Q_1^+)$ and $F \in C_p^{k+1,\alpha}(Q_1^+)$, and let $u \in C_p^{k+2,\alpha}(Q_1^+)$ be a weak solution to (1.1). Then there exists $C > 0$, depending on $N, a, \lambda, \Lambda, r, \alpha$ and $\|A\|_{C_p^{k+1,\alpha}(Q_1^+)}$, such that*

$$(5.4) \quad \|u\|_{C_p^{k+2,\alpha}(Q_r^+)} \leq C(\|u\|_{L^2(Q_1^+, y^a)} + \|f\|_{C_p^{k,\alpha}(Q_1^+)} + \|F\|_{C_p^{k+1,\alpha}(Q_1^+)}).$$

The proof of Proposition 5.1 crucially uses Lemma 5.3 below. In turn, in the proof of Lemma 5.3, we exploit an approximation argument which relies on the following auxiliary result.

Lemma 5.2. *Let $N \geq 1, a > -1, r \in (0, 1), k \in \mathbb{N}$. Let $A \in C_p^{k+2,\alpha}(Q_1^+)$ satisfy (1.2), let $f \in C^\infty(Q_1^+)$, $F \in C_p^{k+2,\alpha}(Q_1^+)$, and let u be a weak solution to (1.1). Then we have that $u \in C_p^{k+3,\alpha}(Q_r^+)$.*

Proof. It is enough to slightly modify the arguments of the proof of Theorem 1.1 in the case $f = 0$. ■

Lemma 5.3. *Let $N \geq 1, a > -1, \alpha \in (0, 1)$ and $k \in \mathbb{N}$. Let $D \in C_p^{k+2,\alpha}(Q_1^+)$ be a diagonal matrix satisfying (1.2), $f \in C_p^{k+1,\alpha}(Q_1^+)$ and $F \in C_p^{k+2,\alpha}(Q_1^+)$. Let $\mu := D_{N+1,N+1}$ and $g := F_{N+1}$. Let $u \in C_p^{k+3,\alpha}(Q_1^+)$ be a weak solution to*

$$(5.5) \quad \begin{cases} y^a \partial_t u - \operatorname{div}(y^a D \nabla u) = y^a f + \operatorname{div}(y^a F) & \text{in } Q_1^+, \\ \lim_{y \rightarrow 0^+} y^a (\mu \partial_y u + g) = 0 & \text{on } \partial^0 Q_1^+. \end{cases}$$

Then the function

$$w := y^{-a} \partial_y \left(y^a \left(\partial_y u + \frac{g}{\mu} \right) \right) \in C_p^{k+1,\alpha}(Q_1^+)$$

is a weak solution to

$$(5.6) \quad \begin{cases} y^a \partial_t w - \operatorname{div}(y^a D \nabla w) = \operatorname{div}(y^a \tilde{F}) & \text{in } Q_r^+, \\ \lim_{y \rightarrow 0^+} y^a (D \nabla w + \tilde{F}) \cdot e_{N+1} = 0 & \text{on } \partial^0 Q_r^+, \end{cases}$$

where

$$(5.7) \quad \tilde{F} := \partial_y D \nabla \left(\partial_y u + \frac{g}{\mu} \right) + \left[\tilde{f} + \partial_y \mu \frac{a}{y} \left(\partial_y u + \frac{g}{\mu} \right) \right] e_{N+1}$$

and

$$(5.8) \quad \tilde{f} := \partial_y f + \partial_y \operatorname{div} g + \operatorname{div}(\partial_y D \nabla u) + \partial_t \left(\frac{g}{\mu} \right) - \operatorname{div} \left(D \nabla \left(\frac{g}{\mu} \right) \right).$$

Moreover,

$$(5.9) \quad \|\tilde{F}\|_{C_p^{k,\alpha}(Q_r^+)} \leq C(\|f\|_{C_p^{k+1,\alpha}(Q_1^+)} + \|F\|_{C_p^{k+2,\alpha}(Q_1^+)} + \|u\|_{C_p^{k+2,\alpha}(Q_r^+)})$$

for some $C > 0$ depending only on $N, a, \lambda, \Lambda, r, \alpha$ and $\|A\|_{C_p^{k+2,\alpha}(Q_1^+)}$.

Proof. Step 1. First, we prove that

$$w := y^{-a} \partial_y \left(y^a \left(\partial_y u + \frac{g}{\mu} \right) \right) = \partial_y \left(\partial_y u + \frac{g}{\mu} \right) + \frac{a}{y} \left(\partial_y u + \frac{g}{\mu} \right) \in C_p^{k+1,\alpha}(Q_1^+).$$

By (1.2), we have $\mu \geq \lambda > 0$ and so

$$\partial_y u + \frac{g}{\mu} \in C_p^{k+2,\alpha}(Q_1^+),$$

thanks to the regularity assumptions on μ, g and μ . By Theorem 2.4, u satisfies the conormal boundary condition

$$(5.10) \quad \lim_{y \rightarrow 0^+} \mu \partial_y u + g = 0,$$

and hence, by Lemma 2.8, we deduce that $\frac{a}{y}(\partial_y u + \frac{g}{\mu}) \in C_p^{k+1,\alpha}(Q_1^+)$, which implies that $w \in C_p^{k+1,\alpha}(Q_1^+)$ by definition of w .

By similar considerations, it follows that $\tilde{f} \in C_p^{k,\alpha}(Q_1^+)$, where \tilde{f} is defined in (5.8). Consequently, $\tilde{F} \in C_p^{k,\alpha}(Q_1^+)$ (defined in (5.7)) and (5.9) directly follows by definition.

Step 2. From this point, we distinguish two cases. If $k = 0$, we assume that $D, f, F \in C^\infty(Q_1^+)$ and thus, by Lemma 5.2, $u \in C^\infty(Q_1^+)$ as well. We will recover our statement under the weaker assumptions $D \in C_p^{2,\alpha}(Q_1^+)$, $f \in C_p^{1,\alpha}(Q_1^+)$ and $F \in C_p^{2,\alpha}(Q_1^+)$ through an approximation argument (see step 3). If $k \geq 1$, such an approximation argument is unnecessary (this is because, when $k \geq 1$, the equation of w is satisfied in the classical sense).

We may rewrite (5.5) as

$$\partial_t u - \operatorname{div}(D \nabla u) - \frac{a}{y}(\mu \partial_y u + g) = f + \operatorname{div} F \quad \text{in } Q_1^+.$$

Differentiating the above equation with respect to y , we obtain

$$(5.11) \quad \begin{aligned} \partial_t(\partial_y u) - \operatorname{div}(D\nabla(\partial_y u)) - \operatorname{div}(\partial_y D\nabla u) - \partial_y \left(\frac{a}{y}(\mu\partial_y u + g) \right) \\ = \partial_y f + \partial_y \operatorname{div} F \quad \text{in } Q_1^+. \end{aligned}$$

Taking into account (5.11) and setting

$$v := y^a \left(\partial_y u + \frac{g}{\mu} \right),$$

we obtain the equation of v :

$$\begin{aligned} y^{-a} \partial_t v - \operatorname{div}(y^{-a} D\nabla v) \\ = \partial_t(\partial_y u) + \partial_t \left(\frac{g}{\mu} \right) - \operatorname{div} \left(D\nabla \left(\partial_y u + \frac{g}{\mu} \right) \right) - \partial_y \left(\frac{a}{y}(\mu\partial_y u + g) \right) \\ = \partial_t(\partial_y u) + \partial_t \left(\frac{g}{\mu} \right) - \operatorname{div}(D\nabla(\partial_y u)) - \operatorname{div} \left(D\nabla \left(\frac{g}{\mu} \right) \right) - \partial_y \left(\frac{a}{y}(\mu\partial_y u + g) \right) \\ = \partial_y f + \partial_y \operatorname{div} \bar{F} + \operatorname{div}(\partial_y D\nabla u) + \partial_t \left(\frac{g}{\mu} \right) - \operatorname{div} \left(D\nabla \left(\frac{g}{\mu} \right) \right) \\ =: \tilde{f} \quad \text{in } Q_1^+, \end{aligned}$$

and thus, recalling that $\mu \geq \lambda > 0$ and (5.10), v satisfies

$$(5.12) \quad \begin{cases} y^{-a} \partial_t v - \operatorname{div}(y^{-a} D\nabla v) = y^{-a}(y^a \tilde{f}) & \text{in } Q_1^+, \\ v = 0 & \text{on } \partial^0 Q_1^+. \end{cases}$$

Differentiating (5.12) with respect to y , we get

$$\partial_t \partial_y v - \operatorname{div}(D\nabla \partial_y v) - \operatorname{div}(\partial_y D\nabla v) - \partial_y \left(\frac{a}{y}(\mu\partial_y v) \right) = \partial_y (y^a \tilde{f}) \quad \text{in } Q_1^+.$$

Consequently, $w = y^{-a} \partial_y v$ and

$$\begin{aligned} y^a \partial_t w - \operatorname{div}(y^a D\nabla w) \\ = \partial_t \partial_y v - \operatorname{div}(D\nabla \partial_y v) + \left(\frac{a}{y}(\mu\partial_y v) \right) = \partial_y (y^a \tilde{f}) + \operatorname{div}(\partial_y D\nabla v) \\ = \partial_y (y^a \tilde{f}) + \operatorname{div} \left(y^a \partial_y D\nabla \left(\partial_y u + \frac{g}{\mu} \right) \right) + \partial_y \left(y^a \partial_y \mu \frac{a}{y} \left(\partial_y u + \frac{g}{\mu} \right) \right) \quad \text{in } Q_1^+. \end{aligned}$$

We need to establish that w satisfies the boundary condition in (5.6). By the regularity assumptions and the fact that $v = 0$ on $\{y = 0\}$, we can take the limit as $y \rightarrow 0^+$ in (5.12) to get

$$\lim_{y \rightarrow 0^+} \left[\mu \partial_{yy} v - \frac{a}{y} \mu \partial_y v + \partial_y \mu \partial_y v + y^a \tilde{f} \right] = \lim_{y \rightarrow 0^+} \left[\partial_t v - \sum_{i=1}^N \partial_{x_i} (D_{i,i} \partial_{x_i} v) \right] = 0,$$

which turns out to be the boundary condition

$$\begin{aligned} 0 &= \lim_{y \rightarrow 0^+} \left[y^a (\mu \partial_y w + \tilde{f}) + \partial_y \mu \partial_y v \right] \\ &= \lim_{y \rightarrow 0^+} y^a \left[\mu \partial_y w + \tilde{f} + \partial_y \mu \partial_y \left(\partial_y u + \frac{g}{\mu} \right) + \partial_y \mu \frac{a}{y} \left(\partial_y u + \frac{g}{\mu} \right) \right]. \end{aligned}$$

Hence, defining \tilde{F} as in (5.7), it follows that w is solution to (5.6) as claimed.

Step 3. In this final step, we present the approximation argument which allows to complete the proof when $k = 0$. First, by Theorem 4.1, we have that

$$\begin{aligned} (5.13) \quad \|\tilde{F}\|_{C_p^{0,\alpha}(Q_1^+)} &\leq C(\|f\|_{C_p^{1,\alpha}(Q_1^+)} + \|F\|_{C_p^{2,\alpha}(Q_1^+)} + \|u\|_{C_p^{2,\alpha}(Q_1^+)}) \\ &\leq C(\|f\|_{C_p^{1,\alpha}(Q_1^+)} + \|F\|_{C_p^{2,\alpha}(Q_1^+)} + \|u\|_{L^2(Q_1^+, y^a)}), \end{aligned}$$

for some $C > 0$ depending only on $N, a, \lambda, \Lambda, r, \alpha$ and $\|D\|_{C_p^{2,\alpha}(Q_1^+)}$.

The proof follows the approximation scheme done in the proof of Theorem 4.1. It is enough to replace the matrix A with the matrix D . Indeed, after regularizing the data (which we call $f_\varepsilon, F_\varepsilon, A_\varepsilon \in C^\infty(Q_1^+)$), and using Lemma 5.2, we can find a family of smooth solutions $v_\varepsilon \in C_c^\infty(Q_r^+)$ to

$$\begin{cases} y^a \partial_t v_\varepsilon - \operatorname{div}(y^a D_\varepsilon \nabla v_\varepsilon) = y^a f_\varepsilon + \operatorname{div}(y^a F_\varepsilon) & \text{in } Q_r^+, \\ \lim_{y \rightarrow 0^+} y^a (D_\varepsilon \nabla v_\varepsilon + F_\varepsilon) \cdot e_{N+1} = 0 & \text{on } \partial^0 Q_r^+. \end{cases}$$

which converges to the original solution u as $\varepsilon \rightarrow 0^+$. Consequently, step 2 yields that

$$w_\varepsilon := y^{-a} \partial_y \left(y^a \left(\partial_y v_\varepsilon + \frac{\tilde{f}_\varepsilon}{\mu_\varepsilon} \right) \right)$$

is a solution to (5.6) (with D and \tilde{F} replaced by D_ε and \tilde{F}_ε) and \tilde{F}_ε , defined analogously to (5.7), satisfies (5.13). By Proposition 4.2 and the Arzelà–Ascoli theorem, one has that $v_\varepsilon \rightarrow u$ in $C_p^{2,\alpha}(Q_r^+)$, which implies that $w_\varepsilon \rightarrow w$ in $C_p^{0,\alpha}(Q_r^+)$. Then a slight modification of the argument in Lemma 4.2 of [4] shows that w_ε converges to w in the energy spaces and that w is a weak solution to (5.6), as claimed. ■

Proof of Proposition 5.1. Let $\partial_i := \partial_{x_i}$ for $i = 1, \dots, N$. We proceed with an induction argument. The step $k = 0$ has been proved in Proposition 4.2. Let us assume that (5.4) holds for $j = 1, \dots, k \in \mathbb{N}$, and let us prove that it is valid for $k + 1$. So, let $u \in C_p^{k+3,\alpha}(Q_1^+)$, $A, F \in C_p^{k+2,\alpha}(Q_1^+)$ and $f \in C_p^{k+1,\alpha}(Q_1^+)$.

Let us fix $0 < r < r' < 1$. First, for every $i = 1, \dots, N$, by Lemma 2.7, one has that $u_i := \partial_i u$ solves (2.12) in $Q_{r'}^+$. Noticing that $u_i \in C_p^{k+2,\alpha}(Q_1^+)$, $A \in C_p^{k+2,\alpha}(Q_1^+)$, $\partial_i f \in C_p^{k,\alpha}(Q_1^+)$ and $\partial_i F, \partial_i A \nabla u \in C_p^{k+1,\alpha}(Q_1^+)$, we can use the inductive step to obtain

$$\begin{aligned} (5.14) \quad \|u_i\|_{C_p^{k+2,\alpha}(Q_r^+)} &\leq C(\|u_i\|_{L^2(Q_{r'}^+, y^a)} + \|\partial_i f\|_{C_p^{k,\alpha}(Q_1^+)} + \|\partial_i F\|_{C_p^{k+1,\alpha}(Q_1^+)}) \\ &\leq C(\|u\|_{L^2(Q_1^+, y^a)} + \|f\|_{C_p^{k+1,\alpha}(Q_1^+)} + \|F\|_{C_p^{k+2,\alpha}(Q_1^+)}) \end{aligned}$$

where $C > 0$ depends only on $N, a, \alpha, \lambda, \Lambda, \|A\|_{C_p^{k+2,\alpha}(Q_1^+)}$. It remains to prove that

$$\begin{aligned} & [u_{yy}]_{C_p^{k,\alpha}(Q_r^+)} + [u_{yy}]_{C_t^{k,(1+\alpha)/2}(Q_r^+)} + [u_{ty}]_{C_p^{k,\alpha}(Q_r^+)} + [u_t]_{C_t^{k,(1+\alpha)/2}(Q_r^+)} \\ & \leq C(\|u\|_{L^2(Q_1^+, y^a)} + \|f\|_{C_p^{k+1,\alpha}(Q_1^+)} + \|F\|_{C_p^{k+2,\alpha}(Q_1^+)}). \end{aligned}$$

Let $D := \text{diag}(A)$. It is immediate to check that u solves

$$\begin{cases} y^a \partial_t u - \text{div}(y^a D \nabla u) = y^a \bar{f} + \text{div}(y^a \bar{F}) & \text{in } Q_r^+, \\ \lim_{y \rightarrow 0^+} y^a (\mu \partial_y u + g) = 0, & \text{on } \partial^0 Q_r^+, \end{cases}$$

where

$$\bar{F} := ((A - D)\nabla u \cdot e_{N+1})e_{N+1} + F$$

and

$$\bar{f} := f + \sum_{i,j=1}^{N+1} \partial_i((A - D)_{i,j} \partial_j u), \quad g = \bar{F}_{N+1} \quad \text{and} \quad \mu = A_{N+1, N+1}.$$

Furthermore, by (5.14) and the definition of \bar{F} and \bar{f} , we have that $\bar{F} \in C_p^{k+2,\alpha}(Q_{r'}^+)$, $\bar{f} \in C_p^{k+1,\alpha}(Q_{r'}^+)$ and

$$(5.15) \quad \begin{aligned} & \|\bar{F}\|_{C_p^{k+2,\alpha}(Q_{r'}^+)} + \|\bar{f}\|_{C_p^{k+1,\alpha}(Q_{r'}^+)} \\ & \leq C(\|u\|_{L^2(Q_1^+, y^a)} + \|f\|_{C_p^{k+1,\alpha}(Q_1^+)} + \|F\|_{C_p^{k+2,\alpha}(Q_1^+)}), \end{aligned}$$

for some $C > 0$ depending only on $N, a, \alpha, \lambda, \Lambda, \|A\|_{C_p^{k+2,\alpha}(Q_1^+)}$.

By Lemma 5.3, the function $w := y^{-a} \partial_y (y^a (\partial_y u + g/\mu))$ belongs to $C_p^{k+1,\alpha}(Q_{r'}^+)$ and is a weak solution to

$$\begin{cases} y^a \partial_t w - \text{div}(y^a D \nabla w) = \text{div}(y^a \tilde{F}) & \text{in } Q_{r'}^+, \\ \lim_{y \rightarrow 0^+} y^a (D \nabla w + \tilde{F}) \cdot e_{N+1} = 0 & \text{on } \partial^0 Q_{r'}^+, \end{cases}$$

where \tilde{F} is defined in (5.7), with f and F replaced by \bar{f} and \bar{F} , respectively. Furthermore, $\tilde{F} \in C_p^{k,\alpha}(Q_{r'}^+)$, so, by the inductive assumption (noticing that in the case $k = 0$ we use Theorem 2.4), (5.9), (5.14) and (5.15), we obtain that $w \in C_p^{k+1,\alpha}(Q_{r'}^+)$ and

$$\|w\|_{C_p^{k+1,\alpha}(Q_{r'}^+)} \leq C(\|u\|_{L^2(Q_1^+, y^a)} + \|f\|_{C_p^{k+1,\alpha}(Q_1^+)} + \|F\|_{C_p^{k+2,\alpha}(Q_1^+)}),$$

for some $C > 0$ which depends only on $N, a, \alpha, \lambda, \Lambda, \|A\|_{C_p^{k+2,\alpha}(Q_1^+)}$. Now, by the same arguments of Lemma 4.3 and by Lemma 2.9, it follows that

$$(\partial_y u + g/\mu)(x, y, t) = \frac{1}{y^a} \int_0^y s^a w(x, s, t) ds,$$

satisfies $\partial_y (\partial_y u + \bar{f}/\mu) \in C^{k+1,\alpha}$ and, by the regularity of g and μ , we deduce

$$\begin{aligned} & [u_{yy}]_{C_p^{k,\alpha}(Q_r^+)} + [u_{yy}]_{C_t^{k,(1+\alpha)/2}(Q_r^+)} \\ & \leq C(\|u\|_{L^2(Q_1^+, y^a)} + \|f\|_{C_p^{k+1,\alpha}(Q_1^+)} + \|F\|_{C_p^{k+2,\alpha}(Q_1^+)}), \end{aligned}$$

for some $C > 0$ depending only on $N, a, \alpha, \lambda, \Lambda, \|A\|_{C_p^{k+2,\alpha}(Q_1^+)}$.

To conclude the proof, it is sufficient to observe that

$$\partial_t u = y^{-a} \operatorname{div}(y^a (A \nabla u + F)) + f \in C_p^{k+1, \alpha}(Q_r^+),$$

which immediately implies

$$\|\partial_t u\|_{C_p^{k+1, \alpha}(Q_r^+)} \leq C(\|u\|_{L^2(Q_1^+, y^a)} + \|f\|_{C_p^{k+1, \alpha}(Q_1^+)} + \|F\|_{C_p^{k+2, \alpha}(Q_1^+)}),$$

for some $C > 0$ depending only on $N, a, \alpha, \lambda, \Lambda, \|A\|_{C_p^{k+2, \alpha}(Q_1^+)}$. ■

Proof of Theorem 1.1. Once Proposition 5.1 and Lemma 5.2 are established, our statement follows by approximation as in Theorem 4.1. ■

6. Cylindrically curved characteristic manifolds

In this section, we show how to extend the $C_p^{k+2, \alpha}$ regularity estimates to weak solutions of a class of equations having weights vanishing or exploding on curved characteristic manifolds Γ , as in (1.8).

Proof of Corollary 1.3. The proof follows the one of Corollary 1.3 in [4]; after composing with a standard local diffeomorphism, one may apply the main Theorem 1.1.

Indeed, let us consider the classical diffeomorphism

$$\Phi(x, y) = (x, y + \varphi(x)),$$

which is of class $C^{k+2, \alpha}$ and then $C_p^{k+2, \alpha}$ extending constantly in the time variable. Up to a dilation, one has that $\tilde{u} := u \circ (\Phi(x), t)$ is a weak solution to

$$\begin{cases} \tilde{\delta}^a \partial_t \tilde{u} - \operatorname{div}(\tilde{\delta}^a \tilde{A} \nabla \tilde{u}) = \tilde{\delta}^a \tilde{f} + \operatorname{div}(\tilde{\delta}^a \tilde{F}) & \text{in } Q_1^+, \\ \lim_{y \rightarrow 0^+} \tilde{\delta}^a (\tilde{A} \nabla \tilde{u} + \tilde{F}) \cdot e_{N+1} = 0 & \text{on } \partial^0 Q_1^+. \end{cases}$$

where $\tilde{\delta} = \delta \circ \Phi$,

$$\tilde{f} = f \circ (\Phi(x), t), \quad \tilde{F} = J_{\Phi}^{-1} F \circ (\Phi(x), t) \quad \text{and} \quad \tilde{A} = (J_{\Phi}^{-1})(A \circ (\Phi(x), t))(J_{\Phi}^{-1})^T.$$

We have that $\tilde{\delta} \in C^{k+2, \alpha}(B_1^+)$, $\tilde{A}, \tilde{F} \in C^{k+1, \alpha}(B_1^+)$ and $\tilde{f} \in C^{k, \alpha}(B_1^+)$. Moreover, by using Lemma 2.8, $\tilde{\delta}$ satisfies

$$\begin{aligned} \tilde{\delta} > 0 & \text{ in } B_1^+, \quad \tilde{\delta} = 0 & \text{ on } \partial^0 B_1^+, \quad \partial_y \tilde{\delta} > 0 & \text{ on } \partial^0 B_1^+, \\ \tilde{\delta}/y & \in C^{k+1, \alpha}(B_1^+), \quad \tilde{\delta}/y \geq \mu > 0 & \text{ in } \bar{B}_1^+, \end{aligned}$$

where the last non-degeneracy condition is a consequence of the assumption $|\nabla \delta| \geq c_0 > 0$.

Defining

$$b(z) := \left(\frac{\tilde{\delta}(z)}{y} \right)^a \in C^{k+1, \alpha}(B_1^+),$$

one has

$$\begin{aligned}
 0 &= \int_{Q_1^+} y^a b (-\tilde{u} \partial_t \phi + \tilde{A} \nabla \tilde{u} \cdot \nabla \phi - \tilde{f} \phi + \tilde{F} \cdot \nabla \phi) \\
 &= \int_{Q_1^+} y^a (-\tilde{u} \partial_t (\phi b) + \tilde{A} \nabla \tilde{u} \cdot \nabla (\phi b) - \tilde{A} \nabla \tilde{u} \cdot \nabla b \phi - \tilde{f} (\phi b) + \tilde{F} \cdot \nabla (\phi b) - \tilde{F} \cdot \nabla b \phi),
 \end{aligned}$$

so, $b\phi$ being an admissible test function, we deduce that \tilde{u} is a weak solution to

$$\begin{cases}
 y^a \partial_t \tilde{u} - \operatorname{div}(y^a \tilde{A} \nabla \tilde{u}) = y^a \tilde{g} + \operatorname{div}(y^a \tilde{F}) & \text{in } Q_1^+, \\
 \lim_{y \rightarrow 0^+} y^a (\tilde{A} \nabla \tilde{u} + \tilde{F}) \cdot e_{N+1} = 0 & \text{on } \partial^0 Q_1^+,
 \end{cases}$$

where

$$\tilde{g} := \tilde{f} + \frac{\tilde{A} \nabla \tilde{u} \cdot \nabla b}{b} + \frac{\tilde{F} \cdot \nabla b}{b}.$$

Finally, we apply a recursive argument to prove the $C_p^{k+2,\alpha}$ -regularity of \tilde{u} , which in turn extends to the same regularity for the original u by composing back with the diffeomorphism.

Let $k = 0$. We notice that $u \in C_p^{1,\alpha}$ by Corollary 1.3 in [4], and hence, after composing with the $C_p^{2,\alpha}$ diffeomorphism, one has $\nabla \tilde{u} \in C_p^{0,\alpha}$, which gives that $\tilde{g} \in C_p^{0,\alpha}$. Then the $C_p^{2,\alpha}$ -regularity of \tilde{u} follows by Theorem 1.1.

Finally, one may iterate this reasoning for any $k \geq 1$ by replacing the use of the starting result Corollary 1.3 in [4] with the present Corollary 1.3 at a lower step. ■

7. Parabolic higher order boundary Harnack principle

This last section, is devoted to the proof of the higher order boundary Harnack principle in Theorem 1.4.

Proof of Theorem 1.4. First, the regularity assumptions of boundaries, coefficients and data for the equations in (1.9) do guarantee that $u, v \in C_{\text{loc}}^{k+2,\alpha}(\bar{\Omega} \cap Q_1)$, by classical theory of uniformly parabolic equations (for instance, see [32]). Hence, the equations in (1.9) are satisfied both in the weak sense and pointwisely in $\Omega \cap Q_1$. From this, we deduce a pointwise equation for the quotient $w = v/u$ in $\Omega \cap Q_1$, that is,

$$(7.1) \quad u^2 \partial_t w - \operatorname{div}(u^2 A \nabla w) = u f - v g + u^2 b \cdot \nabla w.$$

Now, let us define the standard diffeomorphism

$$\Phi(x, y, t) := (x, y + \varphi(x, t), t),$$

which is of class $C_p^{k+2,\alpha}$. Let us set

$$\tilde{u} = u \circ \Phi, \tilde{v} = v \circ \Phi, \quad \tilde{f} = f \circ \Phi, \quad \tilde{g} = g \circ \Phi,$$

and define

$$\tilde{A} = (J_{z,\Phi}^{-1})^T (A \circ \Phi) J_{z,\Phi}^{-1}, \quad \tilde{b} = J_{z,\Phi}^{-1} b \circ \Phi,$$

where $J_{z,\Phi}$ is the square block $[c_{ij}]_{i,j=1,\dots,N+1}$ of the Jacobian $J_\Phi := [c_{ij}]_{i,j=1,\dots,N+2}$.

Since w solves (7.1), then, up to dilations, $\tilde{w} = w \circ \Phi = \tilde{v}/\tilde{u}$ solves

$$(7.2) \quad y^2 \mu^2 \partial_t \tilde{w} - \operatorname{div}(y^2 \mu^2 \tilde{A} \nabla \tilde{w}) = y \left(\mu \tilde{f} - \frac{\tilde{v}}{y} \tilde{g} \right) + y^2 \mu^2 \tilde{b} \cdot \nabla \tilde{w} + y^2 \mu^2 c \cdot \nabla \tilde{w},$$

pointwisely in Q_1^+ , where $\mu = \tilde{u}/y$ and $c = \partial_t \varphi e_{N+1}$. Now some remarks on the regularity of the data of the weighted equation above are in order. First, by Lemma 2.8 and the non-degeneracy condition $u(z, t) \geq c_0 d_p((z, t), \partial\Omega \cap Q_1)$ in (1.9), we can infer that

$$0 < \bar{c}_0 \leq \mu \in C_p^{k+1, \alpha}(B_1^+).$$

Thanks to the previous information, we can rewrite (7.2) dividing by μ^2 as

$$(7.3) \quad y^2 \partial_t \tilde{w} - \operatorname{div}(y^2 \tilde{A} \nabla \tilde{w}) = y h + y^2 \tilde{b} \cdot \nabla \tilde{w},$$

where

$$\tilde{b} = \tilde{b} + c + 2\tilde{A}^T \frac{\nabla \mu}{\mu} \in C_p^{k, \alpha} \quad \text{and} \quad h = \frac{\mu \tilde{f} - \frac{\tilde{v}}{y} \tilde{g}}{\mu^2} \in C_p^{k+1, \alpha}.$$

Moreover, since

$$\tilde{w} = \frac{\tilde{v}/y}{\tilde{u}/y},$$

again by Lemma 2.8 and the $C_p^{k+2, \alpha}$ -regularity of \tilde{u}, \tilde{v} , we have $\tilde{w} \in C_p^{k+1, \alpha}(Q_1^+)$ which has two implications. First, the drift term in (7.3) can be considered as a forcing term, that is, $\tilde{b} \cdot \nabla \tilde{w} = \tilde{f} \in C_p^{k, \alpha}(Q_1^+)$. Second, $\tilde{w} \in L^2(I_1; H^1(B_1^+, y^a)) \cap L^\infty(I_1; L^2(B_1^+, y^a))$ and, by multiplying the equation (7.3) by test functions $\phi \in C_c^\infty(Q_1^+)$ and integrating by parts, one gets that \tilde{w} is a weak solution to

$$\begin{cases} y^2 \partial_t \tilde{w} - \operatorname{div}(y^2 \tilde{A} \nabla \tilde{w}) = \operatorname{div}(y^2 H) + y^2 \tilde{f} & \text{in } Q_1^+, \\ \lim_{y \rightarrow 0^+} y^2 (\tilde{A} \nabla \tilde{w} + H) \cdot e_{N+1} = 0 & \text{on } \partial^0 Q_1^+, \end{cases}$$

where the field

$$H(x, y, t) = \frac{e_{N+1}}{y^2} \int_0^y s h(x, s, t) ds$$

belongs to $C_p^{k+1, \alpha}(Q_1^+)$ by Lemma 2.9.

Then the regularity $C_p^{k+2, \alpha}$ -regularity of \tilde{w} follows by Theorem 1.1. Finally, the same regularity is inherited by w by composing back with the diffeomorphism. ■

Funding. The three authors are research fellows of Istituto Nazionale di Alta Matematica INDAM group GNAMPA. The project has been partially funded by the project ‘‘Anomalous diffusion equations: regularity and geometric properties of solutions and free boundaries’’ (MUR funding for Young Researchers-Seal of Excellence SOE_0000194 (ADE)), the PRIN project ‘‘NO3-Nodal Optimization, NONlinear elliptic equations, NONlocal geometric problems, with a focus on regularity’’ (2022R537CS) and the INDAM-GNAMPA projects ‘‘Regolarità e singolarità in problemi di frontiere libere’’ (CUP_E53C22001930001), ‘‘Teoria della regolarità per problemi ellittici e parabolici con diffusione anisotropa e pesata’’ (CUP_E53C22001930001) and ‘‘PDE ellittiche che degenerano su varietà di dimensione bassa e frontiere libere molto sottili’’ (CUP_E5324001950001).

References

- [1] Allen, M., Kriventsov, D. and Shahgholian, H.: The free boundary for semilinear problems with highly oscillating singular terms. Preprint 2024, arXiv:2405.10418v1.
- [2] Athanasopoulos, I., Caffarelli, L. and Milakis, E.: [On the regularity of the non-dynamic parabolic fractional obstacle problem](#). *J. Differential Equations* **265** (2018), no. 6, 2614–2647. Zbl [1394.35588](#) MR [3804726](#)
- [3] Audrito, A.: [On the existence and Hölder regularity of solutions to some nonlinear Cauchy–Neumann problems](#). *J. Evol. Equ.* **23** (2023), no. 3, article no. 58, 45 pp. Zbl [1521.35065](#) MR [4629496](#)
- [4] Audrito, A., Fioravanti, G. and Vita, S.: [Schauder estimates for parabolic equations with degenerate or singular weights](#). *Calc. Var. Partial Differential Equations* **63** (2024), no. 8, article no. 204, 46 pp. Zbl [1547.35123](#) MR [4788276](#)
- [5] Audrito, A. and Terracini, S.: [On the nodal set of solutions to a class of nonlocal parabolic equations](#). *Mem. Amer. Math. Soc.* **301** (2024), no. 1512, v+118 pp. Zbl [07927092](#) MR [4808714](#)
- [6] Banerjee, A., Danielli, D., Garofalo, N. and Petrosyan, A.: [The structure of the singular set in the thin obstacle problem for degenerate parabolic equations](#). *Calc. Var. Partial Differential Equations* **60** (2021), no. 3, article no. 91, 52 pp. Zbl [1469.35130](#) MR [4249869](#)
- [7] Banerjee, A. and Garofalo, N.: [A parabolic analogue of the higher-order comparison theorem of De Silva and Savin](#). *J. Differential Equations* **260** (2016), no. 2, 1801–1829. Zbl [1331.35066](#) MR [3419746](#)
- [8] Banerjee, A. and Garofalo, N.: [Monotonicity of generalized frequencies and the strong unique continuation property for fractional parabolic equations](#). *Adv. Math.* **336** (2018), 149–241. Zbl [1400.35045](#) MR [3846151](#)
- [9] Banerjee, A. and Garofalo, N.: [On the space-like analyticity in the extension problem for nonlocal parabolic equations](#). *Proc. Amer. Math. Soc.* **151** (2023), no. 3, 1235–1246. Zbl [1505.35009](#) MR [4531651](#)
- [10] Biswas, A. and Stinga, P. R.: [Regularity estimates for nonlocal space-time master equations in bounded domains](#). *J. Evol. Equ.* **21** (2021), no. 1, 503–565. Zbl [1464.35392](#) MR [4238215](#)
- [11] Bucur, C. and Karakhanyan, A. L.: [Potential theoretic approach to Schauder estimates for the fractional Laplacian](#). *Proc. Amer. Math. Soc.* **145** (2017), no. 2, 637–651. Zbl [1355.26007](#) MR [3577867](#)
- [12] Caffarelli, L., Mellet, A. and Sire, Y.: [Traveling waves for a boundary reaction-diffusion equation](#). *Adv. Math.* **230** (2012), no. 2, 433–457. Zbl [1255.35072](#) MR [2914954](#)
- [13] Caffarelli, L. and Silvestre, L.: [An extension problem related to the fractional Laplacian](#). *Comm. Partial Differential Equations* **32** (2007), no. 8, 1245–1260. Zbl [1143.26002](#) MR [2354493](#)
- [14] Caffarelli, L. A. and Stinga, P. R.: [Fractional elliptic equations, Caccioppoli estimates and regularity](#). *Ann. Inst. H. Poincaré C Anal. Non Linéaire* **33** (2016), no. 3, 767–807. Zbl [1381.35211](#) MR [3489634](#)
- [15] Chiarenza, F. and Serapioni, R.: [A remark on a Harnack inequality for degenerate parabolic equations](#). *Rend. Sem. Mat. Univ. Padova* **73** (1985), 179–190. Zbl [0588.35013](#) MR [799906](#)

- [16] Danielli, D., Garofalo, N., Petrosyan, A. and To, T.: [Optimal regularity and the free boundary in the parabolic Signorini problem](#). *Mem. Amer. Math. Soc.* **249** (2017), no. 1181, v+103 pp. Zbl [1381.35249](#) MR [3709717](#)
- [17] De Silva, D. and Savin, O.: [A note on higher regularity boundary Harnack inequality](#). *Discrete Contin. Dyn. Syst.* **35** (2015), no. 12, 6155–6163. Zbl [1336.35096](#) MR [3393271](#)
- [18] De Silva, D. and Savin, O.: [A short proof of boundary Harnack principle](#). *J. Differential Equations* **269** (2020), no. 3, 2419–2429. Zbl [1439.35110](#) MR [4093736](#)
- [19] De Silva, D. and Savin, O.: [On the parabolic boundary Harnack principle](#). *Matematica* **1** (2022), no. 1, 1–18. Zbl [07968274](#) MR [4482724](#)
- [20] Dong, H., Jeon, S. and Vita, S.: [Schauder type estimates for degenerate or singular elliptic equations with DMO coefficients](#). *Calc. Var. Partial Differential Equations* **63** (2024), no. 9, article no. 239, 42 pp. Zbl [07942221](#) MR [4821888](#)
- [21] Dong, H. and Kim, D.: [Parabolic and elliptic systems in divergence form with variably partially BMO coefficients](#). *SIAM J. Math. Anal.* **43** (2011), no. 3, 1075–1098. Zbl [1230.35045](#) MR [2800569](#)
- [22] Dong, H. and Kim, D.: [Schauder estimates for a class of non-local elliptic equations](#). *Discrete Contin. Dyn. Syst.* **33** (2013), no. 6, 2319–2347. Zbl [1263.45008](#) MR [3007688](#)
- [23] Dong, H. and Phan, T.: [Parabolic and elliptic equations with singular or degenerate coefficients: the Dirichlet problem](#). *Trans. Amer. Math. Soc.* **374** (2021), no. 9, 6611–6647. Zbl [1477.35099](#) MR [4302171](#)
- [24] Dong, H. and Phan, T.: [On parabolic and elliptic equations with singular or degenerate coefficients](#). *Indiana Univ. Math. J.* **72** (2023), no. 4, 1461–1502. Zbl [1529.35141](#) MR [4637368](#)
- [25] Fabes, E. B., Kenig, C. E. and Serapioni, R. P.: [The local regularity of solutions of degenerate elliptic equations](#). *Comm. Partial Differential Equations* **7** (1982), no. 1, 77–116. Zbl [0498.35042](#) MR [643158](#)
- [26] Fernández-Real, X. and Ros-Oton, X.: [Regularity theory for elliptic PDE](#). Zur. Lect. Adv. Math. 28, EMS Press, Berlin, 2022. Zbl [1522.35001](#) MR [4560756](#)
- [27] Garofalo, N. and Ros-Oton, X.: [Structure and regularity of the singular set in the obstacle problem for the fractional Laplacian](#). *Rev. Mat. Iberoam.* **35** (2019), no. 5, 1309–1365. Zbl [1433.35466](#) MR [4018099](#)
- [28] Hyder, A., Segatti, A., Sire, Y. and Wang, C.: [Partial regularity of the heat flow of half-harmonic maps and applications to harmonic maps with free boundary](#). *Comm. Partial Differential Equations* **47** (2022), no. 9, 1845–1882. Zbl [1496.35132](#) MR [4472924](#)
- [29] Jeon, S. and Vita, S.: [Higher order boundary Harnack principles in Dini type domains](#). *J. Differential Equations* **412** (2024), 808–856. Zbl [07950709](#) MR [4794088](#)
- [30] Kukuljan, T.: [Higher order parabolic boundary Harnack inequality in \$C^1\$ and \$C^{k,\alpha}\$ domains](#). *Discrete Contin. Dyn. Syst.* **42** (2022), no. 6, 2667–2698. Zbl [1490.35061](#) MR [4421508](#)
- [31] Ladyženskaja, O. A., Solonnikov, V. A. and Ural'ceva, N. N.: [Linear and quasi-linear equations of parabolic type](#). Transl. Math. Monogr. 23, American Mathematical Society, Providence, RI, 1968; translation from Izdat. “Nauka”, Moscow, 1967. Zbl [0174.15403](#) MR [0241822](#)
- [32] Lieberman, G. M.: [Second order parabolic differential equations](#). World Scientific, River Edge, NJ, 1996. Zbl [0884.35001](#) MR [1465184](#)

- [33] Lions, J.-L. and Magenes, E.: *Non-homogeneous boundary value problems and applications. Vol. I.* Grundlehren Math. Wiss. 181, Springer, New York-Heidelberg, 1972. Zbl [0223.35039](#) MR [350177](#)
- [34] Mazzeo, R.: *Elliptic theory of differential edge operators. I.* *Comm. Partial Differential Equations* **16** (1991), no. 10, 1615–1664. Zbl [0745.58045](#) MR [1133743](#)
- [35] Mazzeo, R. and Vertman, B.: *Elliptic theory of differential edge operators, II: Boundary value problems.* *Indiana Univ. Math. J.* **63** (2014), no. 6, 1911–1955. Zbl [1311.58014](#) MR [3298726](#)
- [36] Metafuno, G., Negro, L. and Spina, C.: *A unified approach to degenerate problems in the half-space.* *J. Differential Equations* **351** (2023), no. 6, 63–99. Zbl [1511.35182](#) MR [4530863](#)
- [37] Metafuno, G., Negro, L. and Spina, C.: *Regularity theory for parabolic operators in the half-space with boundary degeneracy.* Preprint 2024, arXiv:[2309.14319v2](#).
- [38] Metafuno, G., Negro, L. and Spina, C.: *Singular parabolic problems in the half-space.* *Studia Math.* **277** (2024), no. 1, 1–44. Zbl [1550.35287](#) MR [4792088](#)
- [39] Negro, L. and Spina, C.: *Kernel bounds for parabolic operators having first-order degeneracy at the boundary.* Preprint 2024, arXiv:[2403.01959v1](#).
- [40] Nyström, K. and Sande, O.: *Extension properties and boundary estimates for a fractional heat operator.* *Nonlinear Anal.* **140** (2016), 29–37. Zbl [1381.35230](#) MR [3492726](#)
- [41] Restrepo, D. and Ros-Oton, X.: *C^∞ regularity in semilinear free boundary problems.* Preprint 2024, arXiv:[2407.20426v1](#).
- [42] Silvestre, L.: *On the differentiability of the solution to an equation with drift and fractional diffusion.* *Indiana Univ. Math. J.* **61** (2012), no. 2, 557–584. Zbl [1308.35042](#) MR [3043588](#)
- [43] Simon, J.: *Compact sets in the space $L^p(0, T; B)$.* *Ann. Mat. Pura Appl. (4)* **146** (1987), 65–96. Zbl [0629.46031](#) MR [916688](#)
- [44] Simon, L.: *Schauder estimates by scaling.* *Calc. Var. Partial Differential Equations* **5** (1997), no. 5, 391–407. Zbl [0946.35017](#) MR [1459795](#)
- [45] Sire, Y., Terracini, S. and Tortone, G.: *On the nodal set of solutions to degenerate or singular elliptic equations with an application to s -harmonic functions.* *J. Math. Pures Appl. (9)* **143** (2020), 376–441. Zbl [1471.35151](#) MR [4163134](#)
- [46] Sire, Y., Terracini, S. and Vita, S.: *Liouville type theorems and regularity of solutions to degenerate or singular problems part I: even solutions.* *Comm. Partial Differential Equations* **46** (2021), no. 2, 310–361. Zbl [1471.35152](#) MR [4207950](#)
- [47] Sire, Y., Terracini, S. and Vita, S.: *Liouville type theorems and regularity of solutions to degenerate or singular problems part II: odd solutions.* *Math. Eng.* **3** (2021), no. 1, article no. 5, 50 pp. Zbl [1496.35126](#) MR [4144100](#)
- [48] Stinga, P. R. and Torrea, J. L.: *Regularity theory and extension problem for fractional nonlocal parabolic equations and the master equation.* *SIAM J. Math. Anal.* **49** (2017), no. 5, 3893–3924. Zbl [1386.35419](#) MR [3709888](#)
- [49] Terracini, S., Tortone, G. and Vita, S.: *Higher order boundary Harnack principle via degenerate equations.* *Arch. Ration. Mech. Anal.* **248** (2024), no. 2, article no. 29, 44 pp. Zbl [1537.35109](#) MR [4726059](#)

- [50] Terracini, S., Tortone, G. and Vita, S.: A priori regularity estimates for equations degenerating on nodal sets. Preprint 2024, arXiv:[2404.06980v2](https://arxiv.org/abs/2404.06980v2).
- [51] Zhikov, V. V.: [Weighted Sobolev spaces](#). *Sb. Math.* **189** (1998), no. 7-8, 1139–1170; translation from *Mat. Sb.* **189** (1998), no. 8, 27–58. Zbl [0919.46026](#) MR [1669639](#)

Received March 27, 2024; revised October 31, 2024.

Alessandro Audrito

Dipartimento di Scienze Matematiche “Giuseppe Luigi Lagrange”, Politecnico di Torino
Corso Duca degli Abruzzi 24, 10129 Torino, Italy;
alessandro.audrito@polito.it

Gabriele Fioravanti

Dipartimento di Matematica “Giuseppe Peano”, Università di Torino
Via Carlo Alberto 10, 10124 Torino, Italy;
gabriele.fioravanti@unito.it

Stefano Vita

Dipartimento di Matematica “Felice Casorati”, Università di Pavia
Via Ferrata 5, 27100 Pavia, Italy;
stefano.vita@unipv.it