

Temporal Analysis of PM Concentrations in Turin: Assessment and Data Elaboration for Future Correlation with Human Health Implications

*Original*

Temporal Analysis of PM Concentrations in Turin: Assessment and Data Elaboration for Future Correlation with Human Health Implications / Gallione, Davide; Mastromatteo, Nicole; Matina, Mariagrazia; Clerico, Marina. - ELETTRONICO. - (2025). ( 10thWorldCongressonCivil,Structural,andEnvironmentalEngineering(CSEE'25) Barcellona (ES) 10-12 April, 2025) [10.11159/iceptp25.151].

*Availability:*

This version is available at: 11583/3001110 since: 2025-06-19T08:42:14Z

*Publisher:*

International ASET Inc.

*Published*

DOI:10.11159/iceptp25.151

*Terms of use:*

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

(Article begins on next page)

# Temporal Analysis of PM Concentrations in Turin: Assessment and Data Elaboration for Future Correlation with Human Health Implications

**Davide Gallione<sup>1</sup>, Nicole Mastromatteo<sup>1</sup>, Mariagrazia Matina<sup>1</sup>, Marina Clerico<sup>1</sup>**

<sup>1</sup>Politecnico Di Torino, DIATI – Department of Environment, Land and Infrastructure Engineering,  
Corso Duca degli Abruzzi, 24, Turin, Italy

davide.gallione@polito.it; [nicole.mastromatteo@polito.it](mailto:nicole.mastromatteo@polito.it); [mariagrazia.matina95@gmail.com](mailto:mariagrazia.matina95@gmail.com); [marina.clerico@polito.it](mailto:marina.clerico@polito.it)

**Abstract** - As urban populations continue to grow, understanding the dynamics of particulate matter (PM) concentrations in these areas is increasingly important. This study investigated the temporal variations of PM<sub>1</sub>, PM<sub>2.5</sub>, and PM<sub>10</sub> in the urban area of Turin, located in the Po Valley, Italy, utilizing high-resolution data from a monitoring campaign over a four-year period (2020–2024), focusing on identifying seasonal and weekly variations. The results revealed significant differences in PM concentrations between different seasons. The findings reveal a strong seasonality, with higher PM levels in winter due to domestic heating, traffic emissions, and adverse meteorological conditions, while summer months show lower concentrations. Winter concentrations often exceed WHO air quality guidelines, with PM<sub>10</sub> levels surpassing EU annual thresholds, emphasizing the need for stricter emission control policies during colder months when pollution poses significant health risks. Weekly fluctuations in PM concentrations were also observed, with peaks mid-week and at the end of each week. These fluctuations are likely influenced by human activities and meteorological factors, suggesting that interventions targeting specific periods could help reduce pollution levels. The use of the Palas Fidas 200S instrument, with its high temporal resolution, allowed for detailed examination of daily and hourly trends, offering insights into the dynamics of PM concentrations. These findings are critical for epidemiological studies examining the link between air pollution and public health outcomes, particularly cardiovascular diseases. This study is part of a future research effort that integrates both concentration and epidemiological data, offering a comprehensive understanding of the effects of air pollution on human health.

**Keywords:** particulate matter, air quality, urban pollution, fix monitoring station, temporal variability, PM<sub>1</sub>, PM<sub>2.5</sub>, PM<sub>10</sub>.

## 1. Introduction

Air pollution can cause a number of harmful health effects, including premature mortality. An estimated 6.7 million deaths per year are attributed to air pollution globally. Italy has the highest number of deaths caused by fine particulate matter; in 2020, premature deaths due to PM<sub>2.5</sub> were reported to be 52,303, according to Eurostat sources. Air pollution impacts various pathophysiological processes both at the target organ level and systemically<sup>1</sup>. Beyond premature mortality, air pollution also contributes to morbidity, which refers to the number of disease cases recorded during a specific period in relation to the total population surveyed<sup>2</sup>. Understanding how particulate matter varies across space and time in different areas is essential for accurately assessing the health risks associated with air pollution<sup>3</sup>. Most epidemiological studies on short-term exposure have relied on daily or hourly variations in pollutant concentrations measured at air quality monitoring stations<sup>4</sup>. In 2004, the American Heart Association released its first scientific statement on “Air Pollution and Cardiovascular Disease”, concluding that exposure to particulate matter contributes to cardiovascular morbidity and mortality. At the national level, Italy’s Istituto Superiore di Sanità (ISS) recently updated the “Sentieri” report, which examines mortality and morbidity among residents in areas of national interest (SIN). Alongside epidemiological evidence linking air pollution to oncological and respiratory diseases, the report highlighted excess hospitalizations for circulatory system diseases in both sexes, as well as ischemic heart disease and cardiovascular diseases, even at a young age<sup>1</sup>. In recent years, the number of time-series studies investigating associations between daily mortality, morbidity, and ambient concentrations of PM<sub>10</sub> and PM<sub>2.5</sub> has grown significantly<sup>2</sup>.

Studies confirm that a 10  $\mu\text{g}/\text{m}^3$  increase in PM10 raises the risk of hospitalization for myocardial infarction (MI) <sup>5</sup>, while short-term exposure to high PM2.5 levels increases MI risk within hours in high-risk populations. Elevated PM2.5 levels are linked to a higher incidence of atherosclerosis. Short-term PM increases are associated with ischemic stroke, cardiac ischemia, and coronary disease, as well as variations in blood pressure <sup>6</sup>. Fine particulate inhalation may also raise the risk of deep vein thrombosis and blood clotting <sup>6</sup>. Mechanisms include endothelial dysfunction, platelet activation, and coagulation changes, contributing to plaque instability, rupture, and clot formation, potentially leading to MI or stroke <sup>6</sup>.

Specifically, a 10  $\mu\text{g}/\text{m}^3$  increase in PM concentrations has been shown to correlate with an increased risk of hospitalization on the same day or the day following exposure in most studies. Studies such as that of <sup>7</sup> have shown effects up to 2-3 days later, and rarely up to 5 days after exposure <sup>8</sup>. It is also significant to note that an increased risk of hospitalization was observed in the hour immediately following a 10  $\mu\text{g}/\text{m}^3$  increase in PM concentration <sup>9</sup>. These results suggest the importance of considering both a daily and hourly time scale in future analyses.

The present study is positioned as a complementary part of a larger study including both concentration and epidemiological data, giving the possibility to provide an in-depth understanding of the impact of air pollution on human health. This study investigated the temporal and spatial variabilities of PM1, PM2.5, and PM10 over the urban area of Turin in the Po Valley, Italy, based on high-resolution data from the monitoring station of University of Turin Polytechnic (DIATI) CC-Green-Roof-Lab working since 2018. In particular, the years 2020 to 2024 were analysed.

## 2. Materials and Methods

PM data, used for this study, were acquired from the CC-Green-Roof-Lab, an open-air laboratory. In particular, the Palas Fidas 200S instrument was used. This is an optical spectrometer designed for air pollution monitoring and certified according to EN 16450. The Palas Fidas 200S analyzer is an instrument that can measure the concentration of solid particles ranging in size from 180 nm to 18 microns. The parameters acquired are PM1, PM2.5, PM4, PM10, and PTS. The external configuration of the instrument includes a stainless-steel container to protect it from the weather and keep the instrument at an optimal temperature for measurements. The instrument is equipped with a sensor and an optical spectrometer that allows the definition of the number of aerosol particles and their size, thanks to the scattering phenomenon. The sampling head can capture a flow of 0.3  $\text{m}^3/\text{h}$  and enables representative measurements even in intense wind, thanks to the Sigma-2 sampling head with VDI 2119-4. The instrument is equipped with an IADS (Intelligent Aerosol Drying System), from Palas GmbH Germany, as well as sensors for measuring the temperature, air pressure, and relative humidity. The IADS prevents incorrect measurements caused by condensation effects in areas with high humidity. The temperature must be kept higher than the dew temperature, which is why a mechanism inside the drying line regulates it according to the external temperature, pressure, and rH <sup>10</sup>.

The objective of the proposed analysis is to identify the exceeding of the concentration limits and the critical periods in which the greatest increases in PM concentrations occurred, in order to make the results available for future epidemiological studies and, specifically, to be able to make the correlation between exceedances and hospital admissions for cardiovascular-related problems. In order to understand the evolution of particulate concentrations over time, graphs representing the trend of the PM1, PM2.5 and PM10 fractions over the period under consideration (1/1/2020-31/7/2024) were created using the Python programming environment. This shows the changes in concentrations on an hourly basis, highlighting any peaks and fluctuations over the course of the different seasons due mainly to domestic heating, heavy traffic and weather conditions. Thanks to the high temporal resolution of the data, it was possible to visualize specific conditions and analyze particular situations.

## 3. Discussion and Results

In this study, trends in PM concentrations over approximately four years, from 2020 to July 2024, are analyzed. Analyzing this very extensive time series, it can be seen that, almost cyclically, PM concentrations are higher in the winter periods and remain lower in the summer months, generating a repetitiveness in the trend of concentrations over the years, this is very evident from Fig. 1. Turin is exposed to the Po Valley in two directions and is surrounded by hilly relief on one side and the Alpine chain on the other <sup>11</sup>. This setting affects the city's atmospheric stability, which in winter contributes to the

high number of pollution events <sup>12</sup>. The increase in concentrations in the winter months may be due to the increase in emissions from domestic heating and vehicular traffic during typical winter weather conditions that favour the accumulation of pollutants.




Fig. 1: Trends in daily PM concentrations between 01/01/2020 and 31/07/2024 measured at the Politecnico di Torino. From top to bottom PM1, PM2.5 and PM10. These last two graphs also show the concentration limits of the main legislative regulations in force.

PM1, PM2.5 and PM10 show rather similar trends, although the concentrations are different. In fact, PM10 has the highest peak concentrations compared to PM2.5 and PM1, which, with a few exceptions, show very close values. This suggests that PM1 and PM2.5 tend to behave more uniformly, whereas PM10, which includes larger particles, reaches more pronounced peaks, probably related to specific emission sources or local events such as dust uplift. Several studies have evaluated the impacts of typical winter heating sources, such as domestic boilers, in the area under study and the entire Po Valley <sup>13</sup>. The concentration was lowest in the summer months, especially in May. The spring months showed a gradual reduction in concentrations, whereas the autumn months showed an increase. The same trend was also found by <sup>14,15</sup>.




Fig. 2: Monthly average concentrations for each year for the years 2020 (blue), 2021 (orange), 2022 (green), 2023 (red), 2024 (purple) for (a) PM<sub>1</sub>; (b) PM<sub>2.5</sub> and (c) PM<sub>10</sub>. It is specified that 2024 is incomplete with the winter period as it is still in progress: the analysis stops in July 2024.

Starting with the daily averages in the graph in Fig. 2, monthly averages are presented to emphasize the already evident seasonality presented and explained above (Fig. 1). This allowed us to extend the analysis to a monthly scale in order to also visualize special air pollution situations. This happened in November 2020, when very high PM values were recorded. According to Arpa Piemonte data, in 2020, Turin reached 36 days of exceeding the limit of  $50 \mu\text{g}/\text{m}^3$  of PM<sub>10</sub>, which is the maximum allowed by current regulations. This data shows how the city has frequently exceeded the legal limits for fine particles, posing a risk to public health. It is important to note that, although the restrictions related to the COVID-19 pandemic led to a reduction in industrial activities and traffic, air pollution continued to pose a significant challenge. Several

studies have found that, during the lockdown, there was a decrease in higher PM10 concentrations, but not a uniform reduction in all pollutants <sup>16</sup>.




Fig. 3: Averages of PM1, PM2.5 and PM10 concentrations, divided between the winter months (October to March) and the summer months (April to September) for the period from 2020 to 2024. It is specified that 2024 is incomplete with the winter period as it is still in progress: the analysis stops in July 2024.

Fig. 3 shows the average concentrations of PM1, PM2.5, and PM10, divided into winter months (October to March) and summer months (April to September) for the period 2020 to 2024. This chart aims to contextualize the data concerning current regulations and WHO air quality guidelines, with the goal of verifying whether winter conditions alone contribute to exceeding the annual limits set by various regulatory agencies. From the analysis of the graph, it is observed that during the winter months, the PM2.5 averages range between 20 and 30  $\mu\text{g}/\text{m}^3$ , consistently exceeding the new European threshold of 10  $\mu\text{g}/\text{m}^3$ . PM10 also shows winter averages well above the 20  $\mu\text{g}/\text{m}^3$  limit established by the European Directive 2024/2881, with values exceeding 40  $\mu\text{g}/\text{m}^3$  in certain years. During the summer months, although concentrations are generally lower, occasional exceedances of the 20  $\mu\text{g}/\text{m}^3$  threshold for PM10 are observed, along with values close to or exceeding the 10  $\mu\text{g}/\text{m}^3$  threshold for PM2.5. Moreover, it is evident that both in summer and winter, PM2.5 and PM10 concentrations significantly exceed the WHO guideline limits of 5  $\mu\text{g}/\text{m}^3$  for PM2.5 and 15  $\mu\text{g}/\text{m}^3$  for PM10. In particular, during the winter months, PM10 concentrations are more than twice the WHO limit. Finally, comparing the data with the current limits set by Legislative Decree 155/2010, PM2.5 concentrations during winter are generally compliant in most years but approach the critical threshold of 25  $\mu\text{g}/\text{m}^3$ . In summer, values remain below this threshold. Regarding PM10, winter averages frequently reach or exceed the 40  $\mu\text{g}/\text{m}^3$  limit, as highlighted in the cases of 2022 and 2024. In summer, values generally remain below the 40  $\mu\text{g}/\text{m}^3$  threshold.

In order to examine the variation of particulate concentrations in relation to seasonality, the following graphs were made showing the monthly trend of average PM1, PM2.5 and PM10 concentrations between 2020 and 2024.




Fig. 4: Trend of daily PM in a given month (above) February; (below) April. The different fractions of particulate matter are indicated with lines of different colors specified in the legend and the end of each week is highlighted with a dashed black vertical line.

Fig. 4 illustrates the daily trends of particulate matter concentrations over one month, with different particulate fractions represented by lines of various colours (specified in the legend). The end of each week is marked by a dashed black vertical line. Each graph starts from the first Monday and extends to the month's end, allowing visualization of entire weeks from Monday to Sunday. This approach helps analyze weekly trends, compare patterns across days within the week, and examine differences across weeks over the five years considered in the analysis.

In February, every years show peaks in PM concentrations around mid and late month. Specifically, in 2020, 2022, and 2024, significant increases are observed between February 11–17 and again around February 23–25. This suggests a possible weekly periodicity, with PM concentrations rising and falling in roughly seven-day intervals. For example, in 2020, a peak around February 12 is followed by a decrease and another increase around February 18. A similar trend is seen in 2022 and 2024. However, in 2021, the pattern appears less regular, with relatively stable concentrations until the latter half of the month, when a single major peak occurs between February 23–25. In 2023, concentrations begin to rise around February 8, remain high with minor fluctuations until February 20, and then drop sharply. While 2021 and 2023 deviate slightly from the weekly trend observed in other years, they still display some oscillatory behavior.

In April, variability is noted across years, but PM10 concentrations are consistently higher than PM2.5 and PM1. Some years exhibit periodic declines aligned with roughly weekly intervals. However, the overall pattern is irregular, making it difficult to establish a clear weekly trend. Notably, peaks often occur between April 10–20 in several years (e.g., 2022 and 2024), hinting at a possible but inconsistent weekly pattern.

#### 4. Conclusion

This study analysed PM1, PM2.5, and PM10 concentrations in Turin over a four-year period (2020–2024), identifying critical periods and trends related to seasonal and weekly variations. The results highlight a clear seasonality in PM levels, with higher concentrations during the winter months due to domestic heating, vehicular emissions, and unfavourable meteorological conditions, and lower concentrations in the summer. Winter PM concentrations consistently exceeded the WHO air quality guidelines, with PM10 levels frequently surpassing the European Union's annual thresholds.

These findings underscore the urgent need for stricter policies to control emissions, particularly during the colder months, when pollution levels pose a significant threat to public health. Moreover, the study observed periodic weekly fluctuations, with noticeable peaks occurring mid-week and towards the end of each week. This cyclic behaviour, likely linked to human activities and meteorological factors, emphasizes the importance of targeted interventions during specific periods to mitigate PM levels. The high temporal resolution of the Palas Fidas 200S instrument allowed for a detailed examination of daily and even hourly trends, providing valuable insights into the dynamics of PM concentrations. These findings are crucial for future epidemiological studies, particularly in assessing the correlation between air pollution exceedances and hospital admissions for cardiovascular diseases.

However, further investigations are needed to evaluate the impact of additional factors, such as industrial emissions and long-range transport of pollutants, as well as to explore mitigation strategies aimed at reducing PM concentrations in urban areas. In conclusion, this research contributes to the growing body of knowledge on air quality in urban environments and highlights the necessity of comprehensive monitoring programs and proactive measures to improve public health and environmental quality in the Po Valley region.

#### References

- [1] (1) Belfiore, A.; Portincasa, P.; Ciaula, A. D. INQUINAMENTO ATMOSFERICO, ATEROSCLEROSI E RISCHIO CARDIOVASCOLARE.
- [2] (2) Cassee, F. R.; Mills, N. L.; Newby, D. E. *Cardiovascular Effects of Inhaled Ultrafine and Nano-Sized Particles*; Wiley, 2011.
- [3] (3) Liu, S.; Tian, H.; Luo, L.; Bai, X.; Zhu, C.; Lin, S.; Zhao, S.; Zhang, K.; Hao, J.; Guo, Z.; Lv, Y. Health Impacts and Spatiotemporal Variations of Fine Particulate and Its Typical Toxic Constituents in Five Urban Agglomerations of China. *Science of The Total Environment* **2022**, *806*, 151459. <https://doi.org/10.1016/j.scitotenv.2021.151459>.
- [4] (4) Atkinson, R. W.; Butland, B. K.; Dimitroulopoulou, C.; Heal, M. R.; Stedman, J. R.; Carslaw, N.; Jarvis, D.; Heaviside, C.; Vardoulakis, S.; Walton, H.; Anderson, H. R. Long-Term Exposure to Ambient Ozone and Mortality: A Quantitative Systematic Review and Meta-Analysis of Evidence from Cohort Studies. *BMJ Open* **2016**, *6* (2), e009493. <https://doi.org/10.1136/bmjopen-2015-009493>.
- [5] (5) Zanobetti A, Schwartz J. The Effect of Particulate Air Pollution on Emergency Admissions for Myocardial Infarction: A Multicity Case-Crossover Analysis. *Environ Health Perspect.* 2005 Aug;113(8):978-82. Doi: 10.1289/Ehp.7550. PMID: 16079066; PMCID: PMC1280336.
- [6] (6) Polichetti, G.; Cocco, S.; Spinali, A.; Trimarco, V.; Nunziata, A. Effects of Particulate Matter (PM10, PM2.5 and PM1) on the Cardiovascular System. *Toxicology* **2009**, *261* (1), 1–8. <https://doi.org/10.1016/j.tox.2009.04.035>.
- [7] (7) Fisher, J. A.; Puett, R. C.; Laden, F.; Wellenius, G. A.; Sapkota, A.; Liao, D.; Yanosky, J. D.; Carter-Pokras, O.; He, X.; Hart, J. E. Case-Crossover Analysis of Short-Term Particulate Matter Exposures and Stroke in the Health Professionals Follow-up Study. *Environment International* **2019**, *124*, 153–160. <https://doi.org/10.1016/j.envint.2018.12.044>.

- [8] (8) Wu, H.; Zhang, B.; Wei, J.; Lu, Z.; Zhao, M.; Liu, W.; Bovet, P.; Guo, X.; Xi, B. Short-Term Effects of Exposure to Ambient PM<sub>1</sub>, PM<sub>2.5</sub>, and PM<sub>10</sub> on Ischemic and Hemorrhagic Stroke Incidence in Shandong Province, China. *Environmental Research* **2022**, *212*, 113350. <https://doi.org/10.1016/j.envres.2022.113350>.
- [9] (9) Tabaghi, S.; Sheibani, M.; Khaheshi, I.; Miri, R.; Haji Aghajani, M.; Safi, M.; Eslami, V.; Pishgahi, M.; Alipour Parsa, S.; Namazi, M. H.; Beyranvand, M. R.; Sohrabifar, N.; Hassanian-Moghaddam, H.; Pourmotahari, F.; Khaiat, S.; Akbarzadeh, M. A. Associations between Short-term Exposure to Fine Particulate Matter and Acute Myocardial Infarction: A Case-crossover Study. *Clinical Cardiology* **2023**, *46* (11), 1319–1325. <https://doi.org/10.1002/clc.24111>.
- [10] (10) Mastromatteo, N.; Drudi, L.; Gallione, D.; Bellopede, R.; Clerico, M. Assessment of Environmental Parameters in Natural Coastal Scenery and Compositional by Means of an Innovative Approach. *Atmosphere* **2024**, *15* (11), 1379. <https://doi.org/10.3390/atmos15111379>.
- [11] (11) Mecca, D.; Boanini, C.; Vaccaro, V.; Gallione, D.; Mastromatteo, N.; Clerico, M. Spatial Variation, Temporal Evolution, and Source Direction Apportionment of PM<sub>1</sub>, PM<sub>2.5</sub>, and PM<sub>10</sub>: 3-Year Assessment in Turin (Po Valley). *Environ Monit Assess* **2024**, *196* (12), 1251. <https://doi.org/10.1007/s10661-024-13446-9>.
- [12] (12) Pernigotti, D.; Georgieva, E.; Thunis, P.; Bessagnet, B. Impact of Meteorology on Air Quality Modeling over the Po Valley in Northern Italy. *Atmospheric Environment* **2012**, *51*, 303–310. <https://doi.org/10.1016/j.atmosenv.2011.12.059>.
- [13] (13) Gilardoni, S.; Massoli, P.; Marinoni, A.; Mazzoleni, C.; Freedman, A.; Lonati, G.; De Iuliis, S.; Gianelle, V. Spatial and Temporal Variability of Carbonaceous Aerosol Absorption in the Po Valley. *Aerosol Air Qual. Res.* **2020**, *20* (12), 2624–2639. <https://doi.org/10.4209/aaqr.2020.03.0085>.
- [14] (14) Chen, T.; He, J.; Lu, X.; She, J.; Guan, Z. Spatial and Temporal Variations of PM<sub>2.5</sub> and Its Relation to Meteorological Factors in the Urban Area of Nanjing, China. *IJERPH* **2016**, *13* (9), 921. <https://doi.org/10.3390/ijerph13090921>.
- [15] (15) Xu, G.; Jiao, L.; Zhang, B.; Zhao, S.; Yuan, M.; Gu, Y.; Liu, J.; Tang, X. Spatial and Temporal Variability of the PM<sub>2.5</sub>/PM<sub>10</sub> Ratio in Wuhan, Central China. *Aerosol Air Qual. Res.* **2017**, *17* (3), 741–751. <https://doi.org/10.4209/aaqr.2016.09.0406>.
- [16] (16) Deserti, M.; Raffaelli, K.; Ramponi, L.; Carbonara, C.; Agostini, C.; Amorati, R.; Arvani, B.; Giovannini, G.; Maccaferri, S.; Poluzzi, V. Report COVID-19-Studio Preliminare Degli Effetti Delle Misure COVID-19 Sulle Emissioni in Atmosfera e Sulla Qualità Dell'aria nel Bacino Padano-Giugno 2020; Technical Report, prepAIR Project, 2020., 2020. Available online: [https://www.lifeprepare.eu/wp-content/uploads/2020/06/COVIDQA-Prepair-19Giugno2020\\_final.pdf](https://www.lifeprepare.eu/wp-content/uploads/2020/06/COVIDQA-Prepair-19Giugno2020_final.pdf).