

Type IV optic nerve and Onodi cell: is there a risk of injury during sphenoid sinus surgery?

*Original*

Type IV optic nerve and Onodi cell: is there a risk of injury during sphenoid sinus surgery? / Fadda, Gian Luca; Urbanelli, Anastasia; Petrelli, Alessio; Trossarello, Marta; Nitro, Letizia; Saibene, Alberto Maria; De Corso, Eugenio; Gned, Dario; Panfili, Marco; Cavallo, Giovanni. - In: ACTA OTORHINOLARYNGOLOGICA ITALICA. - ISSN 1827-675X. - 44:1(2024), pp. 36-41. [10.14639/0392-100x-n2462]

*Availability:*

This version is available at: 11583/3002558 since: 2025-08-26T11:16:57Z

*Publisher:*

Pacini Editore

*Published*

DOI:10.14639/0392-100x-n2462

*Terms of use:*

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

(Article begins on next page)

RHINOLOGY

# Type IV optic nerve and Onodi cell: is there a risk of injury during sphenoid sinus surgery?

Gian Luca Fadda<sup>1</sup>, Anastasia Urbanelli<sup>1</sup>, Alessio Petrelli<sup>2</sup>, Marta Trossarelli<sup>1</sup>, Letizia Nitro<sup>3</sup>, Alberto Maria Saibene<sup>3</sup>, Eugenio De Corso<sup>4</sup>, Dario Gned<sup>5</sup>, Marco Panfil<sup>6</sup>, Giovanni Cavallo<sup>1</sup>

<sup>1</sup> Department of Otorhinolaryngology, University of Turin, "San Luigi Gonzaga" Hospital, Orbassano (TO), Italy; <sup>2</sup> National Institute for Health, Migration and Poverty (INMP), Rome, Italy; <sup>3</sup> Otolaryngology Unit, "Santi Paolo e Carlo" Hospital, Department of Health Sciences, University of Milan, Milan, Italy; <sup>4</sup> Otorhinolaryngology, "A. Gemelli" University Hospital IRCCS, Rome, Italy; <sup>5</sup> Department of Diagnostic Imaging, University of Turin, "San Luigi Gonzaga" Hospital, Orbassano (TO), Italy; <sup>6</sup> Unit of Radiology, "A. Gemelli" University Hospital Foundation IRCCS, Rome, Italy

## SUMMARY

**Objective.** This study aims to determine the prevalence and types of Onodi cells through computed tomography and investigate the relationship between Onodi cell and the surrounding structures, paying particular attention to the risky proximity to the optic nerve canal.

**Methods.** In this study, 430 computed tomography scans of paranasal sinuses were analysed to establish the prevalence and different types of Onodi cells. Furthermore, the relationship between Onodi cell and different patterns of sphenoid sinus pneumatization and surrounding structures were investigated. Special attention was paid to the relationship between Onodi cell and the optic nerve canal, particularly in cases when the optic nerve canal was bulging by more than 50% into the Onodi cell (Type IV).

**Results.** The Onodi cell was detected in 21.6% of cases, with the most common being Type I (48.5% right, 54.3% left). Type IV bulging of the optic nerve canal into the Onodi cell was observed in 47.1% of cases on the right side, 41.2% on the left side and bilateral in 11.7% of cases.

**Conclusions.** In our series, we observed a high prevalence of Type IV optic nerve bulging into the Onodi cell. For this reason, we suggest that clinicians should always try to identify it in a pre-operative setting with computed tomography to avoid catastrophic consequences during endoscopic sinus surgery approaching the sphenoid area.

**KEY WORDS:** Onodi cell, optic nerve, endoscopic sinus surgery, sphenoid sinus, computed tomography

## Introduction

Adolf Ónodi (1857-1919), professor of rhinology, described *cellula sphenoidaloidalis* (after that, known as "Onodi cell") in 1903 as a condition in which the most posterior or lateral ethmoid air cell extends posteriorly to lie lateral and/or superior to the sphenoid sinus (SS) <sup>1</sup>.

Onodi cell (OC) is of particular interest during endoscopic sinus surgery (ESS) because of its close relationship with the optic nerve canal (ONC) and the internal carotid artery (ICA), making it crucial to examine computed tomography (CT) scans pre-operatively to avoid serious injuries <sup>2-4</sup>.

Another reason why OC must be detected prior to surgery is because an OC often causes the natural ostium of the sphenoid sinus to be located more superiorly, and it is necessary for the surgeon to be aware of this anatomical variant when operating.

Multiple studies have demonstrated an increased prevalence of OC identification when examining CT scans on two to three axial planes <sup>4-7</sup>.

Our study aims to investigate, through the analysis of 430 CT scans (860 sides), the prevalence and the patterns of OC and sphenoid sinus pneumatization

Received: January 7, 2023

Accepted: June 13, 2023

Published online: December 29, 2023

### Correspondence

Gian Luca Fadda

E-mail: dott.fadda@gmail.com

**How to cite this article:** Fadda GL, Urbanelli A, Petrelli A, et al. Type IV optic nerve and Onodi cell: is there a risk of injury during sphenoid sinus surgery? Acta Otorhinolaryngol Ital 2024;44:36-41. <https://doi.org/10.14639/0392-100X-N2462>

© Società Italiana di Otorinolaringoiatria e Chirurgia Cervico-Facciale



OPEN ACCESS

This is an open access article distributed in accordance with the CC-BY-NC-ND (Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International) license. The article can be used by giving appropriate credit and mentioning the license, but only for non-commercial purposes and only in the original version. For further information: <https://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>


(SSP) and the relationship between OC and the ONC, with particular attention to its bulging into the OC as it represents a relevant risk factor for critical and potentially irreversible consequences during ESS.

## Materials and methods

This is a retrospective study including the analyses of paranasal sinus CT images of 430 patients, analysing a total of 860 SS), collected at a 1.5 mm slice thickness as part of the pre-surgical assessment for patients undergoing ESS for chronic rhinosinusitis. Patients were enrolled between November 2017 and December 2022 at “San Luigi Gonzaga” University Hospital. Inclusion criteria were individuals older than 18 years with sinonasal symptoms (nasal obstruction, chronic sinusitis, rhinorrhoea) undergoing sinus surgery for the first time. Patients with sinonasal malignancies, facial bone fractures, previous head trauma and massive nasal polyposis that could modify normal anatomy were excluded. High-resolution CT images were performed in three axial planes and without contrast medium, as is normally done in pre-operative assessment protocols. Images were reviewed and analysed independently by three sinus surgeons and two radiologists with extensive and long-lasting sinus surgery experience using a commercially available software viewer (window OsiriX).

Based on CT scans, the following variables were assessed:

- sagittal, coronal, and axial planes of all analysed CT were used to identify the presence of OC and its localisation (right, left, or bilateral). OC was identified as an ethmoidal cells extending posteriorly or laterally to the anterior wall of the SS;
- according to Thimmaiah et al. <sup>8</sup>, three patterns of OC pneumatisation were assessed: Type I (superior) if the sphenothmoidal cell was located above and medial to the sphenoid sinus (Fig. 1), Type II (superolateral) if the air cell was located both above and below to a horizontal line drawn at the uppermost part of SS in a coronal plane (Fig. 2), and Type III (lateral) if sphenothmoidal cell was located below the horizontal line (Fig. 3);
- the relationship between ONC and OC wall was evaluated according to the studies of Ozdemir et al. <sup>2</sup> and Chmielik et al. <sup>24</sup>. The grade of protrusion of ONC was calculated by measuring the diameter of the ONC on the axial and sagittal plane, focusing specifically on Type IV which defines ONC bulging into the OC by more than 5 mm or more than 50% of the circumference of the ONC in two planes (axial or sagittal plane). It is worth to note that this classification of types of ONC are defined only in relation to an OC, and not to any other cells or the SS;
- the presence of the ICA bulging into an OC;



**Figure 1.** Onodi cell (\*) Type I pattern of pneumatisation is above and medial to the horizontal line. (1a) Coronal, (1b) Axial and (1c) Sagittal, paranasal CT scan. A horizontal line was drawn at the uppermost part of the sphenoid sinus (s) in the coronal image (1a). Optic canals (white arrows) with bulging > 5 mm (Type IV) are shown.


- the relationship between OC and different patterns of SSP (sellar, presellar, conchal and postsellar type) was evaluated considering the location of posterior sinus wall concerning the position of the sella turcica on the sagittal plane <sup>9</sup>.

### Statistical analysis


The associations between variables of interest were evaluated using the Chi-square test. If the expected frequencies were less than 5 for more than 25% of the cells in the contingency tables, Fisher’s exact test was performed. The analyses were conducted using SAS System 9.4 (SAS Inc, Cary, NC, USA).

## Results

CT scans of 430 patients were analysed retrospectively (mean age,  $53 \pm 18.9$  years; range, 18-92). 57% of patients were male ( $n = 245$ ) and 43% female ( $n = 185$ ). OC was identified in 93 of 430 scans (21.6%). Of these, 35.5% were detected on the right side, 37.6% on the left side, and 26.9% were bilateral. Type I was the most prevalent, accounting for 16 scans (48.5%) on the right side, 19 (54.3%) on the left side, and 35 scans considering the totality of OC detected simultaneously on the right and left sides (Tab. I); 18.3% of



**Figure 2.** Onodi cell (\*) Type II is above and below the horizontal line. Axial (2b) CT image showing protrusion of ICA (double arrow). (2a) Coronal, (2b) Axial and (2c) Sagittal, paranasal CT scan. A horizontal line was drawn at the uppermost part of the sphenoid sinus (s) in the coronal image (2a). Optic canals (white arrows) with bulging > 5 mm (Type IV) are shown.



**Figure 3.** Onodi cell (\*) Type III is below the horizontal line. **(3d)** Endoscopic right sphenoid sinus, Onodi cell and optic canal are shown. **(3a)** Coronal, **(3b)** Axial and **(3c)** Sagittal, paranasal CT scan. A horizontal line was drawn at the uppermost part of the sphenoid sinus (s) in the coronal image **(3a)**. Optic canals (white arrows) with bulging > 5 mm (Type IV) are shown.

cases presented a bulging of the ONC into the OC by more than 50% of the circumference (Type IV). Of these, 47.1% occurred on the right side, 41.2% on the left side and 11.7% (only 2 cases) on both left and right side. ICA protruded into the OC in 8 scans (8.6%) (Tab. I). The presence of OC was significantly associated with the presence of a Type IV-ONC for both sides ( $p$  value < 0.0001). The presence of bilateral OCs also showed a statistically significant association with Type IV-ONC coexistence ( $p$  value < 0.0001).

The association between the presence of Type IV-ONC and all the different types of OC was also investigated. Our re-

sults demonstrated a significant association between the Type IV-ONC and all OC sub-types ( $p$  value < 0.01) for both sides, except for Type III-OC on the left ( $p$  value = 0.59).

Analysis of SSP associated to the presence of OC identified a predominance of the sellar type (60.6% on the right side, 65.7% on the left side), followed by the postsellar type (27.3% right and 20% left) and presellar type (12.1% right and 11% left) (Tab. II). Type I-OC was more represented among cases associated with the sellar type of SSP, both on the right side (9 cases) and on the contralateral side (14 cases). Only one case of Type I-OC was associated with the presence of the conchal type of SSP (left-sided).

## Discussion

In the present investigation, we reported an incidence of OC of 21.6%. Similar percentages were described by Oz-turan et al. (16.6%)<sup>10</sup>, while many previous studies showed extremely variable percentages ranging between 8% and 65%<sup>3,11-19</sup>. This inconsistency in the prevalence of OC may be due to the different methods used for its identification (radiological vs anatomical). Tanaviratananich et al.<sup>17</sup> found a prevalence of 60% during studies conducted on cadavers, and similar results were reported in the works by Kantarci et al. (60%)<sup>13</sup> and Tomovic et al. (65.3%)<sup>12</sup>.

**Table I.** Relation between side and type of Onodi cell and bulging Type IV of the optic nerve canal on 430 paranasal sinus CT scans.

|                     | N (%)                | Men<br>N (%)        | Women<br>N (%)      | Bulging<br>ONC (type IV)     | Protrusion ICA    |
|---------------------|----------------------|---------------------|---------------------|------------------------------|-------------------|
| <b>R-Onodi cell</b> | <b>33 (35.5)</b>     | <b>18 (29)</b>      | <b>15 (48.4)</b>    | <b>8 (47.1)</b>              | <b>3 (37.5)</b>   |
|                     |                      |                     |                     | <b>(p value &lt; 0.0001)</b> |                   |
| Type I              | 16 (48.5)            | 9 (50.0)            | 7 (46.7)            | 5 (62.5)                     | 2 (66.7)          |
|                     |                      |                     |                     | (p value < 0.0001)           |                   |
| Type II             | 11 (33.3)            | 6 (33.3)            | 5 (33.3)            | 2 (25.0)                     | 1 (33.3)          |
|                     |                      |                     |                     | (p value < 0.0001)           |                   |
| Type III            | 6 (18.2)             | 3 (16.7)            | 3 (20.0)            | 1 (12.5)                     | -                 |
|                     |                      |                     |                     | (p value < 0.001)            |                   |
| <b>L-Onodi cell</b> | <b>35 (37.6)</b>     | <b>27 (43.5)</b>    | <b>8 (25.8)</b>     | <b>7 (41.2)</b>              | <b>3 (37.5)</b>   |
|                     |                      |                     |                     | <b>(p value &lt; 0.0001)</b> |                   |
| Type I              | 19 (54.3)            | 14 (51.9)           | 5 (62.5)            | 4 (57.1)                     | 2 (66.7)          |
|                     |                      |                     |                     | (p value < 0.0001)           |                   |
| Type II             | 9 (25.7)             | 7 (25.9)            | 2 (25)              | 3 (42.9)                     | 1 (33.3)          |
|                     |                      |                     |                     | (p value < 0.0001)           |                   |
| Type III            | 7 (20.0)             | 6 (22.2)            | 1 (12.5)            | -                            | -                 |
|                     |                      |                     |                     | (p value = 0.59)             |                   |
| <b>B-Onodi cell</b> | <b>25 (26.9)</b>     | <b>17 (27.5)</b>    | <b>8 (25.8)</b>     | <b>2 (11.7)</b>              | <b>2 (25.0)</b>   |
|                     |                      |                     |                     | <b>(p value &lt; 0.0001)</b> |                   |
| <b>Total</b>        | <b>93/430 (21.6)</b> | <b>62/93 (66.7)</b> | <b>31/93 (33.3)</b> | <b>17/93 (18.3)</b>          | <b>8/93 (8.6)</b> |

ONC: optic nerve canal; R: right; L: left; B: bilateral.

Furthermore, many of the studies were carried out analysing only one<sup>12</sup> or two CT planes (axial and/or coronal). In our study, the identification of OC was performed on all three CT scan planes (coronal, axial and sagittal) to obtain a more accurate evaluation of this anatomical variant.

Regarding the prevalence of sub-types of OC, Thimmaiah et al.<sup>8</sup> described three main patterns of pneumatization, with Type II being the most common, followed by Type I and Type III. In our study, Type I was the most common followed by Type II. Our percentages correlate with those found in the previous literature: in the study by Ali et al.<sup>3</sup>, Type I was the most prevalent, while Chmielik et al.<sup>4</sup> reported that Type II was the most common pattern (Tab. III). The improvement of systematic pre-operative CT analysis to detect the presence of OC and their pattern of pneumatization can decrease the risk of ONC injury during ESS, thus avoiding catastrophic consequences such as blindness<sup>8</sup>. Özdemir et al.<sup>2</sup> reported four types of relationships between the ONC and OC, using the classification first described by Chmielik et al. in their study<sup>4</sup>. Type IV-ONC in relationship with the OC was described by some authors as the most dangerous pattern, as the potential risk during ESS is greater when more than 50% of the circumference of ONC is bulging into the OC<sup>20</sup>. We found 17 cases (18.3%) of ONC presenting a bulging into the OC for more than 50% (Type IV), representing a relevant risk factor for injury during ESS. The dangerous relationship between OC and ONC has been pointed out previously by several authors<sup>2,4,11,12,21</sup>, but most considered the totality of ONC bulging, without differentiating those that protruded more than 50% (Type IV).

On the other hand, Yeoh et al.<sup>20</sup> found a prevalence of 51% of ONC protruding into the OC and concluded that the most dangerous type of ONC was the one with no bulging, because of the extremely thin wall separating the sphenoidal air cell and the ONC itself.

Another risk of injury during sphenoid sinus surgery is represented by the proximity of OC to the ICA. In this study, 8 cases were found (8.6%) where the ICA protruded into the OC, thus exposing the endoscopic procedure to the risk of a catastrophic haemorrhage. Ozturan et al.<sup>10</sup> reported an ICA protrusion rate of 2.8% in absence of an OC, but a prevalence of protruding ICA of 59% when the OC was present. They also concluded that increased dimension of OC significantly correlated with higher rates of protrusion of ICA. The relationship between different types of OC and different patterns of SSP were also evaluated. According to the Güldner classification<sup>9</sup>, four types (sellar, presellar, conchal, and postsellar types) of SSP were considered. The sellar type was most commonly associated with the presence of OC, followed by the postsellar type. We did not find any similar correspondence in all previous analysed studies.

It is important to analyse CT scans before sinus surgery on all three axial planes, as stated before and based on evidence in the literature. The sagittal plane is the most important for OC identification, since it provides information on the antero-posterior extension of the most posterior ethmoidal cell in relation to the sphenoid sinus.

Regarding complications during ESS in the presence of OC, in addition to accurate pre-operative review of the patient's surgical anatomy, the surgeon's experience and care-

**Table II.** Relation between the type of Onodi cell and type of sphenoid sinus on 430 paranasal sinus CT scans.

|                     | Sellar<br>N (%)      |                      | Presellar<br>N (%)  |                    | Conchal<br>N (%)    |                     | Postsellar<br>N (%) |                     | Total                |
|---------------------|----------------------|----------------------|---------------------|--------------------|---------------------|---------------------|---------------------|---------------------|----------------------|
| <b>R-Onodi cell</b> | <b>20/33 (60.6)</b>  |                      | <b>4/33 (12.1)</b>  |                    | <b>-</b>            |                     | <b>9/33 (27.3)</b>  |                     | <b>33/93 (35.5)</b>  |
| Type I              | 9/20 (45.0)          |                      | 2/4 (50.0)          |                    |                     |                     | 5/9 (55.6)          |                     |                      |
| Type II             | 8/20 (40.0)          |                      | 2/4 (50.0)          |                    |                     |                     | 1/9 (11.1)          |                     |                      |
| Type III            | 3/20 (15.0)          |                      | -                   |                    |                     |                     | 3/9 (33.3)          |                     |                      |
| <b>L-Onodi cell</b> | <b>23/35 (65.7)</b>  |                      | <b>4/35 (11.4)</b>  |                    | <b>1/35 (2.9)</b>   |                     | <b>7/35 (20.0)</b>  |                     | <b>35/93 (37.6)</b>  |
| Type I              | 14/23 (60.9)         |                      | 3/4 (75.0)          |                    | 1/1 (100.0)         |                     | 1/7 (14.3)          |                     |                      |
| Type II             | 4/23 (17.4)          |                      | 1/4 (25.0)          |                    | -                   |                     | 4/7 (57.1)          |                     |                      |
| Type III            | 5/23 (21.7)          |                      | -                   |                    | -                   |                     | 2/7 (28.6)          |                     |                      |
| <b>B-Onodi cell</b> | <b>R</b>             | <b>L</b>             | <b>R</b>            | <b>L</b>           | <b>R</b>            | <b>L</b>            | <b>R</b>            | <b>L</b>            |                      |
|                     | <b>13/25* (52.0)</b> | <b>14/25* (56.0)</b> | <b>3/25* (12.0)</b> | <b>2/25* (8.0)</b> | <b>3/25* (12.0)</b> | <b>5/25* (20.0)</b> | <b>6/25* (24.0)</b> | <b>4/25* (16.0)</b> | <b>25*/93 (26.9)</b> |
| Type I              | 11/13 (84.6)         | 12/14 (85.7)         | 2/3 (66.7)          | 1/2 (50.0)         | 2/3 (66.7)          | 1/5 (20.0)          | 4/6 (66.7)          | 2/4 (50.0)          |                      |
| Type II             | 2/13 (15.4)          | -                    | 1/3 (33.3)          | 1/2 (50.0)         | -                   | 4/5 (80.0)          | -                   | 2/4 (50.0)          |                      |
| Type III            | -                    | 2/14 (14.3)          | -                   | -                  | 1/3 (33.3)          | -                   | 2/6 (33.3)          | -                   |                      |
| <b>Total</b>        | <b>93/430 (21.6)</b> |                      |                     |                    |                     |                     |                     |                     |                      |

R: right; L: left; B: bilateral.

\* (25\*) This refers to the total number of patients with bilateral OC (in total 50 OC in this group).

**Table III.** Comparison of Onodi cell incidence in the literature.

|                               | No. PNSCT | No. PNSCT sides | No. OC (%)  | No. R-side (%) | No. L-side (%) | No. Bilateral (%) | Bulging ONC (%) |
|-------------------------------|-----------|-----------------|-------------|----------------|----------------|-------------------|-----------------|
| Kasemsiri, 2011 <sup>16</sup> | 187       | 374             | 92 (49.7)*  | -              | -              | -                 | -               |
| Shin, 2011 <sup>14</sup>      | 162       | 324             | 54 (33.3)   | 16 (29.6)      | 15 (27.8)      | 23 (42.6)         | -               |
| Tomovic, 2012 <sup>12</sup>   | 170       | 340             | 117 (68.8)  | 45 (38.5)      | 28 (23.9)      | 44 (37.6)         | -               |
| Ozturan, 2013 <sup>10</sup>   | 999       | 1998            | 160 (16.6)  | 58 (36.3)      | 50 (31.2)      | 52 (32.5)         | 80.1%           |
| Wada, 2015 <sup>18</sup>      | 261       | 522             | 132 (50.6)* | -              | -              | -                 | 34.8%           |
| Senturk, 2017 <sup>19</sup>   | 618       | 1236            | 326 (52.7)  | 94 (28.8)      | 78 (23.9)      | 154 (47.3)        | -               |
| Chmielik, 2017 <sup>4</sup>   | 196       | 392             | 78 (39.8)   | -              | -              | -                 | 25%             |
| Özdemir, 2019 <sup>2</sup>    | 508       | 1016            | 108 (21.2)  | 44 (40.7)      | 28 (25.9)      | 36 (33.4)         | 5.3%**          |
| Thimmaiah, 2017 <sup>8</sup>  | 1080      | 2160            | 260 (24.1)  | -              | -              | -                 | -               |
| Ali, 2020 <sup>3</sup>        | 201       | 402             | 86 (42.8)   | -              | -              | -                 | -               |
| Fadda, 2022 <sup>22</sup>     | 430       | 860             | 93 (21.6)   | 33 (35.5)      | 35 (37.6)      | 25 (26.9)         | 18.3%**         |

PNSCT: paranasal CT; OC: Onodi cell; ONC: optic nerve canal. \* sides; \*\* only Type IV bulging.

ful use of appropriate surgical instrumentation also play a key role in the successful management of potential iatrogenic injuries.

To minimise intra-operative injuries during sphenoid surgery in the presence of Type IV optic nerve into the OC, Fadda et al. <sup>22</sup> recommend using a neuroinvestigation system and to not use drills and surgical chisels to avoid injuries of vital structures, in particular in presence of protruded ICA or ONC.

In addition, Humphreys et al. <sup>23</sup> reports use of through-cutting forceps instead of traction-instruments to remove bony partitions to avoid possible damage of vital neurovascular structures, to preserve the mucosal layer, and to reduce the risk of neo-osteogenesis and scarring. Shaver dissection along the sphenoid area should be carefully used to minimise the risk of dural violation and consequent brain injury. Strength and level of wall suction should be well regulated. Lastly, use of a neuroinvestigation system is also recommended similar to Fadda et al. <sup>22</sup>.

## Conclusions

The identification of OC and its relationship with adjacent structures in pre-operative assessment represents a prerogative of ESS because of its proximity to important anatomical structures such as the ONC and ICA. Its identification is crucial to avoid potential complications during surgery <sup>11</sup>. Furthermore, the presence of an OC, if not previously detected, can also be a source of a more difficult approach to the sphenoid sinus for the surgeon, since it may cause the natural ostium of the sphenoid sinus to be located more superiorly, and the asymmetry between sides may steer the surgeon in the wrong direction while opening the sinus. This anatomic variant can also cause

difficulties in locating the skull base during trans-sphenoid exposure.

In the most challenging cases, given the risk by Type IV of ONC bulging into the OC, use of neuronavigation systems to assist the surgeon during ESS may be helpful to reduce the risk of complications; especially in cases where the widely enunciated risky variants of ONC may place the surgeon in a dangerous intra-operative setting, exposing the patient to potentially catastrophic consequences. Finally, in such cases, traumatic surgical tools such as drill or chisel should be avoided when approaching the OC if the ONC is bulging into it (particularly in cases of Type IV-ONC).

## Conflict of interest statement

The authors declare no conflict of interest.

## Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

## Author contributions

GLF: substantial contributions to the work reported in this manuscript and surgeon, final approval of the version to be published; AU: substantial contributions to the conception or design of the work, final approval of the version to be published; AP: substantial contribution to the statistical analysis and interpretation of the data, final approval of the version to be published; MT, EdC, GC: substantial contributions to the preparation of patients undergoing surgery, editing, revision and final approval of the version to be published; LN, AMS, MP, DG: substantial contributions to the analysis of the CT scans, final approval of the version to be published.

### Ethical consideration

This study was approved by the “San Luigi Gonzaga” University Hospital Ethics Committee (Date: 18.03.2022, protocol number: 4526). The research was conducted ethically, with all study procedures being performed in accordance with the requirements of the World Medical Association’s Declaration of Helsinki.

Written informed consent was obtained from each participant for study participation and data publication.

### References

- 1 Onodi A. The optic nerve and the accessory sinuses of the nose. *Der Sehnerv und die NebenOhlen der Nase*. Wien, Germany: Alfred Holder; 1907. pp. 1-6.
- 2 Özdemir A, Bayar Muluk N, Asal N, et al. Is there a relationship between Onodi cell and optic canal? *Eur Arch Otorhinolaryngol* 2019;276:1057-1064. <https://doi.org/10.1007/s00405-019-05284-0>
- 3 Ali IK, Sansare K, Karjodkar F, et al. Imaging analysis of Onodi cells on cone-beam computed tomography. *Int Arch Otorhinolaryngol* 2020;24:E319-E322. <https://doi.org/10.1055/s-0039-1698779>
- 4 Chmielik LP, Chmielik A. The prevalence of the Onodi cell – most suitable method of CT evaluation in its detection. *Int J Pediatr Otorhinolaryngol* 2017;97:202-205. <https://doi.org/10.1016/j.ijporl.2017.04.001>
- 5 Driben JS, Bolger WE, Robles HA, et al. The reliability of computerized tomographic detection of the Onodi (Sphenoethmoid) cell. *Am J Rhinol* 1998;12:105-111. <https://doi.org/10.2500/105065898781390325>
- 6 Bansberg SF, Harner SG, Forbes G. Relationship of the optic nerve to the paranasal sinuses as shown by computed tomography. *Otolaryngol Head Neck Surg* 1987;96:331-335. <https://doi.org/10.1177/019459988709600405>
- 7 Batra PS, Citardi MJ, Gallivan RP, et al. Software-enabled CT analysis of optic nerve position and paranasal sinus pneumatization patterns. *Otolaryngol Head Neck Surg* 2004;131:940-945. <https://doi.org/10.1016/j.otohns.2004.07.013>
- 8 Thimmaiah VT, Anupama C. Pneumatization patterns of Onodi cell on multidetector computed tomography. *J Oral Maxillofac Radiol* 2017;5:63-66.
- 9 Güldner C, Pistorius SM, Diogo I, et al. Analysis of pneumatization and neurovascular structures of the sphenoid sinus using cone-beam tomography (CBT). *Acta Radiol* 2012;53:214-219. <https://doi.org/10.1258/ar.2011.110381>
- 10 Ozturan O, Yenigun A, Degirmenci N, et al. Co-existence of the Onodi cell with the variation of perisphenoidal structures. *Eur Arch Otorhinolaryngol* 2013;270:2057-2063. <https://doi.org/10.1007/s00405-012-2325-8>
- 11 Kim JY, Kim HJ, Kim CH, et al. Optic nerve injury secondary to endoscopic sinus surgery: an analysis of three cases. *Yonsei Med J* 2005;46:300-304. <https://doi.org/10.3349/ymj.2005.46.2.300>
- 12 Tomovic S, Esmaeili A, Chan NJ, et al. High-resolution computed tomography analysis of the prevalence of Onodi cells. *Laryngoscope* 2012;122:1470-1473. <https://doi.org/10.1002/lary.23346>
- 13 Kantarci M, Karasen RM, Alper F, et al. Remarkable anatomic variations in paranasal sinus region and their clinical importance. *Eur J Radiol* 2004;50:296-302. <https://doi.org/10.1016/j.ejrad.2003.08.012>
- 14 Shin JH, Kim SW, Hong YK, et al. The Onodi cell: an obstacle to sellar lesions with a transsphenoidal approach. *Otolaryngol Head Neck Surg* 2011;145:1040-1042. <https://doi.org/10.1177/0194599811418040>
- 15 Nitinavakarn B, Thanaviratananich S, Sangslip N. Anatomical variations of the lateral nasal wall and paranasal sinuses: a CT study for endoscopic sinus surgery (ESS) in Thai patients. *J Med Assoc Thai* 2005;88:763-768.
- 16 Kasemsiri P, Thanaviratananich S, Puttharak W. The prevalence and pattern of pneumatization of Onodi cell in Thai patients. *J Med Assoc Thai* 2011;94:1122-1126.
- 17 Thanaviratananich S, Chaisiwamongkol K, Kraitrakul S, et al. The prevalence of an Onodi cell in adult Thai cadavers. *Ear Nose Throat J* 2003;82:200-204.
- 18 Wada K, Moriyama H, Edamatsu H, et al. Identification of Onodi cell and new classification of sphenoid sinus for endoscopic sinus surgery. *Int Forum Allergy Rhinol* 2015;5:1068-1076. <https://doi.org/10.1002/alr.21567>
- 19 Senturk M, Guler I, Azgin I, et al. The role of Onodi cells in sphenoiditis: results of multiplanar reconstruction of computed tomography scanning. *Braz J Otorhinolaryngol* 2017;83:88-93. <https://doi.org/10.1016/j.bjorl.2016.01.011>
- 20 Yeoh KH, Tan KK. The optic nerve in the posterior ethmoid in Asians. *Acta Otolaryngol* 1994;114:329-336. <https://doi.org/10.3109/00016489409126065>
- 21 Dessi P, Moulin G, Castro F, et al. Protrusion of the optic nerve into the ethmoid and sphenoid sinus: prospective study of 150 CT studies. *Neuroradiology* 1994;36:515-516. <https://doi.org/10.1007/BF00593511>
- 22 Fadda GL, Petrelli A, Urbanelli A, et al. Risky anatomical variations of sphenoid sinus and surrounding structures in endoscopic sinus surgery. *Head Face Med* 2022;18:29. <https://doi.org/10.1186/s13005-022-00336-z>
- 23 Humphreys IM, Hwang PH. Avoiding complications in endoscopic sinus surgery. *Otolaryngol Clin North Am* 2015;48:871-881. <https://doi.org/10.1016/j.otc.2015.05.013>