

Weakly weighted generalised quasi-metric spaces and semilattices

Original

Weakly weighted generalised quasi-metric spaces and semilattices / Castellano, I., Giordano Bruno, A., Zava, N.. - In: THEORETICAL COMPUTER SCIENCE. - ISSN 0304-3975. - 977:(2023), pp. 1-29. [10.1016/j.tcs.2023.114129]

Availability:

This version is available at: 11583/3003346 since: 2025-09-25T13:20:58Z

Publisher:

Elsevier

Published

DOI:10.1016/j.tcs.2023.114129

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

Elsevier postprint/Author's Accepted Manuscript

© 2023. This manuscript version is made available under the CC-BY-NC-ND 4.0 license
<http://creativecommons.org/licenses/by-nc-nd/4.0/>. The final authenticated version is available online at:
<http://dx.doi.org/10.1016/j.tcs.2023.114129>

(Article begins on next page)

Weakly weighted generalised quasi-metric spaces and semilattices

Ilaria Castellano^a, Anna Giordano Bruno^b, Nicolò Zava^c

^a*Fakultät für Mathematik, Universität Bielefeld, D-33501, Bielefeld, Germany*

^b*Dipartimento di Scienze Matematiche, Informatiche e Fisiche, Università degli Studi di Udine, 33100, Udine, Italy*

^c*Institute of Science and Technology Austria (ISTA), 3400, Klosterneuburg, Austria*

Abstract

Motivated by recent applications to entropy theory in dynamical systems, we generalise notions introduced by Matthews and define weakly weighted and componentwise weakly weighted (generalised) quasi-metrics. We then systematise and extend to full generality the correspondences between these objects and other structures arising in theoretical computer science and dynamics. In particular, we study the correspondences with weak partial metrics and, if the underlying space is a semilattice, with invariant (generalised) quasi-metrics satisfying the descending path condition, and with strictly monotone semi(-co-)valuations.

We conclude discussing, for endomorphisms of generalised quasi-metric semilattices, a generalisation of both the known intrinsic semilattice entropy and the semigroup entropy.

Keywords: generalised quasi-metric, weak partial metric, weak weight, weakly weighted quasi-metric, quasi-metric semilattice, semivaluation, intrinsic entropy.

2020 MSC: 54E35, 06A12, 06B35, 20M10, 54C70, 54E15.

1. Introduction

The notion of metric space has been generalised in several directions with the aim of producing non-Hausdorff topologies. For instance, Scott's breakthrough works [43, 44] used non-Hausdorff topologies to describe partial objects in computation (see also the survey [1] and the monograph [27]). Moreover, partial metrics were introduced by Matthews ([34]) and allow the distance from a point to itself to be non-zero. On the other hand, quasi-metrics abandon the symmetry axiom; they were introduced and studied for the first time by Wilson ([51]), but this notion can be traced back to Hausdorff ([28]).

Matthews introduced the notion of a *weighted quasi-metric*, that is a quasi-metric d on a non-empty set X such that there exists a function $w: X \rightarrow \mathbb{R}_{\geq 0}$ with

$$d(x, y) + w(x) = d(y, x) + w(y), \quad \text{for every } x, y \in X. \quad (1.1)$$

Indeed, he showed a precise correspondence between partial metrics and weighted quasi-metrics, where the specialisation orders of the partial metric and the corresponding weighted quasi-metric coincide:

$$\text{partial metrics} \xleftarrow{[34]} \text{weighted quasi-metrics}. \quad (1.2)$$

As immediately noticed by Matthews, not every quasi-metric is weighted, and moreover the question he posed of which quasi-metrizable spaces admit weighted quasi-metrics is still open – see the paper [31] by Künzi and Vajner on topological aspects of weighted quasi-metrics.

The complexity spaces were introduced in [39] as a part of the development of a topological foundation for the complexity analysis of algorithms, while the dual complexity spaces were investigated in [36, 32] as an appropriate tool to carry out a mathematical analysis of complexity spaces. Schellekens ([42]) noticed that these and several other examples of partial metric spaces arising in quantitative domain theory share a further underlying structure: the specialisation order associated to the weighted quasi-metric (equivalently, to the partial metric) turns them into semilattices. In this case we speak about *quasi-metric semilattices* with a slight difference with the terminology used in [42], which adopts the classical one for quasi-metric lattices from [48, 49] (see Definition 4.9 and the comment after it).

Inspired by the classical connection between (strictly increasing) valuations and (pseudo-)metrics on a lattice (see Birkhoff's monograph [5]), a correspondence between partial metrics and valuations on semilattices was studied by O'Neill ([33]) and further discussed in [7, 8].

As previously done by Nakamura ([35]), Schellekens ([42]) introduced semi(-co-)valuations on semilattices, showing how these notions precisely extend the classical concept of valuations for lattices. Moreover, he provided a correspondence between

Email addresses: ilaria.castellano@math.uni-bielefeld.de (Ilaria Castellano), anna.giordanobruno@uniud.it (Anna Giordano Bruno), nicolo.zava@gmail.com (Nicolò Zava)

invariant (see Definition 4.9) (co-)weighted quasi-metrics and strictly increasing semi(-co-)valuations ([42]), and a characterisation of invariant quasi-metrics on semilattices as those satisfying the *descending path condition* (briefly, (DPC) – see Definition 4.6) and the extension property ([41]):

$$\begin{array}{ccccc}
\text{invariant} & \xleftrightarrow{[34]} & \text{invariant weighted} & \xleftrightarrow{[41]} & \text{invariant quasi-metrics with} \\
\text{partial metrics} & & \text{quasi-metrics} & & \text{(DPC) and extension property} \\
& & \updownarrow [42] & & \\
& & \text{strictly monotone} & & \\
& & \text{positive semi(-co-)valuations.} & &
\end{array} \tag{1.3}$$

The main aim of this paper is to complete, extend and systematise the above relations by Matthews and Schellekens. In order to achieve this, inspired by the notion of *weak partial metrics* due to O’Neill ([33], see Definition 3.1), we introduce *weakly weighted quasi-metrics* d on a space X by allowing the function $w: X \rightarrow \mathbb{R}$, which satisfies (1.1), to take any real value (see Definition 2.3). This new notion immediately allows us to extend the correspondence in (1.2) to weak partial metrics and weakly weighted quasi-metrics:

$$\text{weak partial metrics} \xleftrightarrow{\text{Cor. 3.3}} \text{weakly weighted quasi-metrics.} \tag{1.4}$$

Then, in the spirit of Schellekens’ work, we extend his results for semilattices collected in (1.3). In particular, we characterise the invariant quasi-metric semilattices that are weakly weighted as those satisfying (DPC), and prove the existence of a correspondence between weak weights for invariant quasi-metrics and strictly monotone semi(-co-)valuations for a semilattice:

$$\begin{array}{ccccc}
\text{invariant weak} & \xleftrightarrow{\text{Rem. 4.17}} & \text{invariant weakly} & \xleftrightarrow{\text{Cor. 5.5}} & \text{invariant quasi-metrics} \\
\text{partial metrics} & & \text{weighted quasi-metrics} & & \text{with (DPC)} \\
& & \updownarrow \text{Cor. 5.17} & & \\
& & \text{strictly monotone} & & \\
& & \text{semi(-co-)valuations.} & &
\end{array} \tag{1.5}$$

Furthermore, our extensions allow for alternative proofs of the classical correspondences represented in (1.3).

A quasi-metric d on a space X is a *generalised quasi-metric* if it can also assume the value ∞ . Generalised quasi-metric semilattices play a central role in the theory of intrinsic entropy. Indeed, the first attempt at a lattice-theoretic approach to entropy is due to Nakamura, who proved in [35] that Shannon’s entropy from information theory ([45]) can be characterised as a semivaluation on a semilattice. Then, a notion of *semigroup entropy* for normed semigroups and their monotone endomorphisms was proposed in [15] in order to find a general scheme for (almost) all classical entropies in mathematics; in most of the specific cases the underlying semigroup is a semilattice. Finally, in [11] a notion of *intrinsic semilattice entropy* was introduced in the category of generalised quasi-metric semilattices and non-expansive homomorphisms as the key to a unifying approach able to cover all (or, at least, most) of the intrinsic-like entropies in literature. Since all of the examples of generalised quasi-metric semilattices appearing in [11] satisfy the property (DPC), it is natural to ask whether we could retrieve a scheme of correspondences similar to (1.5) in this more general context.

To this end, first we introduce *generalised weak partial metrics* and further extend the correspondence in (1.4) to generalised weak partial metrics and weakly weighted generalised quasi-metrics:

$$\text{generalised weak partial metrics} \xleftrightarrow{\text{Cor. 3.8}} \text{weakly weighted generalised quasi-metrics.}$$

We then investigate possible extensions of (1.5) to the generalised setting. It is clear that weakly weighted generalised quasi-metrics and generalised weak partial metrics are not the right tool to characterise generalised quasi-metric semilattices with (DPC). In fact, a weakly weighted generalised quasi-metric space is actually a quasi-metric space provided that its specialisation order induces a semilattice (see Remark 5.1). A more suitable alternative descends from the following observation. The absorbing element ∞ forces the generalised quasi-metric space (X, d) to split in large-scale connected components $\mathcal{Q}(x)$ (see Definition 2.2), with $x \in X$, and so it induces an equivalence relation \cong_d on X . Each $\mathcal{Q}(x)$ is a quasi-metric space with the quasi-metric $d \upharpoonright_{\mathcal{Q}(x)}$ given by the restriction of d . We then call (X, d) *componentwise weakly weighted* when there exists a function $w: X \rightarrow \mathbb{R}$ such that, for every $x \in X$, $w \upharpoonright_{\mathcal{Q}(x)}$ is a weak weight for $d \upharpoonright_{\mathcal{Q}(x)}$ (see Definition 2.19).

Moreover, we introduce the notion of *generalised semi(-co-)valuation* (see Definition 6.4) for generalised quasi-metric semilattices, and extend the correspondences in (1.5) to the most general framework:

$$\begin{array}{ccccc}
\text{invariant generalised} & \xleftrightarrow{\text{Th. 6.2}} & \text{invariant componentwise} & \xleftrightarrow{\text{Cor. 6.11}} & \text{strictly monotone} \\
\text{quasi-metrics} & & \text{weakly weighted} & & \text{generalised} \\
\text{satisfying (DPC)} & & \text{generalised quasi-metrics} & & \text{semi(-co-)valuations.}
\end{array}$$

At the end of this paper, as an application of the above listed results we introduce a new notion of entropy. For an endomorphism ϕ of a semilattice X with an equivalence relation \cong , first we introduce the concept of (ϕ, \cong) -inert element, which extends that of ϕ -inert element for a non-expansive endomorphism of a generalised quasi-metric semilattice introduced in [11]. Then, using (ϕ, \cong) -inert elements, we propose a new intrinsic entropy for ϕ with respect to \cong . This entropy generalises both the semigroup entropy for normed semilattices from [15] and the intrinsic semilattice entropy from [11].

The paper is organised as follows.

Section 2 collects the main definitions, some basic properties of (componentwise) weakly weighted (generalised) quasi-metric spaces and provides several examples. Those coming from entropy theory will be exploited throughout the entire paper. In particular, §2.1 treats the weakly weighted quasi-metric spaces and the comparison to the standard notions of (co-)weighted quasi-metric spaces, §2.2 concerns weakly weighted generalised quasi-metric spaces, while §2.3 the even more general notion of componentwise weakly weighted generalised quasi-metric spaces.

Section 3 is dedicated to partial metrics. In §3.1 we recall the notion of a weak partial metric and provide the above mentioned correspondence between weakly weighted quasi-metrics and weak partial metrics. In §3.2 this correspondence is extended to weakly weighted generalised quasi-metrics and generalised weak partial metrics.

In Section 4, we turn our attention to generalised quasi-metric semilattices: in §4.1 we analyse the properties of the specialization order and (DPC), while §4.2 concerns the invariance property.

Section 5 is dedicated to the characterization of weakly weighted invariant quasi-metric semilattices as those that satisfy (DPC) (§5.1) and to the correspondence between weakly weighted invariant quasi-metrics and strictly monotone semi(-co-)valuations (§5.2).

In Section 6, we extend the results from the previous section to generalised quasi-metric semilattices, by defining suitable notions of generalised semi(-co-)valuations.

In Section 7, we introduce the new intrinsic entropy for an endomorphism ϕ of a meet-semilattice X with an equivalence relation \cong , and we recover the semigroup entropy (§7.1) and the intrinsic semilattice entropy (§7.2) as particular cases.

Notation. We denote by \mathbb{R} and \mathbb{N} the real and natural numbers (including 0), respectively. Moreover, we set, for a real constant c , $\mathbb{R}_{\geq c} = \{x \in \mathbb{R} \mid x \geq c\}$ and $\mathbb{R}_{> c} = \{x \in \mathbb{R} \mid x > c\}$. Finally, $\mathbb{N}_+ = \mathbb{N} \setminus \{0\}$. As usual, for a set X , we denote its cardinality by $|X|$.

Acknowledgement. We thank the referees for their comments that helped us improve our manuscript.

2. Weakly weighted (generalised) quasi-metric spaces

Definition 2.1. A *generalised quasi-metric* on a non-empty set X is a function $d: X \times X \rightarrow \mathbb{R}_{\geq 0} \cup \{\infty\}$ satisfying the properties:

(QM1) for $x, y \in X$, $d(x, y) = d(y, x) = 0$ if and only if $x = y$;

(QM2) $d(x, z) \leq d(x, y) + d(y, z)$, for every $x, y, z \in X$, with $r < r + \infty = \infty + \infty = \infty$ for every $r \in \mathbb{R}_{\geq 0}$.

Whenever the function d admits only finite values, i.e., $d: X \times X \rightarrow \mathbb{R}_{\geq 0}$, we simply call it *quasi-metric*.

The pair (X, d) is a (*generalised*) *quasi-metric space* provided that d is a (generalised) quasi-metric on X .

A (generalised) quasi-metric d on a set X is said to be a (*generalised*) *metric* if it satisfies the following axiom:

(QM3) d is *symmetric* (i.e., $d(x, y) = d(y, x)$ for every $x, y \in X$).

In this case, the pair (X, d) is a (*generalised*) *metric space*.

Given a (generalised) quasi-metric d on a set X , we define the *conjugated (generalised) quasi-metric* d^{-1} by

$$d^{-1}(x, y) := d(y, x), \quad \text{for every } x, y \in X, \quad (2.1)$$

and the *symmetrisation* d^s of d , which is a (generalised) metric, by the law

$$d^s(x, y) := \max\{d(x, y), d^{-1}(x, y)\} = \max\{d(x, y), d(y, x)\}, \quad \text{for every } x, y \in X. \quad (2.2)$$

If (X, d) is a generalised quasi-metric space and Y is a non-empty subset of X , then Y becomes a generalised quasi-metric space with the restriction of d to Y , that we denote by $d \upharpoonright_Y$ with some abuse of notation.

Definition 2.2. Let (X, d) be a generalised quasi-metric space. We define the equivalence relation \cong_d on X as follows:

$$x \cong_d y \text{ if } d^s(x, y) < \infty, \quad \text{for every } x, y \in X.$$

For every $x \in X$, the *large-scale connected component* of x is the equivalence class $[x]_{\cong_d}$ of x , which we denote by $\mathcal{Q}_X(x)$ for consistency with [52].

If there is no risk of ambiguity, we simply refer to $\mathcal{Q}_X(x)$ as $\mathcal{Q}(x)$. The family of large-scale connected components provides a partition of X with the following further property: for every $x \in X$, $(\mathcal{Q}(x), d \upharpoonright_{\mathcal{Q}(x)})$ is a quasi-metric space.

Notice that quasi-metric spaces can be regarded as generalised quasi-metric spaces with only one large-scale connected component.

A map $f: (X, d_X) \rightarrow (Y, d_Y)$ between (generalised) quasi-metric spaces is *non-expansive* (or *1-Lipschitz*) if, for every $x, y \in X$, $d_Y(f(x), f(y)) \leq d_X(x, y)$.

2.1. Weakly weighted quasi-metric spaces

We start with one of the central notions of the paper.

Definition 2.3. Let (X, d) be a generalised quasi-metric space. Then (X, d) and d are called *weakly weighted* if there exists a function $w: X \rightarrow \mathbb{R}$, called *weak weight* for d , such that, for every $x, y \in X$,

$$d(x, y) + w(x) = d(y, x) + w(y). \quad (2.3)$$

We also say that d is weakly weighted by w .

Clearly, every generalised metric is weakly weighted. In fact, (2.3) is fulfilled if we take w to be the constant function 0 (or, in general, any constant function). Moreover, a generalised quasi-metric is weakly weighted by a function w if and only if its conjugate is weakly weighted by $-w$.

The definition above is inspired by the following concepts of Matthews ([34]) and Schellekens ([42]), respectively. A quasi-metric space (X, d) and its quasi-metric d are said to be:

- *weighted* if there exists a function $w: X \rightarrow \mathbb{R}_{\geq 0}$, called *weight*, satisfying (2.3) for every pair $x, y \in X$;
- *co-weighted* if there exists a function $w: X \rightarrow \mathbb{R}_{\geq 0}$, called *co-weight*, such that, for every $x, y \in X$,

$$d(x, y) + w(y) = d(y, x) + w(x); \quad (2.4)$$

equivalently, the conjugated quasi-metric d^{-1} is weighted by w .

In this subsection, we first focus on weakly weighted quasi-metric spaces, motivated by the connection with weighted and co-weighted quasi-metrics that we formalise in Remark 2.4. We leave to the subsequent subsection considerations regarding weakly weighted generalised quasi-metric spaces.

Remark 2.4. Let (X, d) be a quasi-metric space. If $w: X \rightarrow \mathbb{R}_{\geq 0}$ is a weight for d , then trivially w is a weak weight for d , and thus (X, d) is weakly weighted. Similarly, if $w: X \rightarrow \mathbb{R}_{\geq 0}$ is a co-weight for d , then $-w$ satisfies (2.3) and so it is a weak weight for (X, d) .

Hence, Definition 2.3 generalises both weighted and co-weighted quasi-metric spaces. We will see in Example 2.10(c) that there are weakly weighted quasi-metric spaces that are neither weighted nor co-weighted.

Proposition 2.5. *Let (X, d) be a (weakly/co-) weighted quasi-metric space, $w: X \rightarrow \mathbb{R}$ be a (weak/co-) weight, and $Y \subseteq X$. Then $(Y, d \upharpoonright_Y)$ is (weakly/co-) weighted and $w \upharpoonright_Y$ is a (weak/co-) weight.*

Weak weights, weights and co-weights are not unique, as stated in the following proposition. On the other hand, the result shows that two weak weights for the same quasi-metric differ only by a constant. For the sake of brevity, if S is a non-empty set, $f: S \rightarrow \mathbb{R}$ a function and $c \in \mathbb{R}$, we denote by $f + c$ the function from S to \mathbb{R} defined by the law $(f + c)(x) = f(x) + c$, for every $x \in S$. Moreover, we write $f \geq 0$ ($f \leq 0$) if, for every $x \in S$, $f(x) \geq 0$ ($f(x) \leq 0$, respectively).

Proposition 2.6. *Let (X, d) be a quasi-metric space and $w, w': X \rightarrow \mathbb{R}$.*

- (a) *Suppose that w is a weak weight for d . Then w' is a weak weight for d if and only if $w' = w + c$ for some $c \in \mathbb{R}$.*
- (b) ([42, Proposition 8]) *Suppose that w is a (co-)weight for d . Then w' is a (co-)weight for d if and only if $w' = w + c$ for some $c \in \mathbb{R}$ and $w' \geq 0$.*

Proof. Let us prove item (a), while item (b) can be similarly derived. Suppose then that w is a weak weight for d . It is easy to see that for every $c \in \mathbb{R}$ the function $w + c$ still satisfies (2.3).

Suppose now that w' is another weak weight for d . Then, for every $x, y \in X$,

$$\begin{aligned} w(x) - w'(x) &= d(x, y) + w(x) - (d(x, y) + w'(x)) = \\ &= d(y, x) + w(y) - (d(y, x) + w'(y)) = w(y) - w'(y), \end{aligned}$$

which is obtained using (2.3) for both w and w' . Hence w and w' differ by a constant, and so the claim follows. \square

Let us introduce an equivalence relation \sim on \mathbb{R}^X , which is the family of all functions from a non-empty set X to \mathbb{R} , by putting,

$$\text{for every } f, g \in \mathbb{R}^X, f \sim g \text{ if there exists a constant } c \in \mathbb{R} \text{ such that } g = f + c. \quad (2.5)$$

Then Proposition 2.6(a) and (b) can be rewritten respectively in the following way:

- (a) if w is a weak weight for d , then $[w]_{\sim}$ is the family of all weak weights for d ;
- (b) if w is a (co-)weight for d , then $[w]_{\sim} \cap \mathbb{R}_{\geq 0}^X$ is the family of all (co-)weights for d .

Using Proposition 2.6, we can give a useful characterisation of those weakly weighted quasi-metric spaces that are (co-)weighted.

Proposition 2.7. *For a weakly weighted quasi-metric space (X, d) the following equivalences hold:*

- (a) *(X, d) is weighted if and only if some (equivalently, every) weak weight $w: X \rightarrow \mathbb{R}$ for d has a lower bound \underline{b}_w , i.e., $w(X) \subseteq [\underline{b}_w, +\infty[$.*

Moreover, given a weak weight w , the weak weight $w - c \in [w]_{\sim}$ is a weight for every $c \leq \underline{b}_w$;

(b) (X, d) is co-weighted if and only if some (equivalently, every) weak weight $w: X \rightarrow \mathbb{R}$ for d has an upper bound \bar{b}_w , i.e., $w(X) \subseteq]-\infty, \bar{b}_w]$.

Moreover, given a weak weight w for d , the weak weight $-w + c \in [-w]_{\sim}$ is a co-weight for d for every $c \geq \bar{b}_w$.

Proof. Let us prove item (a), while the dual item (b) can be similarly shown.

If (X, d) is weighted, then there is a weak weight w for d with non-negative values (see Proposition 2.6(b)) and so $\underline{b}_w = 0$ is a lower bound. By Proposition 2.6(a), if a particular weak weight for d is bounded from below, then every weak weight for d has a lower bound.

Conversely, suppose that there is a weak weight w for d bounded from below by a constant \underline{b}_w . If $\underline{b}_w \geq 0$, then w is a weight for d by definition and (X, d) is weighted. Otherwise, we can use $w - c \in [w]_{\sim}$ to weight (X, d) whenever $c \leq \underline{b}_w$. \square

Remark 2.8. Let (X, d) be a quasi-metric space. In [42], a function $f: X \rightarrow \mathbb{R}$ such that $\inf_{x \in X} f(x) = 0$ is called *fading* and [42, Proposition 8] (see Proposition 2.6(b)) shows that, if w is a (co-)weight for d , then there exists a unique fading (co-)weight for d . Therefore, we can say that a weakly weighted quasi-metric space (X, d) is weighted if and only if the class of all its weak weights $[w]_{\sim}$ contains a fading element. In particular, the fading representative can be explicitly obtained as follows: if w is a weak weight for d , the function $w' = w - \inf_{x \in X} w(x)$ has the desired property.

A dual observation holds for co-weights.

Before presenting some examples, we state a technical property of weakly weighted quasi-metric spaces that will be useful in the sequel:

Lemma 2.9. Let (X, d) be a generalised quasi-metric space that is weakly weighted by w . For every $x, y, z \in X$ such that $d(x, z) = d(x, y) + d(y, z)$, one has $d(z, x) = d(z, y) + d(y, x)$.

Proof. Let $x, y, z \in X$ be such that $d(x, z) = d(x, y) + d(y, z)$. Then (2.3) yields

$$\begin{aligned} d(z, x) &= d(z, x) + w(z) - w(z) = d(x, z) + w(x) - w(z) = \\ &= d(x, y) + d(y, z) + w(x) - w(z) = d(y, x) + w(y) + d(y, z) - w(z) = \\ &= d(y, x) + d(z, y) + w(z) - w(z) = d(y, x) + d(z, y). \end{aligned} \quad \square$$

Below we provide several examples of weak weights. In particular, Example 2.10(c) shows that Proposition 2.7 makes it is easier to produce weakly weighted quasi-metric spaces that are neither weighted nor co-weighted. Moreover, it exposes a method to produce (co-)weights: it suffices to bound the weak weight involved by restricting the domains.

Example 2.10. (a) Let $\mathbb{S} = \{0, 1\}$ be a two-point set endowed with the quasi-metric $d_{\mathbb{S}}$ defined as follows: $d_{\mathbb{S}}(0, 1) = 0$ and $d_{\mathbb{S}}(1, 0) = 1$. The topological space $(\mathbb{S}, d_{\mathbb{S}})$ is also known as *Sierpiński space*. Then $d_{\mathbb{S}}$ is both weighted and co-weighted.

Take, for example, the weight $w(0) = 1$ and $w(1) = 0$ and the co-weight $w'(0) = 0$ and $w'(1) = 1$.

(b) On the three point set $\mathbb{T} = \{0, 1, 2\}$, define the quasi-metric $d_{\mathbb{T}}$ as follows:

$$d_{\mathbb{T}}(0, 1) = d_{\mathbb{T}}(0, 2) = d_{\mathbb{T}}(1, 2) = 0, \quad d_{\mathbb{T}}(2, 1) = d_{\mathbb{T}}(1, 0) = 1 \text{ and } d_{\mathbb{T}}(2, 0) = 2.$$

Then $d_{\mathbb{T}}$ is both weighted and co-weighted. We can indeed define a weight $w(0) = 2$, $w(1) = 1$ and $w(2) = 0$, and a co-weight $w'(0) = 0$, $w'(1) = 1$ and $w'(2) = 2$.

(c) Let us endow \mathbb{R} with the quasi-metric d defined as follows:

$$d_{\mathbb{R}}(x, y) = \max\{x - y, 0\}.$$

Then $d_{\mathbb{R}}$ is weakly weighted by the weak weight w defined by $w(x) = -x$, for every $x \in X$. By virtue of Proposition 2.7, since w is neither lower nor upper bounded, $(\mathbb{R}, d_{\mathbb{R}})$ is neither weighted nor co-weighted. Again Proposition 2.7 implies that $(\mathbb{R}_{\geq 0}, d_{\mathbb{R}} \upharpoonright_{\mathbb{R}_{\geq 0}})$ is co-weighted and $(\mathbb{R}_{\leq 0}, d_{\mathbb{R}} \upharpoonright_{\mathbb{R}_{\leq 0}})$ is weighted.

Notice that $d_{\mathbb{S}}$ and $d_{\mathbb{T}}$ coincide with the quasi-metric induced by $d_{\mathbb{R}}$ on the subsets $\{0, 1\}$ and $\{0, 1, 2\}$, respectively.

Next we see that not all quasi-metrics are weakly weighted. Please mind that the quasi-metrics of Examples 2.11 and 2.17 are different from $d_{\mathbb{T}}$ in Example 2.10(b), although they are defined on the same three-point set.

Example 2.11. On the three-point set $\mathbb{T} = \{0, 1, 2\}$, define a quasi-metric d as follows:

$$d(0, 1) = d(0, 2) = d(1, 2) = 0, \quad \text{and } d(2, 1) = d(2, 0) = d(1, 0) = 1.$$

Then d is not weakly weighted. Suppose by contradiction that a weak weight w for d exists. Using (2.3), we get

$$\begin{aligned} w(0) &= d(0, 2) + w(0) = d(2, 0) + w(2) = 1 + w(2) \quad \text{and} \\ w(1) &= d(1, 2) + w(1) = d(2, 1) + w(2) = 1 + w(2), \end{aligned}$$

so $w(0) = w(1)$. This is a contradiction, as

$$w(0) = d(0, 1) + w(0) = d(1, 0) + w(1) = 1 + w(1).$$

The following examples come from entropy theory and will be exploited throughout the entire paper.

Example 2.12. (a) Let S be a finite set. On the power set $\mathcal{P}(S)$, we define the quasi-metric $d_{\mathcal{P}(S)}$ as follows:

$$d_{\mathcal{P}(S)}(A, B) := |B \setminus A| = |(A \cup B) \setminus A|, \quad \text{for every } A, B \in \mathcal{P}(S). \quad (2.6)$$

Then $d_{\mathcal{P}(S)}$ is weighted by $w: \mathcal{P}(S) \rightarrow \mathbb{R}_{\geq 0}$ defined by $w(A) = |A|$, for every $A \subseteq S$. Moreover, a co-weight for $d_{\mathcal{P}(S)}$ is $w': \mathcal{P}(S) \rightarrow \mathbb{R}_{\geq 0}$ defined by $w'(A) = |X| - |A|$ for every $A \in \mathcal{P}(S)$.

(b) Let G be a finite abelian group and $L(G)$ be the family of all subgroups of G . We define on $L(G)$ the quasi-metric $d_{L(G)}$ as follows:

$$d_{L(G)}(H, K) := \log|H + K : H|, \quad \text{for every } H, K \in L(G). \quad (2.7)$$

Then $d_{L(G)}$ is weighted by $w: L(G) \rightarrow \mathbb{R}_{\geq 0}$ defined by $w(H) = \log|H|$, for every $H \in L(G)$. Moreover, a co-weight for $d_{L(G)}$ is $w': L(G) \rightarrow \mathbb{R}_{\geq 0}$ defined by $w'(H) = \log|G| - \log|H|$.

The quasi-metric spaces defined in the above example will be generalised in Example 2.21.

2.2. Weakly weighted generalised quasi-metric spaces

Let us focus on generalised quasi-metric spaces that are weakly weighted.

Proposition 2.13. *Let (X, d) be a generalised quasi-metric space. Then d is weakly weighted if and only if the following two properties hold:*

- (a) for every $x \in X$, $d \upharpoonright_{\mathcal{Q}(x)}$ is weakly weighted;
- (b) for every $x, y \in X$, $d(x, y) = \infty$ if and only if $d(y, x) = \infty$.

Proof. Let $\{X_i\}_{i \in I}$ be the family of large-scale connected components of X and denote $d \upharpoonright_{X_i}$ by d_i . If d is weakly weighted by w , then every d_i is weakly weighted (by $w \upharpoonright_{X_i}$) and (2.3) implies property (b) as w assumes only finite values.

Conversely, suppose that (a) and (b) are fulfilled. Define, for every $x \in X$, $w(x) = w_i(x)$, where $x \in X_i$ and w_i is a weak weight for d_i . Then w trivially satisfies (2.3), for every $x, y \in X$ with $x \cong_d y$. Moreover, condition (b) implies that w satisfies the desired property also for points $x, y \in X$ with $x \not\cong_d y$. \square

To discuss a corollary of Proposition 2.13, let us describe the coproducts of the category \mathbf{QM}_{Met} of generalised quasi-metric spaces and non-expansive maps between them.

Remark 2.14. Let $\{(X_i, d_i)\}_{i \in I}$ be a family of generalised quasi-metric spaces. Their coproduct in \mathbf{QM}_{Met} is given by the disjoint union of the underlying sets $X = \bigsqcup_{i \in I} X_i$ together with the generalised quasi-metric d defined as follows: if, for every $i \in I$, $j_i: X_i \rightarrow X$ represents the canonical inclusion of X_i into X , then

$$d(j_i(x), j_k(y)) = \begin{cases} d_i(x, y) & \text{if } i = k, \\ \infty & \text{otherwise,} \end{cases}$$

for every $j_i(x), j_k(y) \in X$. Indeed, it can be proved that (X, d) satisfies the universal property of coproducts.

If each d_i is weakly weighted by w_i , then d is weakly weighted by the function w making all the triangles in the following diagram commute:

$$\begin{array}{ccccc} & & X & & \\ & & \uparrow & & \uparrow \\ & & \mathbb{R} & & \mathbb{R} \\ & & \downarrow & & \downarrow \\ \cdots & X_i & \cdots & X_k & \cdots \\ & \uparrow & & \uparrow & \\ & j_i & & j_k & \\ & \uparrow & & \uparrow & \\ & w_i & & w_k & \end{array}$$

Thanks to the previous remark, Proposition 2.13 immediately implies the following characterisation.

Corollary 2.15. *A generalised quasi-metric space is weakly weighted if and only if it is the coproduct of a family of weakly weighted quasi-metric spaces.*

To conclude the subsection, let us focus on a specific example of generalised quasi-metric space. Given a directed graph $\Gamma = (V, E)$ and two vertices $x, y \in V$, a *directed path* P connecting x to y is a finite subset of edges $\{(z_{i-1}, z_i)\}_{i=1, \dots, n}$ such that $z_0 = x$ and $z_n = y$. We define the *path generalised quasi-metric* d_{Γ} on V as follows: for every $x, y \in V$,

$$d_{\Gamma}(x, y) := \begin{cases} \inf\{|P| \mid P \text{ is a directed path connecting } x \text{ to } y\} & \text{if } x \neq y, \\ 0 & \text{otherwise,} \end{cases}$$

with the usual notation that $\inf \emptyset = \infty$, and thus $d_{\Gamma}(x, y) = \infty$ in case there is no directed path from x to y in Γ . So, d_{Γ} is a generalised quasi-metric as there may be no directed path connecting a vertex to another one. More precisely, d_{Γ} is a quasi-metric if and only if Γ is *strongly connected*, i.e., for every pair of vertices x, y there is a path connecting x to y and a path connecting y to x .

Proposition 2.16. *Let $\Gamma = (V, E)$ be a strongly connected directed graph. Then the quasi-metric d_Γ is weakly weighted if and only if d_Γ is a metric, that is, Γ is a non-directed graph (i.e., $(x, y) \in E$ if and only if $(y, x) \in E$).*


Proof. Clearly, if d_Γ is a metric, then it is weakly weighted (a weak weight is any constant function). Moreover, d_Γ is a metric if and only if Γ is non-directed. In fact, for every distinct $x, y \in V$, $(x, y) \in E$ if and only if $d_\Gamma(x, y) = 1$ if and only if $d_\Gamma(y, x) = 1$, which is equivalent to $(y, x) \in E$.

Assume now that d_Γ is weakly weighted by w . Define

$$C = \{d_\Gamma(x, y) + d_\Gamma(y, x) \mid (x, y) \in V \times V : d(x, y) > d(y, x)\} \subseteq \mathbb{N}$$

(recall that d_Γ has integer values). Suppose, by contradiction, that C is non-empty. Then there exist $x, y \in V$ such that $d_\Gamma(x, y) > d_\Gamma(y, x)$ and $d_\Gamma(x, y) + d_\Gamma(y, x)$ is the minimum of C . Let P and Q be two directed paths going from x to y and from y to x , respectively, of minimum length. Note that $d_\Gamma(x, y) \geq 2$, and so we can take a point $z \notin \{x, y\}$ that is crossed by the path P . Hence, P is divided into two directed paths whose concatenation gives P : P_1 and P_2 going from x to z and from z to y , respectively. Note that $|P_1| = d_\Gamma(x, z)$ and $|P_2| = d_\Gamma(z, y)$ as, otherwise, we could provide a directed path going from x to y that is strictly shorter than P . Moreover, $d_\Gamma(x, y) = d_\Gamma(x, z) + d_\Gamma(z, y)$. By Lemma 2.9, one has $d_\Gamma(y, x) = d_\Gamma(y, z) + d_\Gamma(z, x)$. Then $d_\Gamma(x, z) + d_\Gamma(z, y) = d_\Gamma(x, y) > d_\Gamma(y, x) = d_\Gamma(y, z) + d_\Gamma(z, x)$ implies that either $d_\Gamma(x, z) > d_\Gamma(z, x)$ or $d_\Gamma(z, y) > d_\Gamma(y, z)$, and so a contradiction as $d_\Gamma(x, y) + d_\Gamma(y, x)$ was taken as minimal with that property. \square

Example 2.17. On \mathbb{T} we define the quasi-metric d' as the path quasi-metric associated with the following strongly connected directed graph:



Then d' is not weakly weighted by Proposition 2.16.

Thanks to Proposition 2.13, we can drop the strong connectivity requirement from the statement of Proposition 2.16.

Corollary 2.18. *Let $\Gamma = (V, E)$ be a directed graph. Then d_Γ is weakly weighted if and only if d_Γ is a metric, that is, Γ is a non-directed graph.*

2.3. Componentwisely weakly weighted generalised quasi-metric spaces

Proposition 2.13 and Corollary 2.18 suggest that the notion of weak weightedness is too restrictive in the realm of generalised quasi-metric spaces having more than one large-scale connected component. So, inspired by Proposition 2.13, we propose a more suitable version.

Definition 2.19. Let (X, d) be a generalised quasi-metric space. Then X is *componentwise weakly weighted* if there exists a function $w: X \rightarrow \mathbb{R}$, called *componentwise weak weight* for (X, d) (or, for d), such that (2.3) is fulfilled for every $x, y \in X$ satisfying $x \cong_d y$ (i.e., for every $x \in X$, $w \upharpoonright_{\mathcal{Q}(x)}$ is a weak weight for $d \upharpoonright_{\mathcal{Q}(x)}$).

Clearly, a weakly weighted generalised quasi-metric space is, in particular, componentwise weakly weighted, and the two notions coincide for quasi-metric spaces.

Fact 2.20. *A generalised quasi-metric space (X, d) is componentwise weakly weighted if and only if each large-scale connected component with the induced quasi-metric is weakly weighted (i.e., condition (a) of Proposition 2.13 holds). More precisely, $w: X \rightarrow \mathbb{R}$ is a componentwise weak weight of d if and only if, for every $x \in X$, $w \upharpoonright_{\mathcal{Q}(x)}$ is a weak weight of the restriction of d to $\mathcal{Q}(x)$.*

Proof. The “only if” implication is trivial. Conversely, the componentwise weak weight can be obtained by gluing together all the weak weights defined on the large-scale connected components. More explicitly, suppose that $\{(X_i, d_i)\}_{i \in I}$ is the partition of a generalised quasi-metric space (X, d) in its large-scale connected components endowed with the inherited quasi-metrics that are weakly weighted by the corresponding functions $w_i: X_i \rightarrow \mathbb{R}$. Then the function $w: X \rightarrow \mathbb{R}$ defined by $w(x) = w_i(x)$ if $x \in X_i$ is a componentwise weak weight. \square

According to Proposition 2.16, a directed graph $\Gamma = (V, E)$ is componentwise weakly weighted if and only if each strongly connected component (i.e., each large-scale connected component of (V, d_Γ)) is a non-directed graph. So, one can easily find an example of such a directed graph not satisfying condition (b) in Proposition 2.13, and so providing an example of a componentwise weakly weighted generalised quasi-metric space that is not weakly weighted.

In the following example we provide two generalised quasi-metric spaces that are componentwise weakly weighted. However, for the proof of the property we wait for stronger results (see Example 6.3).

Example 2.21. (a) For a set S , we define a generalised quasi-metric $d_{\mathcal{P}(S)}$ on $\mathcal{P}(S)$ as in (2.6).

(b) For an abelian group G , we define a generalised quasi-metric $d_{L(G)}$ on $L(G)$ as in (2.7), with the abuse of notation $\log \infty = \infty$. The large-scale geometry of the metric space $(L(G), d_{L(G)}^s)$ is studied in [20].

Even though the two examples provided in Example 2.21 are componentwise weakly weighted (we prove it in Example 6.3), they are not weakly weighted unless the set S (respectively, the group G) is finite, i.e., unless they are quasi-metric spaces according to Proposition 2.13. In fact, if S (respectively, G) is infinite, then the pair \emptyset and S (respectively, $\{0\}$ and G) does not satisfy condition (b) of the mentioned result.

The equivalence relation \sim from (2.5) can be extended in the following way. Given a family $\{S_i\}_{i \in I}$ of pairwise disjoint subsets of a set S (most commonly, a partition of S), two functions $f, g: S \rightarrow \mathbb{R}$ are equivalent relatively to the family $\{S_i\}_{i \in I}$, and we write

$$f \approx_{\{S_i\}_i} g, \quad \text{if } f \upharpoonright_{S_i} \sim g \upharpoonright_{S_i} \text{ for every } i \in I. \quad (2.8)$$

If the family is clear from the context, we simply write \approx for $\approx_{\{S_i\}_i}$.

Proposition 2.6(a) implies its extension to componentwise weak weights for generalised quasi-metric spaces.

Corollary 2.22. *Let X be a generalised quasi-metric space, and w be a componentwise weak weight for X . Then the set of all componentwise weak weights for X is $[w]_{\approx}$ where \approx is relative to $\{\mathcal{Q}(x)\}_{x \in X}$.*

3. Generalisations of partial metrics

3.1. Partial metrics and their relation with weakly weighted quasi-metrics

Partial metrics were introduced by Matthews ([34]). Then, O'Neill ([33]) extended the notion maintaining the same terminology. In this paper we prefer to distinguish the two notions, and we call them partial metrics and weak partial metrics, respectively.

Definition 3.1 ([33]). A *weak partial metric* p on a non-empty set X is a function $p: X \times X \rightarrow \mathbb{R}$ satisfying the following properties:

- (PM1) for every $x, y \in X$, $x = y$ if and only if $p(x, x) = p(y, y) = p(x, y)$;
- (PM2) for every $x \in X$, the function $p(x, \cdot)$ is minimised at x (i.e., $p(x, x) \leq p(x, y)$, for every $y \in X$);
- (PM3) p is symmetric (i.e., for every $x, y \in X$, $p(x, y) = p(y, x)$);
- (PM4) for every $x, y, z \in X$, $p(x, z) \leq p(x, y) + p(y, z) - p(y, y)$.

A *partial metric*, according to [34], is a weak partial metric p on X satisfying the further property $p \geq 0$.

The pair (X, p) is a *(weak) partial metric space* if p is a (weak) partial metric on X .

If we consider the equivalence relation \sim defined in (2.5), it is trivial to check that, if (X, p) is a weak partial metric space, every function $p': X \times X \rightarrow \mathbb{R}$ satisfying $p' \sim p$ is actually a weak partial metric. A similar result holds for partial metrics with the further assumption that $p' \geq 0$.

Before introducing some examples, let us show how weak partial metrics are tightly connected to weakly weighted quasi-metrics. The corresponding connection between partial metrics and weighted quasi-metrics was proved in [34], where the two notions were introduced.

Theorem 3.2. *Let X be a non-empty set.*

(a) *If d is a quasi-metric on X that is weakly weighted by w , then the function $p_{d,w}: X \times X \rightarrow \mathbb{R}$ defined by the law*

$$p_{d,w}(x, y) := d(x, y) + w(x), \quad \text{for every } x, y \in X,$$

is a weak partial metric on X . If w' is another weak weight for d , then $p_{d,w'} \sim p_{d,w}$.

In particular, if w is a weight (and so d is weighted) then $p_{d,w}$ is a partial metric.

(b) *If p is a weak partial metric on X , then the function $d_p: X \times X \rightarrow \mathbb{R}_{\geq 0}$ defined by the law*

$$d_p(x, y) := p(x, y) - p(x, x), \quad \text{for every } x, y \in X,$$

is a quasi-metric on X which is weakly weighted by $w_p: X \rightarrow \mathbb{R}$, where w_p is defined by $w_p(x) = p(x, x)$ for every $x \in X$. Moreover, if $p': X \times X \rightarrow \mathbb{R}$ satisfies $p' \sim p$, then p' is a weak partial metric, $d_{p'} = d_p$ and $w_{p'} \sim w_p$.

In particular, if p is a partial metric, then w_p is a weight for d_p .

Proof. (a) Let w be a weak weight for the quasi-metric d on X . Properties (PM1) and (PM2) are easily verified for $p_{d,w}$ and (PM3) descends from (2.3). Let us show (PM4). If $x, y, z \in X$, then the triangle inequality (QM2) yields

$$\begin{aligned} p_{d,w}(x, z) &= d(x, z) + w(x) \leq d(x, y) + w(x) + d(y, z) + w(y) - w(y) = \\ &= p_{d,w}(x, y) + p_{d,w}(y, z) - p_{d,w}(y, y). \end{aligned}$$

The second assertion can be easily verified, in view of Proposition 2.6, whereas the last assertion is clear by definition.

(b) Let p be a weak partial metric. Property (PM2) implies that $d_p \geq 0$.

Property (QM1) for d_p is satisfied because of (PM1) and (PM3). In fact, for $x, y \in X$, assuming that $d_p(x, y) = d_p(y, x) = 0$, we have that $p(x, x) = p(x, y)$ and $p(y, x) = p(y, y)$; in view of (PM3) $p(x, y) = p(y, x)$, so $p(x, x) = p(x, y) = p(y, x) = p(y, y)$ and we get $x = y$ by (PM1).

Let us prove property (QM2) by using (PM4). For every $x, y, z \in X$ one has

$$d_p(x, z) = p(x, z) - p(x, x) \leq p(x, y) + p(y, z) - p(y, y) - p(x, x) = d_p(x, y) + d_p(y, z).$$

Finally, by construction and (PM3), w_p is a weak weight for d_p .

The second assertion can be easily verified, also using Proposition 2.6, whereas the last assertion is clear by definition. \square

As mentioned in the introduction, the particular case of the following correspondence between weighted quasi-metrics and partial metrics was already pointed out by Matthews in [34].

Corollary 3.3. *Let X be a non-empty set. If d is a quasi-metric on X that is weakly weighted by w and if p is a weak partial metric on X , then*

$$d_{p_{d,w}} = d \quad \text{and} \quad p_{d_p, w_p} = p.$$

So, there exists a one-to-one correspondence between weakly weighted quasi-metrics (respectively, weighted quasi-metrics) on X and equivalence classes of weak partial metrics (respectively, partial metrics) on X .

Proof. Let us consider a quasi-metric d weakly weighted by w . By Theorem 3.2(a), $[p_{d,w}]_{\sim}$ does not depend on the choice of w , so the map $d \mapsto [p_{d,w}]_{\sim}$ is well-defined. Moreover, for every $x, y \in X$,

$$d_{p_{d,w}}(x, y) = p_{d,w}(x, y) - p_{d,w}(x, x) = d(x, y) + w(x) - d(x, x) - w(x) = d(x, y).$$

In particular, $d_{p_{d,w}}$ does not depend on the choice of w .

Conversely, let us take two weak partial metrics $p \sim p'$ on X . According to Theorem 3.2(b), $d_p = d_{p'}$ and it is weakly weighted, so the map $[p]_{\sim} \mapsto d_p$ is well-defined.

Moreover, for every $x, y \in X$,

$$p_{d_p, w_p}(x, y) = d_p(x, y) + w_p(x) = p(x, y) - p(x, x) + p(x, x) = p(x, y). \quad \square$$

Example 3.4. Let Σ^* be the family of all strings, both finite and infinite, on the alphabet Σ . For every pair of strings $s, s' \in \Sigma$, we denote by $|s|$ the length of s and by $l(s, s')$ the length of the longest common prefix of s and s' . Then we define a partial metric p on Σ^* by the law $p(s, s') = 2^{-l(s, s')}$, for every $s, s' \in \Sigma^*$.

Using Theorem 3.2 we can associate to p the quasi-metric d_p (that is, $d_p(s, s') = 2^{-l(s, s')} - 2^{-|s|}$, for every $s, s' \in \Sigma^*$), which is weighted by the function w_p (namely, $w_p(s) = 2^{-|s|}$, for every $s \in \Sigma^*$).

The following example from [3] provides a weak partial metric space (X, p) coming from a biological setting, satisfying the following stronger condition with respect to (PM2):

(PM2S) for every $x, y \in X$ with $x \neq y$, $p(x, x) < p(x, y)$.

Example 3.5. Let $\Sigma^{<\infty}$ be the family of all finite strings on the (finite) alphabet Σ . For two strings $\bar{x} = (x_1, \dots, x_n), \bar{y} = (y_1, \dots, y_m) \in \Sigma^{<\infty}$, an *alignment of \bar{x} and \bar{y}* is given by two strings $\bar{x}' = (x'_1, \dots, x'_k)$ and $\bar{y}' = (y'_1, \dots, y'_k)$ of the same length that can be obtained from \bar{x} and \bar{y} , respectively, by adding occurrences of a new, blank, character $\# \notin \Sigma$ at any positions. Given such an alignment \bar{x}' and \bar{y}' , we compute its *score* $s'(\bar{x}', \bar{y}')$ by comparing the two strings at each position $i \in \{1, \dots, k\}$ and adding the following real values:

- α if $x'_i = y'_i \neq \#$;
- β if $x'_i \neq y'_i$ and both $x'_i \neq \#$ and $y'_i \neq \#$;
- γ if precisely one of the two characters x'_i and y'_i is $\#$;
- 0 if $x'_i = y'_i = \#$.

For example, the alignment

$$\begin{aligned} \bar{x}' &= G \# A \# T T A C A \# \quad \text{and} \\ \bar{y}' &= G C A \# T C A C G A \end{aligned}$$

of the strings $\bar{x} = GATTACA$ and $\bar{y} = GCATCACGA$ on the alphabet $\Sigma = \{G, A, T, C\}$ has score $s'(\bar{x}', \bar{y}') = 5\alpha + 2\beta + 2\gamma + 0$.

Then we assign to the two strings \bar{x} and \bar{y} the following value:

$$s(\bar{x}, \bar{y}) = \max\{s'(\bar{x}', \bar{y}') \mid \bar{x}' \text{ and } \bar{y}' \text{ form an alignment of } \bar{x} \text{ and } \bar{y}\}.$$

The *score scheme* just described is a useful technique to compare partial DNA strands (see [21]).

The function $p: \Sigma^{<\infty} \times \Sigma^{<\infty} \rightarrow \mathbb{R}$ defined by $p(\bar{x}, \bar{y}) = -s(\bar{x}, \bar{y})$, for every $\bar{x}, \bar{y} \in \Sigma^{<\infty}$, is a weak partial metric with (PM2S) provided that $\alpha > \beta$, $\alpha > \gamma$, $\beta \geq 2\gamma$ and $\gamma < 0$ (see [3, Proposition 2.1]).

If we note that, for every $\bar{x} \in \Sigma^{<\infty}$, \bar{x} and \bar{y} provide the best alignment of \bar{x} and \bar{y} themselves (the one with the highest score), we can characterise the weakly weighted quasi-metric d_p induced by p as follows: for every $\bar{x}, \bar{y} \in \Sigma^{<\infty}$, $d_p(\bar{x}, \bar{y}) = \alpha|\bar{x}| - s(\bar{x}, \bar{y})$.

3.2. Generalised partial metrics and their relation with weakly weighted generalised quasi-metrics

Inspired by the notions of weak partial metric and generalised quasi-metric space, we introduce the following.

Definition 3.6. A *generalised weak partial metric* p on a non-empty set X is a function $p: X \times X \rightarrow \mathbb{R} \cup \{\infty\}$ satisfying the properties (PM1)–(PM4) of Definition 3.1 (with the usual convention that $b < \infty + a = a + \infty = \infty$, for every $a, b \in \mathbb{R}$) and the following further one:

(PM5) for every $x \in X$, $p(x, x) < \infty$.

The pair (X, p) is called *generalised weak partial metric space*.

Note that property (PM4) is well-defined thanks to (PM5).

Similarly to what we have done for generalised quasi-metric spaces, we can introduce an equivalence relation \cong_p on a generalised partial metric space (X, p) as follows: for every $x, y \in X$, $x \cong_p y$ if $p(x, y) < \infty$. Then the equivalence classes of \cong_p are called *p -connected components*. Moreover, for every $x \in X$, $\mathcal{Q}_X^p(x)$ denotes the p -connected component to which x belongs, i.e., $[x]_{\cong_p}$. To distinguish between these two notions and the corresponding ones defined for generalised quasi-metric spaces, we emphasise the role of p in the notation.

As one may expect, also weakly weighted generalised quasi-metric spaces and generalised weak partial metric spaces are two faces of the same coin.

Theorem 3.7. *Let X be a non-empty set.*

(a) *If d is a generalised quasi-metric on X which is weakly weighted by w , then the function $p_{d,w}: X \times X \rightarrow \mathbb{R} \cup \{\infty\}$ defined by putting*

$$p_{d,w}(x, y) := d(x, y) + w(x), \quad \text{for every } x, y \in X,$$

is a generalised weak partial metric on X . Moreover, $\cong_d = \cong_{p_{d,w}}$.

(b) *If p is a generalised weak partial metric on X , then the function $d_p: X \times X \rightarrow \mathbb{R}_{\geq 0} \cup \{\infty\}$ defined by*

$$d_p(x, y) := p(x, y) - p(x, x), \quad \text{for every } x, y \in X,$$

is a generalised quasi-metric weakly weighted by the function $w_p: X \rightarrow \mathbb{R}$ defined by $w_p(x) := p(x, x)$, for every $x \in X$. Moreover, $\cong_{d_p} = \cong_p$.

Proof. The proof is an easy adaptation of that of Theorem 3.2. To verify the coincidence of the equivalence relations, use the symmetry of the partial metrics. \square

In order to fully extend Theorem 3.2, we need to discuss the uniqueness part of the statement. Let d be a generalised quasi-metric on a non-empty set X weakly weighted by w and w' , and let $\{X_i\}_{i \in I}$ represent the partition of X in its large-scale connected components with respect to d . They coincide with the p -connected components with respect to $p_{d,w}$ and $p_{d,w'}$ in view of Theorem 3.7(a). Then $p_{d,w} \approx_{\{X_i \times X_i\}_i} p_{d,w'}$ (see (2.8) for the definition of \approx).

On the other hand, let p and p' be two generalised weak partial metrics on X , with $\cong_p = \cong_{p'}$ and $p \approx_{\{X_i \times X_i\}_i} p'$, where $\{X_i\}_{i \in I}$ is the family of all p -connected components of X . They coincide with the large-scale connected components with respect to d_p and $d_{p'}$ by Theorem 3.7(b). Then $d_p = d_{p'}$ and $w_p \approx_{\{X_i\}_i} w_{p'}$.

Now, as in Corollary 3.3, we obtain the following correspondence.

Corollary 3.8. *Let X be a non-empty set. If d is a generalised quasi-metric on X weakly weighted by w and p is a generalised weak partial metric on X , then $d_{p_w} = d$ and $p_{d_p, w_p} = p$.*

Hence, there is a one-to-one correspondence between weakly weighted generalised quasi-metrics on X and equivalence classes with respect to $\approx_{\{X_i \times X_i\}_i}$ of generalised weak partial metrics on X inducing the same partition into large-scale connected components and p -connected components $\{X_i\}_{i \in I}$.

Let us provide an example of a generalised weak partial metric (and so, according to Theorem 3.7, of a weakly weighted generalised quasi-metric) inspired by Example 3.4.

Example 3.9. Let Σ be an alphabet. Define the set

$$\Sigma_{-\infty}^* = \{(x_n)_{n \in (-\infty, m) \cap \mathbb{Z}} \mid m \in \mathbb{Z} \cup \{\infty\}, \forall n < m, x_n \in \Sigma\}.$$

Let us fix some notation. For every $\bar{x} = (x_n)_{n \in (-\infty, m) \cap \mathbb{Z}} \in \Sigma_{-\infty}^*$, set $L_{\bar{x}} = m - 1$ if $m < \infty$, and $L_{\bar{x}} = \infty$ otherwise. One can think of $L_{\bar{x}}$ as the last index for which \bar{x} is defined. Then, for every $k \leq L_{\bar{x}}$, $\bar{x}[k]$ denotes the substring obtained by cutting \bar{x} at the level k , i.e., $\bar{x}[k] = (x_n)_{n \in (-\infty, k] \cap \mathbb{Z}}$. Set now, for every $\bar{x}, \bar{y} \in \Sigma_{-\infty}^*$,

$$l(\bar{x}, \bar{y}) = \sup\{k \in \mathbb{Z} \mid k \leq \min\{L_{\bar{x}}, L_{\bar{y}}\}, \bar{x}[k] = \bar{y}[k]\},$$

where the supremum is computed in the totally ordered set $\mathbb{Z} \cup \{+\infty, -\infty\}$ with the usual conventions $\sup \emptyset = -\infty$ and $\sup \mathbb{Z} = +\infty$. Then the function $p: \Sigma_{-\infty}^* \times \Sigma_{-\infty}^* \rightarrow \mathbb{R} \cup \{\infty\}$ defined by putting, for every $\bar{x}, \bar{y} \in \Sigma_{-\infty}^*$, $p(\bar{x}, \bar{y}) = 2^{-l(\bar{x}, \bar{y})}$ is a generalised weak partial metric on $\Sigma_{-\infty}^*$. If Σ has at least two elements σ_1, σ_2 , it is not a weak partial metric; in fact, the distance between the constant sequences $(\sigma_1)_{n \in \mathbb{Z}}$ and $(\sigma_2)_{n \in \mathbb{Z}}$ is infinite. Moreover, according to Theorem 3.7, d_p is a weakly weighted generalised quasi-metric.

Remark 3.10. Let now (X, d) be a generalised quasi-metric space and w be a componentwise weak weight for it. Then it can be easily shown that the function $p_{d,w}$ defined as

$$p_{d,w}(x, y) = \begin{cases} d(x, y) + w(x) & \text{if } x \cong_d y, \\ \infty & \text{otherwise,} \end{cases}$$

for every $x, y \in X$, is indeed a generalised weak partial metric. By Corollary 3.8, it follows that every componentwise weak weight induces a weakly weighted generalised quasi-metric on X which can be explicitly described as follows: for every $x, y \in X$,

$$d_{p_{d,w}}(x, y) = \begin{cases} d(x, y) & \text{if } x \cong_d y, \\ \infty & \text{otherwise.} \end{cases}$$

In particular, the identity map $id_X : (X, d_{p_{d,w}}) \rightarrow (X, d)$ is non-expansive and $d_{p_{d,w}}$ is the coarsest weakly weighted generalised quasi-metric on X having this property, i.e., if d' is a weakly weighted generalised quasi-metric on X such that $id_X : (X, d') \rightarrow (X, d)$ is non-expansive, then $id_X : (X, d') \rightarrow (X, d_{p_{d,w}})$ is non-expansive.

For example, let $\Gamma = (V, E)$ be a directed graph endowed with the generalised quasi-metric d_Γ and suppose that each strongly connected component of Γ is a non-directed graph (see §2.2 and §2.3). In other words, d_Γ is componentwise weakly weighted, say by w_Γ . In this case, the quasi-metric $d_{p_{d_\Gamma, w_\Gamma}}$ coincides with $d_{\Gamma'}$, where Γ' is obtained as the disjoint union of all strongly connected components of Γ .

4. Generalised quasi-metric semilattices

Let (X, \leq) be a *partially ordered set* (briefly, *poset*), i.e., a pair of a set X and a *partial order* \leq on it, which is a reflexive, transitive and antisymmetric relation. The *dual poset* (X, \geq) consists of the same set X and the *inverse order* \geq (i.e., $x \geq y$ if and only if $y \leq x$). A subset Y of X is said to be *convex* if, for every $x, y, z \in X$, $y \in Y$ provided that $x \leq y \leq z$ and $x, z \in Y$.

A *meet-semilattice* (respectively, *join-semilattice*) is a poset (X, \leq) such that for every $x, y \in X$ the *meet* $x \wedge y = \inf\{x, y\}$ (respectively, the *join* $x \vee y = \sup\{x, y\}$) exists. If a poset (X, \leq) is simultaneously a meet- and a join-semilattice, then it is a *lattice*. Vice versa, any lattice can be seen separately as a join-semilattice or as meet-semilattice.

Remark 4.1. The poset (X, \leq) is a meet-semilattice if and only if (X, \geq) is a join-semilattice. More precisely, we have that $x \wedge_{\leq} y = x \vee_{\geq} y$, for every $x, y \in X$.

We write simply *semilattice* to indicate either a meet-semilattice or a join-semilattice when the statement holds for both of them. In such a case we use an $*$ to indicate the semilattice operation. Indeed, a semilattice can be seen as a commutative semigroup where every object is idempotent, and we can retrieve the partial order from the semilattice operation $*$: if $* = \wedge$, then $x \leq y$ if and only if $x = x \wedge y$, and if $* = \vee$, then $x \leq y$ if and only if $y = x \vee y$.

When maintaining the distinction between meet- and join-semilattices is instead necessary, sometimes we write results explicitly only for meet-semilattices, while the dual statements can be similarly deduced using Remark 4.1.

An equivalence relation \cong on a semilattice $(X, *)$ is a *congruence* if, for every $x, y, z, w \in X$, $x * z \cong y * w$ provided that $x \cong y$ and $z \cong w$.

Fact 4.2. Let $(X, *)$ be a semilattice and \cong a congruence on it. Then every equivalence class of \cong is a convex subsemilattice of X with respect to the partial order \leq of $(X, *)$.

Proof. Let $x, y \in X$ satisfying $x \cong y$. Then $x * y \cong x * x = x$, thus $[x]_{\cong}$ is a subsemilattice of X . To prove that $[x]_{\cong}$ is convex, take two points $y, z \in X$ such that $x \leq y \leq z$ and $z \cong x$. Then $x * y \cong z * y$, and thus $x \cong y$ (if $* = \wedge$, then $x \wedge y = x$ and $z \wedge y = y$, while, if $* = \vee$, then $x \vee y = y$ and $z \vee y = z \cong x$). \square

4.1. The specialisation order of a generalised quasi-metric

Every generalised quasi-metric space (X, d) comes with the partial order \leq_d , called *specialisation order*, i.e., for $x, y \in X$, $x \leq_d y$ if and only if $d(x, y) = 0$.

Remark 4.3. The partial order \leq_d is indeed nothing but the specialisation order of the quasi-metric topology τ_d induced by d on X . Let us recall that τ_d is the T_0 -topology generated by the base $\{B_\varepsilon^d(x) \mid x \in X, \varepsilon > 0\}$, where $B_\varepsilon^d(x) = \{y \in X \mid d(x, y) < \varepsilon\}$ is the ε -ball around the point $x \in X$. Moreover, if τ is a topology on a set X , then the *specialisation order* \leq_τ of τ is defined as follows: for $x, y \in X$, $x \leq_\tau y$ if y is contained in the closure of $\{x\}$.

Fact 4.4 (Monotonicity). Let (X, d) be a generalised quasi-metric space and $x_1, x_2, y_1, y_2 \in X$. Then

$$d(x_1, y_1) \leq d(x_2, y_2) \text{ provided } x_1 \leq_d x_2 \text{ and } y_2 \leq_d y_1.$$

Proof. Take four points $x_1, x_2, y_1, y_2 \in X$ such that $x_1 \leq_d x_2$ and $y_2 \leq_d y_1$. Then the triangle inequality (QM2) implies

$$d(x_1, y_1) \leq d(x_1, x_2) + d(x_2, y_2) + d(y_2, y_1) = d(x_2, y_2),$$

since $d(x_1, x_2) = 0 = d(y_2, y_1)$. \square

Remark 4.5. Given a generalised quasi-metric space (X, d) , for every $x \in X$, $\mathcal{Q}(x)$ is convex with respect to \leq_d . In fact, fix $\bar{x} \in X$, take $x, y, z \in X$, and suppose that $x, z \in \mathcal{Q}(\bar{x})$ and $x \leq_d y \leq_d z$. Since $z \in \mathcal{Q}(\bar{x}) = \mathcal{Q}(x)$ and $d(y, z) = 0$ by hypothesis, we get $d(y, x) \leq d(y, z) + d(z, x) = d(z, x) < \infty$. Hence, $y \in \mathcal{Q}(x)$ as $d(x, y) = 0$ by assumption.

We extend the following definition from [40] (see also [41] where it is called *order-convexity*, and [22, Definition VI-1.17] and [27, Exercise 6.7.19] where it is called *radial convexity*) to every generalised quasi-metric space.

Definition 4.6. For a generalised quasi-metric space (X, d) , the *descending path condition* is: (DPC) for every $x, y, z \in X$ satisfying $x \geq_d y \geq_d z$, one has $d(x, z) = d(x, y) + d(y, z)$.

The following result immediately descends from the definition of (DPC).

Fact 4.7. Let (X, d) be a generalised quasi-metric space and Y be a subset of X . If (X, d) satisfies (DPC), then $(Y, d \upharpoonright_Y)$ satisfies (DPC).

As a straightforward consequence of Fact 4.7, if a generalised quasi-metric space satisfies (DPC), then so does each large-scale connected component. Moreover, (DPC) can be separately verified in each large-scale connected component.

Proposition 4.8. Let (X, d) be a generalised quasi-metric space. Then (X, d) satisfies (DPC) if and only if each large-scale connected component $(\mathcal{Q}(x), d \upharpoonright_{\mathcal{Q}(x)})$ of X satisfies it.

Proof. The ‘only if’ implication follows from Fact 4.7. Conversely, suppose that $\mathcal{Q}(x)$ satisfies (DPC) for every $x \in X$. Let $x \geq_d y \geq_d z$ be elements of X , for which (QM2) yields $d(x, z) \leq d(x, y) + d(y, z)$. If $d(x, z) = \infty$ then $d(x, z) = d(x, y) + d(y, z)$. Assume now that $d(x, z) < \infty$, and so $z \in \mathcal{Q}(x)$. Since $\mathcal{Q}(x)$ is convex by Remark 4.5, also $y \in \mathcal{Q}(x)$. Thus, $d(x, z) = d(x, y) + d(y, z)$ since $\mathcal{Q}(x)$ satisfies (DPC). \square

Proposition 4.8 was proved in a more restrictive context in [16].

4.2. Invariant generalised quasi-metric semilattices

Given a poset (X, \leq) , we say that a (generalised) quasi-metric d on X is *compatible* with the partial order of X if $\leq_d = \leq$. In the rest of this section, we focus on generalised quasi-metrics defined on semilattices that are compatible with the algebraic structure of the underlying space.


Definition 4.9. A *generalised quasi-metric meet-semilattice* (respectively, *generalised quasi-metric join-semilattice*) is a generalised quasi-metric space (X, d) such that the poset (X, \leq_d) is a meet-semilattice (respectively, join-semilattice).

A generalised quasi-metric meet-semilattice (respectively, join-semilattice) satisfying $d(x, y) = d(x, x \wedge y)$ (respectively, $d(x, y) = d(x \vee y, y)$) for all $x, y \in X$ is said to be *invariant*

In [32] invariant semilattices are called *optimal*.

Remark 4.10. Given a generalised quasi-metric space (X, d) , it may occur that (X, \leq_d) is a lattice and that the invariance property holds for both the meet and the join, as in Example 4.16. Anyway, one can consider in the lattice (X, \leq_d) only the meet (respectively, join) and the invariance property for the meet (respectively, join), in other words one can use only the structure of invariant generalised quasi-metric meet-semilattice (respectively, join-semilattice).

This remark is particularly relevant since there are generalised quasi-metric spaces (X, d) inducing lattices (X, \leq_d) that are invariant for only one of the two lattice operations. Consider, for example, the four-point lattice $X = \{x, y, z, t\}$ represented in the following diagram:



It can be induced by a quasi-metric d defined by setting $d(x, y) = d(x, z) = d(x, t) = 2$ and $d(y, z) = d(y, t) = d(z, y) = d(z, t) = 1$ (all the other values are 0). Then, X is an invariant quasi-metric meet-semilattice, but it is not invariant if we consider the join operation. Similarly, (X, d^{-1}) is invariant with respect to the join operation, but it is not invariant as a meet-semilattice.

In the above definition of (generalised) quasi-metric semilattice, we do not require the semilattice operation to be quasi-uniformly continuous. This slight abuse of terminology is motivated by the fact that we mainly deal with invariant quasi-metric semilattices, which automatically meet the continuity requirement (see Proposition 4.13(b)).

In the sequel, when we work with generalised quasi-metric semilattices (X, \leq_d) , we simply denote by \leq the specialisation order \leq_d if there is no risk of ambiguity.

Remark 4.11. As a consequence of Remark 4.1, (X, d) is a generalised quasi-metric meet-semilattice if and only if (X, d^{-1}) is a generalised quasi-metric join-semilattice. More explicitly, $\leq_{d^{-1}}$ is the inverse order of \leq_d and $x \wedge_{\leq_d} y = x \vee_{\leq_{d^{-1}}} y$, for every $x, y \in X$. Moreover, (X, d) is invariant if and only if so is (X, d^{-1}) .

Similarly to (DPC) that is inherited by subspaces, invariance is inherited by subsemilattices as stated in the following result.

Fact 4.12. *Let X be a generalised quasi-metric semilattice and Y a subsemilattice of X endowed with the induced generalised quasi-metric. If X is invariant, then so is Y .*

Let us see some useful characterisations of invariance. The proof can be obtained by adjusting that provided in [16].

Proposition 4.13. *Let (X, d) be a generalised quasi-metric meet-semilattice. Then the following properties are equivalent:*

- (a) X is invariant;
- (b) for all $x, y, z \in X$, $d(z \wedge x, z \wedge y) \leq d(x, y)$ (i.e., for every $z \in X$, the shift $s_z: x \mapsto z \wedge x$ is non-expansive);
- (c) for all $x, y, z \in X$, $d(x, y \wedge z) \leq d(x, y) + d(x, z)$ (i.e., for every $x \in X$, $d(x, \cdot): X \rightarrow \mathbb{R}_{\geq 0} \cup \{\infty\}$ is subadditive).

For quasi-metric join-semilattices a similar result holds (as for item (c), it should be replaced with the following condition: for all $x, y, z \in X$, $d(y \vee z, x) \leq d(y, x) + d(z, x)$).

By invariance, we see that the large-scale connected components are subsemilattices:

Proposition 4.14. *If (X, d) is an invariant generalised quasi-metric semilattice, then \cong_d is a congruence. In particular, every large-scale connected component $\mathcal{Q}(x)$ of (X, d) is a convex invariant subsemilattice.*

Proof. To prove that \cong_d is a congruence, let $x, y, z, w \in X$ satisfying $x \cong_d y$ and $z \cong_d w$. Then, applying (QM2) and Proposition 4.13(b), we obtain that

$$d(x * z, y * w) \leq d(x * z, x * w) + d(x * w, y * w) \leq d(z, w) + d(x, y) < \infty,$$

and similarly $d(y * w, x * z) < \infty$. Finally, Facts 4.2 and 4.12 imply the second part of the statement. \square

The convexity of the large-scale connected components of a generalised quasi-metric semilattice was already verified in Remark 4.5.

Let us mention that a (generalised) quasi-metric semilattice satisfying (DPC) is not necessarily invariant (see [16] and Example 4.15(a) below).

Next we review Examples 2.10, 2.17, 3.4 and 3.5 to see whether the quasi-metric spaces introduced there are generalised quasi-metric semilattices and what properties they satisfy.

Example 4.15. (a) Let us assume the notation used in Example 2.10. The specialisation orders $\leq_{d_{\mathbb{S}}}$, $\leq_{d_{\mathbb{T}}}$ and $\leq_{d_{\mathbb{R}}}$ coincide with the usual orders. Moreover, it is easy to see that $(\mathbb{S}, d_{\mathbb{S}})$, $(\mathbb{T}, d_{\mathbb{T}})$ and $(\mathbb{R}, d_{\mathbb{R}})$ are both invariant quasi-metric meet-semilattices and invariant quasi-metric join-semilattices satisfying (DPC).

(b) Let (\mathbb{T}, d) be the quasi-metric space defined in Example 2.11. Then the specialisation order \leq_d coincides with the usual order $0 \leq 1 \leq 2$. Moreover, (\mathbb{T}, d) is both an invariant quasi-metric meet-semilattice and an invariant quasi-metric join-semilattice. However, it does not satisfy (DPC).

(c) As for the family of all strings Σ^* endowed with the quasi-metric d_p (see Example 3.4), we can see that, for every $s, s' \in \Sigma^*$, $s \leq_{d_p} s'$ if and only if s is a prefix of s' (i.e., s coincides with an initial substring of s'). Thus the meet of two strings s and s' is given by the longest common prefix of s and s' . The pair (Σ^*, d_p) is then an invariant quasi-metric meet-semilattice with (DPC).

(d) Let $(\Sigma^{<\infty}, d_p)$ be the quasi-metric space defined in Example 3.5. Since the weak partial metric p satisfies property (PM2S), d_p satisfies the following strengthened version of (QM1): for every $x, y \in X$, $d_p(x, y) = 0$ if and only if $x = y$. This property implies that the topology induced by d_p satisfies the separation axiom T_1 . Hence, in particular, the specialisation order \leq_{d_p} coincides with equality, and so $\Sigma^{<\infty}$ is not a semilattice.

Let us now consider the generalised quasi-metric spaces defined in Example 2.21 and already studied in [11].

Example 4.16. (a) For a set X , the specialisation order $\leq_{d_{\mathcal{P}(X)}}$ coincides with the containment \supseteq , which means, more explicitly, that, for every $A, B \subseteq X$, $d_{\mathcal{P}(X)}(A, B) = 0$ (i.e., $A \leq_{d_{\mathcal{P}(X)}} B$) if and only if $A \supseteq B$. With this partial order, $\mathcal{P}(X)$ is a lattice, where the meet of A and B is their union, while the join is their intersection. Moreover, $(\mathcal{P}(X), d_{\mathcal{P}(X)})$ is both an invariant generalised quasi-metric meet-semilattice with (DPC) and an invariant generalised quasi-metric join-semilattice with (DPC).

(b) Given an abelian group G and two subgroups $H, K \in L(G)$, $d_{L(G)}(H, K) = 0$ (and so $H \leq_{d_{L(G)}} K$) if and only if $H \supseteq K$. Thus $L(G)$ with this partial order is a lattice, where, for every $H, K \in L(G)$, the meet of H and K is $H + K$ and the join is $H \cap K$. Then $(L(G), d_{L(G)})$ is both an invariant generalised quasi-metric meet-semilattice with (DPC) and an invariant generalised quasi-metric join-semilattice with (DPC).

Remark 4.17. Let X be a non-empty set. Any generalised weak partial metric p on X induces a partial order \leq_p on X defined as follows: if $x, y \in X$, then $x \leq_p y$ whenever $p(x, x) = p(x, y)$. It is trivial to see that $\leq_p = \leq_{p+c}$ for every constant $c \in \mathbb{R}$. The partial order \leq_p is just the other side of the coin of the specialisation order. In fact, we have that $\leq_p = \leq_{d_p}$, and, moreover, if d is a generalised quasi-metric on X weakly weighted by w , then $\leq_d = \leq_{p_{d,w}}$ (see Theorem 3.7 for the definitions

of d_p and $p_{d,w}$). In order to show it, let $x, y \in X$ and w be a weak weight for d . Then the claim follows from the following two chains:

$$p(x, y) - p(x, x) = d_p(x, y), \text{ and } p_{d,w}(x, y) - p_{d,w}(x, x) = d(x, y) + w(x) - d(x, x) - w(x) = d(x, y).$$

Therefore, (X, \leq_p) is a semilattice, and in this case we say that (X, p) is a *generalised weak partial metric semilattice*, if and only if (X, \leq_{d_p}) is a semilattice. Furthermore, if $*$ is the semilattice operation, $p(x, y) = p(x, x * y)$ for every $x, y \in X$ if and only if d_p is invariant. For consistency, we refer to this property by saying that p is *invariant*. Similarly, a generalised quasi-metric semilattice (X, d) weakly weighted by w is invariant if and only if so is $p_{d,w}$. Thus, Corollary 3.8 leads to a similar correspondence between invariant weakly weighted (generalised) quasi-metric semilattices and invariant (generalised) weak partial metric semilattices.

5. Weakly weighted quasi-metric semilattices and their characterisations

The notion of weak weightedness for generalised quasi-metric semilattices (and so that of generalised weak partial metric semilattice according to Remark 4.17) is too restrictive, as the following remark shows. Motivated by that observation, in this section we focus on weakly weighted quasi-metric semilattices.

Remark 5.1. Let (X, d) be a generalised quasi-metric meet-semilattice (the other case can be similarly treated). Suppose that there exist $x, y \in X$ such that $d(x, y) = \infty$ (and so d is not a quasi-metric). Fact 4.4 implies that $d(x, y) \leq d(x, x \wedge y)$, and so $d(x, x \wedge y) = \infty$. Moreover, $d(x \wedge y, x) = 0$, and so (X, d) cannot be weakly weighted in view of Proposition 2.13.

5.1. Weakly weighted quasi-metric semilattices and (DPC)

Proposition 5.2. *A quasi-metric space (X, d) satisfies (DPC) provided that it is weakly weighted.*

Proof. Let w be a weak weight for d , and $x, y, z \in X$ such that $x \geq_d y \geq_d z$. By Lemma 2.9, one has that $d(x, z) = d(x, y) + d(y, z)$, which implies that (DPC) holds. \square

We show in Theorem 5.4 that the statement of Proposition 5.2 can be reversed for invariant quasi-metric semilattices. To this end, we use the following functions and the subsequent lemma describing them.

Let (X, d) be an invariant quasi-metric semilattice satisfying (DPC), and $x \in X$. We define the function $w_x: X \rightarrow \mathbb{R}$ as follows:

$$w_x(y) := d(x, y) - d(y, x), \quad \text{for every } y \in X. \quad (5.1)$$

Lemma 5.3. *Let (X, d) be an invariant quasi-metric meet-semilattice satisfying (DPC). Then,*

$$w_x(y) = d(x, x \wedge y \wedge z) - d(y, x \wedge y \wedge z), \quad \text{for every } x, y, z \in X.$$

Proof. Let $x, y, z \in X$. Then invariance and (DPC) imply that

$$\begin{aligned} w_x(y) &= d(x, y) - d(y, x) = d(x, x \wedge y) - d(y, x \wedge y) = \\ &= d(x, x \wedge y) + d(x \wedge y, x \wedge y \wedge z) - d(y, x \wedge y) - d(x \wedge y, x \wedge y \wedge z) = \\ &= d(x, x \wedge y \wedge z) - d(y, x \wedge y \wedge z). \end{aligned} \quad \square$$

Theorem 5.4. *Let (X, d) be an invariant quasi-metric semilattice satisfying (DPC). Then, for every $x \in X$, w_x is a weak weight for d (so, (X, d) is weakly weighted). In particular, for all $x, y \in X$, $w_x \sim w_y$.*

Proof. We prove the statement assuming that (X, d) is a meet-semilattice. Fixing $x \in X$, we claim that, for every $y, z \in X$, $d(y, z) + w_x(y) = d(z, y) + w_x(z)$. Applying Lemma 5.3, we obtain that

$$\begin{aligned} w_x(y) - w_x(z) &= d(x, x \wedge y \wedge z) - d(y, x \wedge y \wedge z) - d(x, x \wedge y \wedge z) + d(z, x \wedge y \wedge z) = \\ &= d(z, x \wedge y \wedge z) - d(y, x \wedge y \wedge z) = w_z(y) = d(z, y) - d(y, z), \end{aligned}$$

which concludes the verification that w_x is a weak weight for d . The last assertion follows from Proposition 2.6(a). \square

So, Proposition 5.2 and Theorem 5.4 immediately give the following result.

Corollary 5.5. *An invariant quasi-metric semilattice is weakly weighted if and only if it satisfies (DPC).*

Let us consider the examples contained in Example 2.10 satisfying (DPC).

Example 5.6. (a) We start from the Sierpiński space $\mathbb{S} = \{0, 1\}$. Since $(\mathbb{S}, d_{\mathbb{S}})$ is an invariant quasi-metric semilattice with (DPC), according to Theorem 5.4, it is weakly weighted. Moreover,

$$w_1(x) = d_{\mathbb{S}}(1, x) - d_{\mathbb{S}}(x, 1) = d_{\mathbb{S}}(1, x) = 1 - x,$$

which is the weight provided in Example 2.10(a).

- (b) Analogously to (a), we recover the weight w defined in Example 2.10(b) for $(\mathbb{T}, d_{\mathbb{T}})$ as w_2 .
(c) In Example 2.10(c) we have provided a weak weight w for the quasi-metric space $(\mathbb{R}, d_{\mathbb{R}})$ with $d_{\mathbb{R}}(x, y) = \max\{x - y, 0\}$ for every $x, y \in \mathbb{R}$. A straightforward computation gives $w = w_0$.

Next we review the examples contained in Example 2.12.

- Example 5.7.** (a) As in Example 2.12(a), let X be a finite set and on the power set $\mathcal{P}(X)$ consider the quasi-metric $d_{\mathcal{P}(X)}$. We know that $d_{\mathcal{P}(X)}$ is weighted by the weight $w(A) = |A|$ for every $A \subseteq X$, and now we recover it as $w = w_{\emptyset}$.
(b) Let G be a finite abelian group and $d_{L(G)}$ be the quasi-metric on $L(G)$ defined in Example 2.12(b). We have already noticed in the mentioned example that $d_{L(G)}$ is weighted by the weight $w(H) = \log|H|$ for every subgroup H of G , and now we have that $w = w_{\{0\}}$.

Now we present some important applications of Theorem 5.4.

Theorem 5.8. *Let (X, d) be an invariant quasi-metric semilattice satisfying (DPC). Then the following properties are equivalent:*

- (a) (X, d) is weighted;
(b) for every $x \in X$, there exists $c_x \in \mathbb{R}$ such that $w_x(y) \geq c_x$, for every $y \in X$;
(c) there exist $x \in X$ and $c \in \mathbb{R}$ such that $w_x(y) \geq c$, for every $y \in X$.

If (X, d) is a join-semilattice, then also the following condition is equivalent:

- (d) (X, d) is an M -space ([41, Definition 2.1]), i.e., for every $x \in X$, there exists $c'_x \in \mathbb{R}_{\geq 0}$ such that, for every $y \in X$ satisfying $x \leq y$, $d(y, x) \leq c'_x$.

Proof. Theorem 5.4 implies that, for every $x \in X$, w_x is a weak weight for d . Then the equivalences (a) \Leftrightarrow (b) \Leftrightarrow (c) follow from Proposition 2.7(a).

(b) \Rightarrow (d) Let $x \in X$ and c_x as in (b). If $y \in X$ is such that $x \leq y$, then $c_x \leq w_x(y) = d(x, y) - d(y, x) = -d(y, x)$; in particular $c_x \leq 0$. Hence, $d(y, x) \leq -c_x =: c'_x \geq 0$.

(d) \Rightarrow (b) Let $x, y \in X$, and $c'_x \in \mathbb{R}$ satisfying the hypothesis in (d). Then invariance implies that

$$w_x(y) = d(x, y) - d(y, x) \geq -d(y, x) = -d(y \vee x, x) \geq -c'_x =: c_x. \quad \square$$

The equivalence of (a) and (d) in the above theorem provides another proof to [41, Theorem 2.31].

We can analogously obtain the counterpart of Theorem 5.8 characterising co-weighted quasi-metric semilattices. The equivalence of (a) and (d) in the theorem below covers [41, Theorem 2.34].

Theorem 5.9. *Let (X, d) be an invariant quasi-metric semilattice satisfying (DPC). Then the following properties are equivalent:*

- (a) (X, d) is co-weighted;
(b) for every $x \in X$, there exists $c_x \in \mathbb{R}$ such that $w_x(y) \leq c_x$, for every $y \in X$;
(c) there exist $x \in X$ and $c \in \mathbb{R}$ such that $w_x(y) \leq c$, for every $y \in X$.

If (X, d) is a meet-semilattice, then also the following condition is equivalent:

- (d) X is an m -space ([41, Definition 2.10]), i.e., for every $x \in X$, there exists $c'_x \in \mathbb{R}_{\geq 0}$ such that, for every $y \in X$ satisfying $y \leq x$, $d(x, y) \leq c'_x$.

Proof. Theorem 5.4 implies that, for every $x \in X$, w_x is a weak weight for d . Then the equivalences (a) \Leftrightarrow (b) \Leftrightarrow (c) follow from Proposition 2.7(b).

(b) \Rightarrow (d) Let $x \in X$ and c_x as in (b). Then, if $y \in X$ satisfies $y \leq x$,

$$c_x \geq w_x(y) = d(x, y) - d(y, x) = d(x, y).$$

(d) \Rightarrow (b) Let $x, y \in X$, and $c'_x \in \mathbb{R}$ satisfying the hypothesis in (d). Then, since X is invariant, and $x \wedge y \leq x$,

$$w_x(y) = d(x, y) - d(y, x) \leq d(x, y) = d(x, x \wedge y) \leq c'_x. \quad \square$$

Remark 5.10. Let us consider two particular cases of Theorems 5.8 and 5.9. Let (X, d) be an invariant quasi-metric semilattice satisfying (DPC) and with a bottom element \perp . The definition of the function w_{\perp} can be simplified as follows: for every $x \in X$, since $\perp \leq x$, $w_{\perp}(x) = d(\perp, x) - d(x, \perp) = -d(x, \perp) \leq 0$. Hence, (X, d) is co-weighted. If, otherwise X has a top element \top , for every $x \in X$, $w_{\top}(x) = d(\top, x) \geq 0$, and so (X, d) is weighted.

Clearly, the functions w_{\perp} and w_{\top} can be considered also in semilattices that do not satisfy (DPC). Then the following results can be proved.

(a) Let X be a quasi-metric meet-semilattice.

(a₁) If X has a bottom element \perp , then w_{\perp} is *supadditive* (i.e., $w_{\perp}(x \wedge y) \geq w_{\perp}(x) + w_{\perp}(y)$ for every $x, y \in X$). In fact, $w_{\perp}(x \wedge y) = -d(x, \perp)$, $w_{\perp}(x) + w_{\perp}(y) = -(d(x, \perp) + d(y, \perp))$, and $d(x \wedge y, \perp) \leq d(x, \perp) \leq d(x, \perp) + d(y, \perp)$ by the monotonicity of d (see Fact 4.4).

(a₂) If X has a top element \top and X is invariant, then w_{\top} is subadditive because of Proposition 4.13.

(b) Let X be a quasi-metric join-semilattice.

(b₁) If X has a top element \top , then w_{\top} is subadditive (the proof of item (a₁) can be easily adapted).

(b₂) If X has a bottom element \perp and X is invariant, then w_{\perp} is supadditive according to Proposition 4.13.

5.2. Semivaluations, semi-co-valuations and weak weights

We start recalling the definition of semivaluation from [35], giving also the corresponding notion of semi-co-valuation. These were proposed with the following terminology in [42] with domain $\mathbb{R}_{\geq 0}$ instead of the whole \mathbb{R} and using equivalent defining conditions.

Definition 5.11. Let (X, \leq) be a meet-semilattice. A function $f: X \rightarrow \mathbb{R}$ is called:

- *meet valuation* if $f(x) + f(x \wedge y \wedge z) \geq f(x \wedge y) + f(x \wedge z)$, for all $x, y, z \in X$;
- *meet co-valuation* if $f(x) + f(x \wedge y \wedge z) \leq f(x \wedge y) + f(x \wedge z)$, for all $x, y, z \in X$.

Let (X, \leq) be a join-semilattice. A function $f: X \rightarrow \mathbb{R}$ is called:

- *join valuation* if $f(x) + f(x \vee y \vee z) \leq f(x \vee y) + f(x \vee z)$, for all $x, y, z \in X$;
- *join co-valuation* if $f(x) + f(x \vee y \vee z) \geq f(x \vee y) + f(x \vee z)$, for all $x, y, z \in X$.

We refer to [16] for other characterisations of join valuations, and the respective equivalent conditions for the other properties can be easily and similarly obtained.

The following definition, as given in [35] and [42], conveniently summarises the notions provided in Definition 5.11.

Definition 5.12. A function $f: X \rightarrow \mathbb{R}$ on a semilattice (X, \leq) is called a *semivaluation* (respectively, *semi-co-valuation*) if it is either a meet or a join valuation (respectively, either a meet or a join co-valuation).

In other words, the prefixes join and meet before the term (co-)valuation appear when we want to emphasise the type of semilattice with which we are working. In the remaining cases we use the more general term semi(-co-)valuation.

Proposition 5.13. Let (X, \leq) be a semilattice and $f: X \rightarrow \mathbb{R}$ a function.

- (a) *The function f is a semivaluation (respectively, semi-co-valuation) if and only if $-f$ is a semi-co-valuation (respectively, semivaluation).*
- (b) *If f is a semi(-co-)valuation, then every $f' \in [f]_{\sim}$ has the same property.*
- (c) *If f is a semivaluation, then f is non-decreasing.*
- (d) *If f is a semi-co-valuation, then f is non-increasing.*

Proof. Items (a) and (b) are trivial, while item (c) follows from items (a) and (d).

We want to prove item (d). Assume that X is a meet-semilattice and f is a meet co-valuation. If $x, y \in X$ satisfy $x \leq y$, then

$$f(y) + f(x) = f(y) + f(y \wedge x \wedge x) \leq f(y \wedge x) + f(y \wedge x) = f(x) + f(x),$$

and so $f(y) \leq f(x)$, that is, f is non-increasing. If X is a join-semilattice and f is a join co-valuation, then the claim can be similarly shown. \square

As for the monotonicity results provided in Proposition 5.13(c) and (d), proofs can be found in [42, Lemma 4] and [35, Lemma 3.1(1)]. The same items of Proposition 5.13 also imply that a semi(-co-)valuation is strictly monotone if and only if it is injective.

Let us now start describing the connection between weak weights and semi(-co-)valuations.

Theorem 5.14. Let (X, d) be an invariant quasi-metric semilattice and let w be a weak weight for d . Then w is a strictly decreasing semi-co-valuation, while $-w$ is a strictly increasing semivaluation.

Proof. We prove the result for (X, d) a meet-semilattice. Let $x, y, z \in X$. Invariance and Proposition 4.13(b) imply that $d(x \wedge z, x \wedge y \wedge z) \leq d(x, x \wedge y)$. Since $d(x \wedge y \wedge z, x \wedge z) = 0 = d(x \wedge y, x)$, (2.3) implies that

$$w(x \wedge y \wedge z) - w(x \wedge z) = d(x \wedge z, x \wedge y \wedge z) \leq d(x, x \wedge y) = w(x \wedge y) - w(x),$$

and so w is a meet co-valuation. Moreover, w is strictly decreasing as, if $x < y$, then $d(y, x) > 0$ thanks to (QM1) since $d(x, y) = 0$, and so

$$w(y) < w(y) + d(y, x) = d(x, y) + w(x) = w(x).$$

That $-w$ is a strictly increasing meet valuation follows from Proposition 5.13(a) and (c). \square

When a semilattice admits a semi(-co-)valuation, it can be equipped with a distance function as follows.

Definition 5.15. If X is a meet-semilattice and f is a meet co-valuation, we define $d_f: X \times X \rightarrow \mathbb{R}_{\geq 0}$ as follows:

$$d_f(x, y) := f(x \wedge y) - f(x), \quad \text{for every } x, y \in X.$$

Dually, if X is a join-semilattice and f is a join co-valuation, we define $d_f: X \times X \rightarrow \mathbb{R}_{\geq 0}$ as follows:

$$d_f(x, y) := f(y) - f(x \vee y), \quad \text{for every } x, y \in X.$$

Definition 5.15 generalises that given in [42, Theorem 11] to co-valuations that admit also negative values.

The following result shows how to produce weakly weighted invariant quasi-metrics from semi(-co-)valuations,

Theorem 5.16. *Let (X, \leq) be a semilattice and $f: X \rightarrow \mathbb{R}$. If f is a strictly decreasing semi-co-valuation, then d_f is an invariant quasi-metric on X satisfying $\leq = \leq_{d_f}$, and weakly weighted by f .*

If f is a strictly increasing semivaluation, then d_{-f} is an invariant quasi-metric on X satisfying $\leq = \leq_{d_{-f}}$, and weakly weighted by $-f$.

Proof. Assume that (X, \leq) is a meet-semilattice and so f is a meet co-valuation. Let us first prove that d_f is a quasi-metric. Since f is decreasing, d_f assumes non-negative values. To prove (QM1), take two points $x, y \in X$ such that $d_f(x, y) = d_f(y, x) = 0$. Then the definition of d_f implies that $f(x) = f(x \wedge y) = f(y)$. As f is strictly monotone, the equalities $x = x \wedge y = y$ follow. To verify the triangular inequality (QM2), let $x, y, z \in X$. Due to the fact that f is a non-increasing meet co-valuation,

$$f(y) + f(x \wedge z) \leq f(y) + f(x \wedge z \wedge y) \leq f(x \wedge y) + f(z \wedge y),$$

and so, again since f is non-increasing,

$$\begin{aligned} d_f(x, z) &= f(x \wedge z) - f(x) = f(x \wedge y) - f(x) + f(x \wedge z) - f(x \wedge y) = \\ &= d_f(x, y) + f(x \wedge z) - f(x \wedge y) \leq d_f(x, y) + f(y \wedge z) - f(y) = d_f(x, y) + d_f(y, z). \end{aligned}$$

Note that $x \leq_{d_f} y$ if and only if $f(x) = f(x \wedge y)$, which is then equivalent to $x = x \wedge y$ and therefore to $x \leq y$. Hence, $\leq = \leq_{d_f}$. Finally, d_f is invariant by construction and it can be easily seen that f is a weak weight for d_f .

The second part of the theorem follows from the first one. In fact, if f is a strictly increasing semivaluation, then $-f$ is a strictly decreasing semi-co-valuation, by Proposition 5.13(a). Thus, we can associate to f the map d_{-f} . \square

Corollary 5.17. *Let (X, \leq) be a semilattice. There are one-to-one correspondences between the three sets consisting of the following objects:*

- (a) *invariant weakly weighted quasi-metrics d on X satisfying $\leq_d = \leq$;*
- (b) *equivalence classes of strictly decreasing semi-co-valuations f on X ;*
- (c) *equivalence classes of strictly increasing semivaluations g on X .*

More precisely, to d we associate $[w]_{\sim}$ and $[-w]_{\sim}$, where w is a weak weight for d , to $[f]_{\sim}$ we associate d_f and $[-f]_{\sim}$, and, finally, to $[g]_{\sim}$ we associate d_{-g} and $[-g]_{\sim}$.

Proof. Let (X, \leq) be a meet-semilattice. We prove the correspondence between (a) and (b) as that between (b) and (c) is trivial by Theorem 5.16 and Proposition 5.13(a). Both associations described in the statement are well-defined, respectively, one by virtue of Theorem 5.14 and Proposition 2.6(a), and the converse one by Theorem 5.16 since two semi-co-valuations f, g of X satisfy $d_f = d_g$ provided that $f \sim g$.

Moreover, we have to verify that the two associating maps are one the inverse of the other. Indeed, given a strictly decreasing semi-co-valuation f of X , we have seen in Theorem 5.16 that f is a weak weight for d_f and with d_f we associate $[f]_{\sim}$. It remains to check that if d is an invariant quasi-metric on X with $\leq_d = \leq$ and weakly weighted by w , then $d = d_w$. For $x, y \in X$, since $d(x \wedge y, x) = 0$ and d is invariant,

$$d_w(x, y) = w(x \wedge y) - w(x) = w(x \wedge y) + d(x \wedge y, x) - w(x) = d(x, x \wedge y) = d(x, y). \quad \square$$

As particular cases of Corollary 5.17, we find the following classical results due to Schellekens [42] since each (co-)weighted quasi-metric admits a unique fading (co-)weight in view of Remark 2.8.

Corollary 5.18. (a) ([42, Theorem 11]) *Let (X, \leq) be a meet-semilattice. There are one-to-one correspondences between the three sets consisting of the following objects:*

- (i) *invariant co-weighted quasi-metrics d on X with $\leq_d = \leq$;*
- (ii) *fading strictly increasing meet valuations on X ;*
- (iii) *fading strictly decreasing meet co-valuations on X .*

(b) ([42, Theorem 10]) *Let (X, \leq) be a join-semilattice. There are one-to-one correspondences between the three sets consisting of the following objects:*

- (i) *invariant weighted quasi-metrics d on X with $\leq_d = \leq$;*
- (ii) *fading strictly decreasing join co-valuations on X ;*
- (iii) *fading strictly increasing join valuations on X .*

6. Componentwisely weakly weighted generalised quasi-metric semilattices and their characterisations

Remark 5.1 shows that the notion of weak weightedness is a bad fit with generalised quasi-metric semilattices. In what follows we prove characterisations of componentwise weakly weighted generalised quasi-metric semilattices similar to those obtained for weakly weighted quasi-metric semilattices.

Proposition 5.2 extends to generalised quasi-metric spaces as follows.

Proposition 6.1. *If (X, d) is a componentwise weakly weighted generalised quasi-metric space, then (X, \leq_d) satisfies (DPC).*

Proof. It immediately follows from Fact 2.20, and Propositions 5.2 and 4.8. \square

Similarly to the previous result, the following theorem can be proved applying Propositions 6.1 and 4.8, Theorem 5.4 and Fact 2.20.

Theorem 6.2. *Let (X, d) be an invariant generalised quasi-metric semilattice. Then (X, d) satisfies (DPC) if and only if it is componentwise weakly weighted. Moreover, w is a componentwise weak weight for d precisely when $w \approx w_X$, where w_X is defined by $w_X \upharpoonright_{\mathcal{Q}(x_i)} = w_{x_i}$ for every $i \in I$, where $\{x_i\}_{i \in I}$ is a fixed family of representatives of the large-scale connected components of X .*

Example 6.3. Using Theorem 6.2 we can prove that the generalised quasi-metric meet-semilattices provided in Example 2.21 are componentwise weakly weighted. It is in fact easier to see that they satisfy (DPC) (see [16]), and so Theorem 6.2 implies that they are componentwise weakly weighted.

(a) Let S be a set. For the reasoning above, the generalised quasi-metric semilattice $(\mathcal{P}(S), d_{\mathcal{P}(S)})$ is componentwise weakly weighted. Let us construct a componentwise weak weight. Fix a family $\{A_i\}_{i \in I}$ of representatives of the equivalence classes of $\cong_{d_{\mathcal{P}(S)}}$. Then define $w: \mathcal{P}(X) \rightarrow \mathbb{R}$ as follows: for every $A \subseteq X$, take the index $i \in I$ such that $A \cong_{d_{\mathcal{P}(S)}} A_i$ and set

$$w(A) = w_{A_i}(A) = d_{\mathcal{P}(S)}(A_i, A) - d_{\mathcal{P}(S)}(A, A_i) = |A \setminus A_i| - |A_i \setminus A|.$$

(b) Let us now explicitly provide the componentwise weak weight for the generalised quasi-metric semilattice $(L(G), d_{L(G)})$, where G is an abelian group. Let $\{K_i\}_{i \in I}$ be a family of representatives of the equivalence classes of $\cong_{d_{L(G)}}$. Then define $w: L(G) \rightarrow \mathbb{R}$ as follows: for every $i \in I$, w coincides with w_{K_i} on $\mathcal{Q}(K_i)$, i.e., for every subgroup H of G such that $H \cong_{d_{L(G)}} K_i$,

$$w(H) = w_{K_i}(H) = d_{L(G)}(K_i, H) - d_{L(G)}(H, K_i) = \log|H + K_i : K_i| - \log|H + K_i : H|.$$

Keeping Theorem 6.2 in mind, we generalise the notion of semivaluations and semi-co-valuations in order to extend Corollary 5.17 to generalised quasi-metrics (see Corollary 6.11).

Definition 6.4. Let (X, \leq) be a meet-semilattice. Given a congruence \cong on X , a function $f: X \rightarrow \mathbb{R}$ is called:

- *generalised meet valuation with respect to \cong* if $f(x) + f(x \wedge y \wedge z) \geq f(x \wedge y) + f(x \wedge z)$, for every $x, y, z \in X$ satisfying $x \cong x \wedge z$ and $y \cong x \wedge y$;
- *generalised meet co-valuation with respect to \cong* if $f(x) + f(x \wedge y \wedge z) \leq f(x \wedge y) + f(x \wedge z)$, for every $x, y, z \in X$ satisfying $x \cong x \wedge z$ and $y \cong x \wedge y$.

Let (X, \leq) be a join-semilattice. Given a congruence \cong , a function $f: X \rightarrow \mathbb{R}$ is called:

- *generalised join valuation with respect to \cong* if $f(x) + f(x \vee y \vee z) \leq f(x \vee y) + f(x \vee z)$, for every $x, y, z \in X$ satisfying $x \cong x \vee z$ and $y \cong x \vee y$;
- *generalised join co-valuation with respect to \cong* if $f(x) + f(x \vee y \vee z) \geq f(x \vee y) + f(x \vee z)$, for every $x, y, z \in X$ satisfying $x \cong x \vee z$ and $y \cong x \vee y$.

Similarly to the non-generalised case, a *generalised semivaluation* is either a generalised join valuation or a generalised meet valuation, and a *generalised semi-co-valuation* is either a generalised join co-valuation or a generalised meet co-valuation.

It is easy to see that the notions provided in Definition 6.4 extend the ones given in Definition 5.11. In fact, for example, if X is a meet-semilattice, then $f: X \rightarrow \mathbb{R}$ is a meet co-valuation if and only if it is a generalised meet co-valuation with respect to the trivial congruence having just one equivalence class X .


Let us provide an immediate, but useful fact.

Fact 6.5. *Let x, y, z be three points in a semilattice X and \cong be a congruence. If $x \cong x * z$ and $y \cong x * y$, then*

$$y \cong x * y \cong x * y * z \cong y * z.$$

Proof. Since $x \cong x * z$ and \cong is a congruence, $x * y \cong x * y * z$. Finally, the convexity of the equivalence class (see Fact 4.2) implies that $y \cong y * z$ since $y \cong x * y * z$. \square

The following diagram represents the statement of Fact 6.5 if X is a meet-semilattice. The hypotheses are marked by the symbol \cong on the edge connecting the two elements. Then the elements of the equivalence class that the fact relates to one another are framed.



Proposition 6.6. *Let (X, \leq) be a semilattice, \cong be a congruence on X and $f: X \rightarrow \mathbb{R}$ be a function.*

- (a) The function f is a generalised semivaluation if and only if $-f$ is a generalised semi-co-valuation.
- (b) If f is a generalised semi(-co-)valuation, then every $f' \approx f$ with respect to $\{[x]_{\cong}\}_{x \in X}$ has the same property.
- (c) If f is a generalised semivaluation, then $f \upharpoonright_{[x]_{\cong}}$ is a semivaluation for every $x \in X$. In particular, $f \upharpoonright_{[x]_{\cong}}$ is non-decreasing.
- (d) If f is a generalised semi-co-valuation, then $f \upharpoonright_{[x]_{\cong}}$ is a semi-co-valuation. In particular, $f \upharpoonright_{[x]_{\cong}}$ is non-increasing.

It is important to note that we cannot extend the final implications of items (c) and (d) of Proposition 6.6 and state that a generalised semi(-co-)valuation is monotone with respect to the whole semilattice. On an intuitive level, item (b) of Proposition 6.6 represents an obstruction to that property. We provide an explicit example of a strictly increasing generalised semi-co-valuation.

Example 6.7. Consider the lattice $\mathbb{S} = \{0, 1\}$ as defined in Example 2.10(a), and the congruence \cong given by equality. It can be verified that the map $f = id_{\mathbb{S}}$ is a generalised meet co-valuation even though it is strictly increasing.

Corollary 6.8. Let (X, d) be an invariant generalised quasi-metric semilattice and w be a componentwise weak weight for d . Then w is a generalised semi-co-valuation with respect to \cong_d such that, for every $x \in X$, $w \upharpoonright_{\mathcal{Q}(x)}$ is strictly decreasing. Thus $-w$ is a generalised semivaluation with respect to \cong_d such that, for every $x \in X$, $w \upharpoonright_{\mathcal{Q}(x)}$ is strictly increasing.

Proof. Assume that (X, \leq_d) is a meet-semilattice. Let us take three points $x, y, z \in X$ such that $x \cong_d x \wedge z$ and $y \cong_d x \wedge y$. By Fact 6.5, $y \cong_d x \wedge y \cong_d x \wedge y \wedge z \cong_d y \wedge z$. Then, thanks to Proposition 4.13(b) and the invariance of d ,

$$\begin{aligned} w(x) &= w(x) + d(x, x \wedge z) - d(x, x \wedge z) = w(x \wedge z) - d(x, x \wedge z) \leq \\ &\leq w(x \wedge z) - d(x \wedge y, x \wedge y \wedge z) - w(x \wedge y) + w(x \wedge y) = \\ &= w(x \wedge z) - w(x \wedge y \wedge z) + w(x \wedge y). \end{aligned}$$

Hence, w is a generalised meet co-valuation with respect to \cong_d . Theorem 5.14 implies the second part of the statement. \square

The second statement follows from the first one by applying Proposition 6.6(a). \square

Definition 6.9. If X is a meet-semilattice, \cong a congruence on X and $f: X \rightarrow \mathbb{R}$ is a generalised meet co-valuation, we define $d_f: X \times X \rightarrow \mathbb{R}_{\geq 0} \cup \{\infty\}$ as follows: for every $x, y \in X$,

$$d_f(x, y) := \begin{cases} f(x \wedge y) - f(x) & \text{if } x \cong x \wedge y, \\ \infty & \text{otherwise.} \end{cases}$$

If X is a join-semilattice, \cong a congruence on X and $f: X \rightarrow \mathbb{R}$ is a generalised join co-valuation, we define $d_f: X \times X \rightarrow \mathbb{R}_{\geq 0} \cup \{\infty\}$ as follows: for every $x, y \in X$,

$$d_f(x, y) := \begin{cases} f(y) - f(x \vee y) & \text{if } y \cong x \vee y, \\ \infty & \text{otherwise.} \end{cases}$$

Theorem 6.10. Let (X, \leq) be a semilattice and \cong a congruence on it.

If f is a generalised semi-co-valuation such that $f \upharpoonright_{[x]_{\cong}}$ is strictly decreasing for every $x \in X$, then d_f is an invariant generalised quasi-metric on X satisfying $\leq = \leq_{d_f}$, and it is componentwise weakly weighted by f . Moreover, $\cong = \cong_{d_f}$.

If f is a generalised semivaluation such that $f \upharpoonright_{[x]_{\cong}}$ is strictly increasing for every $x \in X$, then d_{-f} is an invariant generalised quasi-metric on X satisfying $\leq = \leq_{d_{-f}}$, and it is componentwise weakly weighted by $-f$. Moreover, $\cong = \cong_{d_{-f}}$.

Proof. We give a proof when (X, \leq) is a meet-semilattice and f is a generalised meet co-valuation.

First of all, we claim that, $\cong = \cong_{d_f}$ (note that \cong_{d_f} can be defined even if d_f is not a priori a generalised quasi-metric). Indeed, let $x, y \in X$. If $x \cong y$, then, since \cong is a congruence, $x \cong x \wedge y \cong y$, and thus $d_f(x, y) < \infty$ and $d_f(y, x) < \infty$. Vice versa, if $d_f(x, y) < \infty$ and $d_f(y, x) < \infty$, then $x \cong x \wedge y$ and $y \cong x \wedge y$, and so $x \cong y$.

Let us now show that d_f is a generalised quasi-metric. Since f is non-increasing on each $[x]_{\cong}$, d_f assumes non-negative values. To prove (QM1), take two points $x, y \in X$ such that $d_f(x, y) = d_f(y, x) = 0$. Then the definition of d_f implies that $x \cong x \wedge y \cong y$ and $f(x) = f(x \wedge y) = f(y)$. As f is strictly monotone on $[x]_{\cong}$, the equalities $x = x \wedge y = y$ follow.

Let us prove that d_f satisfies (QM2). For every triple of points $x, y, z \in X$, we claim that

$$d_f(y, z) \leq d_f(y, x) + d_f(x, z). \quad (6.1)$$

If either $d_f(y, x) = \infty$ or $d_f(x, z) = \infty$, there is nothing to prove. Therefore, we can assume that $y \cong x \wedge y$ and $x \cong x \wedge z$. Then Fact 6.5 implies that also $y \cong y \wedge z$. Since f is a generalised meet co-valuation and f is non-increasing on the equivalence classes of \cong , we obtain the following chain:

$$\begin{aligned} f(x \wedge y) - f(y) + f(x \wedge z) - f(x) &\geq f(x \wedge y) - f(y) + f(x \wedge z) + f(x \wedge y \wedge z) - f(x \wedge y) - f(x \wedge z) = \\ &= f(x \wedge y \wedge z) - f(y) \geq f(y \wedge z) - f(y). \end{aligned}$$

Hence, (6.1) follows by plugging in the definition of d_f .

Note that, for $x, y \in X$, $x \leq_{d_f} y$ if and only if $x \cong x \wedge y$ and $f(x) = f(x \wedge y)$, which is then equivalent to $x = x \wedge y$ and therefore to $x \leq y$. Hence, $\leq = \leq_{d_f}$. Finally, d_f is invariant by construction and it can be easily seen that f is a componentwise weak weight for d_f .

The second statement can be derived from the first one by using Proposition 6.6. \square

Using the steps provided in this subsection, we can now extend Corollary 5.17 to generalised quasi-metrics.

Corollary 6.11. *Let (X, \leq) be a semilattice and \cong a congruence on it. There are one-to-one correspondences between the three sets consisting of the following objects:*

- (a) *invariant componentwise weakly weighted generalised quasi-metrics d on X satisfying $\leq_d = \leq$;*
- (b) *equivalence classes of generalised semi-co-valuations f strictly decreasing on each $[x]_{\cong}$;*
- (c) *equivalence classes of generalised semivaluations g strictly increasing on each $[x]_{\cong}$.*

More explicitly, to d we associate $[w]_{\approx}$ and $[-w]_{\approx}$ where w is a componentwise weak weight for d and \approx is relative to $\{\mathcal{Q}_d(x)\}_{x \in X}$. To $[f]_{\approx}$, where \approx is relative to $\{[x]_{\cong}\}_{x \in X}$, we associate d_f and $[-f]_{\approx}$. Finally, to $[g]_{\approx}$, where \approx is relative to $\{[x]_{\cong}\}_{x \in X}$, we associate d_{-g} and $[-g]_{\approx}$.

Proof. Let (X, \leq) be a meet-semilattice with a congruence \cong . We prove the correspondence between (a) and (b) since that between (b) and (c) is trivial by Theorem 6.10 and Proposition 6.6(a).

Both associations described in the statement are well-defined by virtue of Corollaries 2.22 and 6.8, Proposition 6.6(b) and Theorem 6.10 since two generalised semi-co-valuations f, f' of X satisfy $d_f = d_{f'}$ provided that $f \approx f'$.

Moreover, we have to verify that the two associating maps are one the inverse of the other. Indeed, given a generalised semi-co-valuation f of X strictly decreasing on each $[x]_{\cong}$, we have seen in Theorem 6.10 that f is a generalised weak weight for d_f ; and then to d_f we associate $[f]_{\approx}$ as in Corollary 6.8. On the other hand, to check that if d is an invariant generalised quasi-metric on X with $\leq_d = \leq$ and componentwise weakly weighted by w , then $d = d_w$, proceed as in the proof of Corollary 5.17, taking into account that, for $x, y \in X$, $d(x, y) = \infty$ if and only if $d_w(x, y) = \infty$ as $\cong = \cong_d$ by Theorem 6.10. \square

7. Entropy for meet-semilattices with respect to an equivalence relation

A self-map $\phi: X \rightarrow X$ of a meet-semilattice X is an *endomorphism* if, for every $x, y \in X$, $\phi(x \wedge y) = \phi(x) \wedge \phi(y)$. For $n \in \mathbb{N}_+$, the n -th *trajectory* of ϕ in $x \in X$ is

$$T_n(\phi, x) := x \wedge \phi(x) \wedge \cdots \wedge \phi^{n-1}(x).$$

Fix a function $f: X \rightarrow \mathbb{R}$ and, for $x \in X$, let

$$h_f(\phi, x) := \limsup_{n \rightarrow \infty} \frac{f(T_n(\phi, x))}{n}. \quad (7.1)$$

Clearly, if the function $g: X \rightarrow \mathbb{R}$ satisfies $g \sim f$, then $h_g(\phi, x) = h_f(\phi, x)$ for every $x \in X$.

We introduce in the following definition a new notion of intrinsic entropy for endomorphisms of meet-semilattices with respect to an equivalence relation.

Definition 7.1. Let X be a meet-semilattice, \cong an equivalence relation on X and ϕ an endomorphism of X . A point $x \in X$ is (ϕ, \cong) -*inert* if $T_n(\phi, x) \cong x$ for every $n \in \mathbb{N}_+$. Let $\mathcal{I}_{\phi}(X, \cong)$ be the set of all (ϕ, \cong) -inert points of X .

For a function $f: X \rightarrow \mathbb{R}$, the *intrinsic semilattice entropy relative to \cong* is defined as

$$\tilde{h}_{f, \cong}(\phi) := \sup\{h_f(\phi, x) \mid x \in X, x \text{ is } (\phi, \cong)\text{-inert}\}.$$

Whenever the equivalence relation \cong is a congruence on X , we can provide the following characterisations of a (ϕ, \cong) -inert point. We say that an endomorphism ϕ of a meet-semilattice X *respects* an equivalence relation \cong if $\phi(x) \cong \phi(y)$ for every $x, y \in X$ satisfying $x \cong y$.

Lemma 7.2. *Let X be a meet-semilattice, \cong a congruence on X , ϕ an endomorphism of X and $x \in X$. Then the following conditions are equivalent:*

- (a) *x is (ϕ, \cong) -inert;*
- (b) *$x \wedge \phi^n(x) \cong x$ for every $n \in \mathbb{N}$.*

Moreover, they imply the following item:

- (c) *$x \wedge \phi(x) \cong x$.*

In addition, if ϕ respects \cong , then (a) and (b) are equivalent to (c).

Proof. (a) \Rightarrow (b) For every $n \in \mathbb{N}_+$ one has that $x \cong T_{n+1}(\phi, x)$ by assumption. By the convexity of $[x]_{\cong}$ provided by Fact 4.2, also $x \wedge \phi^n(x) \cong x$.

(b) \Rightarrow (a) For $n = 1$ one has $T_1(\phi, x) = x \cong x$. Thus, we proceed by induction and assume that $T_n(\phi, x) \cong x$ for $n \in \mathbb{N}_+$. Notice that $T_{n+1}(\phi, x) = \phi^n(x) \wedge T_n(\phi, x)$. Thus, since \cong is a congruence, (b) and the induction hypothesis imply that $T_{n+1}(\phi, x) \cong \phi^n \wedge x \cong x$, which concludes the proof.


The implication (b) \Rightarrow (c) is trivial.

(c) \Rightarrow (a) Assume that $x \cong x \wedge \phi(x)$. As $T_1(\phi, x) = x \cong x$, we proceed by induction: if $T_n(\phi, x) \cong x$ for some $n \in \mathbb{N}_+$, then $\phi(T_n(\phi, x)) \cong \phi(x)$, and so, since \cong is a congruence, we have

$$T_{n+1}(\phi, x) = x \wedge \phi(T_n(\phi, x)) \cong x \wedge \phi(x) \cong x. \quad \square$$

Item (c) of Lemma 7.2 is strictly weaker than (a) and (b) if ϕ does not respect \cong . We provide a small counterexample.

Example 7.3. Consider the lattice X described by the following Hasse diagram



the endomorphism ϕ defined by $\phi(\top) = x$, $\phi(x) = \phi(y) = \phi(\perp) = \perp$, and the congruence \cong partitioning X into the two subsets $\{\top, x\}$ and $\{y, \perp\}$. Note that ϕ does not respect \cong since $\phi(\top) \not\cong \phi(x)$ even though $\top \cong x$. The point \top satisfies item (c) of Lemma 7.2, but $\top \wedge \phi^2(\top) = \perp \not\cong \top$.

Remark 7.4. Let X be a meet-semilattice together with a congruence \cong , and let ϕ be an endomorphism of X that respects \cong .

- (a) If $x \in X$ is (ϕ, \cong) -inert, then every $y \in [x]_{\cong}$ is (ϕ, \cong) -inert. Indeed, $y \cong x \cong x \wedge \phi(x) \cong y \wedge \phi(y)$ and then Lemma 7.2(c) applies.
- (b) Consequently, $\mathcal{I}_{\phi}(X, \cong)$ is a disjoint union of congruence classes of X .

7.1. Entropy for generalised normed semilattices

Following [15, 16], a *generalised normed meet-semilattice* is a meet-semilattice X with a function $v: X \rightarrow \mathbb{R}_{\geq 0} \cup \{\infty\}$, and so v is called *generalised norm*. Let \cong_v be the equivalence relation induced by the following partition:

$$\mathcal{F}_v(X) := \{x \in X \mid v(x) < \infty\} \quad \text{and} \quad X_{\infty} := X \setminus \mathcal{F}_v(X).$$

Define $f_v: X \rightarrow \mathbb{R}_{\geq 0}$ by letting $f_v(x) = v(x)$ if $x \in \mathcal{F}_v(X)$ and $f_v(x) = 0$ otherwise.

The following result shows that $h_{f_v, \cong_v}(\phi)$ coincides with the *semigroup entropy* $h_v(\phi)$ from [16] by slightly generalising that invented in [15]. It is worth to recall that many of the classical entropy functions in mathematics, such as the measure entropy ([30, 46]), several instances of topological entropy ([2, 29, 23]), many kinds of (adjoint) algebraic entropy ([2, 14, 17, 19, 26, 38, 50]) and the two set-theoretic entropies ([4, 12]), are found to be semigroup entropies, and in almost all these cases the underlying semigroup is a semilattice.

Proposition 7.5. *Let $v: X \rightarrow \mathbb{R}_{\geq 0} \cup \{\infty\}$ be a generalised norm on a meet-semilattice X that is subadditive on $\mathcal{F}_v(X)$, and let $\phi: X \rightarrow X$ be an endomorphism that is monotone with respect to v (i.e., for every $x \in X$, $v(\phi(x)) \leq v(x)$). Then, for every $x \in \mathcal{F}_v(X)$, x is (ϕ, \cong_v) -inert and*

$$h_{f_v}(\phi, x) = \limsup_{n \rightarrow \infty} \frac{v(T_n(\phi, x))}{n}. \quad (7.2)$$

In particular, $\tilde{h}_{f_v, \cong_v}(\phi) = \sup\{h_{f_v}(\phi, x) \mid x \in \mathcal{F}_v(X)\}$ and so $\tilde{h}_{f_v, \cong_v}(\phi) = h_v(\phi)$.

Proof. Let $x \in \mathcal{F}_v(X)$. Since ϕ is monotone with respect to v , for all $n \in \mathbb{N}_+$, $v(\phi^n(x)) \leq v(x) < \infty$. We have to prove that for every $n \in \mathbb{N}_+$, also $T_n(\phi, x) \in \mathcal{F}_v(X)$, that is, $v(T_n(\phi, x)) < \infty$. We can proceed by induction, and the case $n = 1$ is clear as $T_1(\phi, x) = x$. Assume that $v(T_n(\phi, x)) < \infty$ for some $n \in \mathbb{N}_+$; then $v(T_{n+1}(\phi, x)) = v(T_n(\phi, x) \wedge \phi^n(x)) \leq v(T_n(\phi, x)) + v(\phi^n(x)) < \infty$ by the subadditivity of v , by inductive hypothesis and by the monotonicity of ϕ with respect to v .

The last assertion follows since every $x \in \mathcal{F}_v(X)$ is (ϕ, \cong_v) -inert and $h_{f_v}(\phi, x) = 0$ for every (ϕ, \cong_v) -inert $x \in X_{\infty}$, and by the definition of semigroup entropy. \square

The limit superior in (7.2) is actually a limit as proved in [15].

Remark 7.6. Let $v: X \rightarrow \mathbb{R}_{\geq 0} \cup \{\infty\}$ be a generalised norm on a meet-semilattice X , and let ϕ be an endomorphism of X .

If v is non-decreasing, then all the elements of $\mathcal{F}_v(X)$ are (ϕ, \cong_v) -inert and the conclusions of Proposition 7.5 hold true.

If, instead, v is non-increasing, all the points of X_{∞} are (ϕ, \cong_v) -inert. Therefore, if v is also subadditive on $\mathcal{F}_v(X)$ and ϕ is monotone with respect to v , every $x \in X$ is (ϕ, \cong_v) -inert in view of Proposition 7.5. Moreover,

$$h_{f_v}(\phi, x) = \begin{cases} \limsup_{n \rightarrow \infty} \frac{v(T_n(\phi, x))}{n} & \text{if } x \in \mathcal{F}_v(X), \\ 0 & \text{if } x \in X_{\infty}. \end{cases}$$

Remark 7.7. Let $v: X \rightarrow \mathbb{R}_{\geq 0} \cup \{\infty\}$ be a generalised norm on a meet-semilattice X , and let ϕ be an endomorphism of X . If v is non-increasing and subadditive, then \cong_v is a congruence: for every $x, y \in X$, $x \wedge y \in \mathcal{F}_v(X)$ if and only if both x and y belong to $\mathcal{F}_v(X)$. Moreover, if ϕ is monotone with respect to v , then $\phi(\mathcal{F}_v(X)) \subseteq \mathcal{F}_v(X)$. However, this is not enough to get that ϕ respects \cong_v : one needs either $\phi(X_{\infty}) \subseteq X_{\infty}$ or $\phi(X_{\infty}) \subseteq \mathcal{F}_v(X)$.

7.2. Entropy for componentwise weakly weighted invariant generalised quasi-metric semilattices

Let (X, d) be an invariant generalised quasi-metric meet-semilattice together with a componentwise weak weight w for d . Let ϕ be a non-expansive endomorphism of X (i.e., $d(\phi(x), \phi(y)) \leq d(x, y)$). Hence, we have the following data:

- the equivalence relation \cong_d is a congruence and so every large-scale connected component $\mathcal{Q}(x)$ is a convex invariant subsemilattice of X (see Proposition 4.14);
- ϕ respects \cong_d by definition of non-expansive endomorphism;
- for every $x \in X$ one has that $w \upharpoonright_{\mathcal{Q}(x)} \sim w_x$, where $w_x = d(x, y) - d(y, x) < \infty$ for every $y \in \mathcal{Q}(x)$ (see Theorem 6.2 and (5.1)).

The following result shows in particular that $\tilde{h}_{w, \cong_d}(\phi)$ coincides with the *intrinsic semilattice entropy* $h_d(\phi)$ from [11]. This entropy was introduced to get a unifying approach able to cover all (or at least, most) of the intrinsic-like entropies in literature: the intrinsic algebraic entropy ([18]), the algebraic and the topological entropies for locally linearly compact vector spaces ([9, 10]), the topological entropy for totally disconnected locally compact abelian groups ([13, 25]) and the algebraic entropy for compactly covered locally compact abelian groups ([13, 24]).

Proposition 7.8. *Let (X, d) be an invariant generalised quasi-metric meet-semilattice componentwise weakly weighted by w and ϕ be a non-expansive endomorphism of X .*

- (a) *An element $x \in X$ is (ϕ, \cong_d) -inert if and only if $d(x, \phi(x)) < \infty$.*
- (b) *For every $x \in \mathcal{I}_\phi(X, \cong_d)$,*

$$h_w(\phi, x) = \limsup_{n \rightarrow \infty} \frac{d(x, T_n(\phi, x))}{n}. \quad (7.3)$$

In particular, $h_w(\phi, x)$ does not depend on the choice of w .

Moreover, $\tilde{h}_{w, \cong_d}(\phi) = \sup\{\tilde{h}_w(\phi, x) \mid x \in X \text{ such that } d(x, \phi(x)) < \infty\}$ and so $\tilde{h}_{w, \cong_d}(\phi) = \tilde{h}_d(\phi)$.

Proof. (a) For every $x \in X$, $d(x, \phi(x)) = d(x, x \wedge \phi(x))$ by the invariance of d . Moreover, as $x \wedge \phi(x) \leq x$, $d(x \wedge \phi(x), x) = 0$ and so $x \wedge \phi(x) \cong_d x$ if and only if $d(x, \phi(x)) < \infty$. Lemma 7.2 concludes the proof.

(b) Let $x \in \mathcal{I}_\phi(X, \cong_d)$. First notice that, for every $n \in \mathbb{N}_+$, $T_n(\phi, x) \in [x]_{\cong_d}$. Since $w \upharpoonright_{\mathcal{Q}(x)} \sim w_x$ as stated above, there exists a constant $c_x > 0$ such that $w(T_n(\phi, x)) = w_x(T_n(\phi, x)) + c_x$ for every $n \in \mathbb{N}_+$. Moreover, since $T_n(\phi, x) \leq x$, $w_x(T_n(\phi, x)) = d(x, T_n(\phi, x))$, and so one computes

$$h_w(\phi, x) = \limsup_{n \rightarrow \infty} \frac{w(T_n(\phi, x))}{n} = \limsup_{n \rightarrow \infty} \frac{w_x(T_n(\phi, x))}{n} = \limsup_{n \rightarrow \infty} \frac{d(x, T_n(\phi, x))}{n}.$$

The last assertion follows from the definitions. □

The limit superior in (7.3) is actually a limit, as proved in [11, Theorem 3.15].

The requirement that the generalised quasi-metric meet-semilattice has to satisfy (DPC) may seem restrictive. However, this property is fulfilled by all the examples of intrinsic entropies collected in [11]. Nevertheless, we explicitly pose the following problem.

Question 7.9. *Let ϕ be a non-expansive endomorphism of an invariant generalised quasi-metric meet-semilattice (X, d) . Fix a family $\{x_i\}_{i \in I}$ of representatives of the large-scale connected components of X , and define w_X as in Theorem 6.2, i.e., by setting $w_X \upharpoonright_{\mathcal{Q}(x_i)} = w_{x_i}$, for every $i \in I$. For a (ϕ, \cong_d) -inert point $x \in X$, is it true that the value $\tilde{h}_{w_X}(\phi, x)$ does not depend on the choice of the representative family? If so, can we conclude that $h_{w_X}(\phi, x) = \limsup_{n \rightarrow \infty} \frac{d(x, T_n(\phi, x))}{n}$?*

7.3. How to recover some classical entropies

We conclude by listing some concrete and classical examples of entropies obtained using Example 2.21 (see also Examples 4.16 and 6.3).

7.3.1. The algebraic entropy

Let G be an abelian group and $(L(G), \supseteq)$ the meet-semilattice consisting of all subgroups of G ordered by reverse inclusion (see Example 4.16(b)). Every group endomorphism $\phi: G \rightarrow G$ induces an endomorphism of $L(G)$, which we call again $\phi: L(G) \rightarrow L(G)$, by setting $H \mapsto \phi(H)$ for every $H \in L(G)$. For $H \in L(G)$,

$$T_n(\phi, H) = H + \phi(H) + \cdots + \phi^{n-1}(H), \quad \text{for every } n \in \mathbb{N}_+.$$

- (a) The meet-semilattice $(L(G), \supseteq)$ can be equipped with the subadditive generalised norm

$$v: L(G) \rightarrow \mathbb{R} \cup \{\infty\}, \quad v(H) := \log |H|.$$

In particular, $\phi: L(G) \rightarrow L(G)$ is monotone with respect to v . By Proposition 7.5, one has

$$\tilde{h}_{f_v, \cong_v}(\phi) = \sup\{h_{f_v}(\phi, H) \mid H \in \mathcal{F}_v(L(G))\}.$$

Since $\mathcal{F}_v(L(G)) = \{H \leq G \mid H \text{ finite}\}$ and $f_v: L(G) \rightarrow \mathbb{R}$ is given by $f_v(H) = \log|H|$ if H is finite and $f_v(H) = 0$ otherwise, we get that

$$h_{f_v}(\phi, H) = \lim_{n \rightarrow \infty} \frac{\log|T_n(\phi, H)|}{n}, \quad \text{and so} \quad \tilde{h}_{f_v, \cong_v}(\phi) = \text{ent}(\phi), \quad (7.4)$$

where $\text{ent}(\phi)$ is the classical *algebraic entropy* of abelian group endomorphisms ([2, 15, 19, 50]).

(b) The pair $(L(G), d_{L(G)})$ is an invariant generalised quasi-metric meet-semilattice satisfying (DPC). Suppose that w is a componentwise weak weight for $d_{L(G)}$. By Proposition 7.8, we obtain that, for every $(\phi, \cong_{d_{L(G)}})$ -inert subgroup H of G ,

$$h_w(\phi, H) = \lim_{n \rightarrow \infty} \frac{\log|T_n(\phi, H) : H|}{n}, \quad \text{and so} \quad \tilde{h}_w(\phi) = \widetilde{\text{ent}}(\phi),$$

where $\widetilde{\text{ent}}(\phi)$ is the *intrinsic algebraic entropy* of ϕ as defined in [18] (see also [11, §5.2.1]).

Remark 7.10. In the above setting, as $\mathcal{F}_v(L(G))$ is a ϕ -invariant subsemilattice of $L(G)$, we can consider on it the restrictions ϕ_{fin} and v_{fin} of ϕ and v , respectively. Restricting the generalised quasi-metric $d_{L(G)}$ defined in Example 2.21(b), we obtain a quasi-metric d on $\mathcal{F}_v(L(G))$ for which v_{fin} is a weight. Therefore, v_{fin} is a strictly decreasing meet co-valuation by Theorem 5.14. Since $(\mathcal{F}_v(L(G)), d)$ has only one large-scale connected component, for every $E \in \mathcal{F}_v(G)$, $w_E \sim v_{fin}$ by Theorem 5.4 and Proposition 2.6(a), where w_E is defined in Example 6.3(b). Thus, given any fixed $E \in \mathcal{F}_v(G)$, for every $H \in \mathcal{F}_v(L(G))$,

$$h_{f_v}(\phi, H) = h_{w_E}(\phi_{fin}, H) = \lim_{n \rightarrow \infty} \frac{\log|T_n(\phi, H) + E : E| - \log|T_n(\phi, H) + E : T_n(\phi, H)|}{n},$$

which has to be compared with (7.4).

Example 7.11. Let now G be a compactly covered locally compact abelian group (i.e., the Pontryagin dual \widehat{G} of G is a totally disconnected locally compact abelian group) and $\phi: G \rightarrow G$ be a continuous endomorphism. Denote by $B(G)$ the subsemilattice of $L(G)$ consisting of all compact open subgroups of G and call $d_{B(G)}$ the restriction of $d_{L(G)}$ to $B(G)$.

In this case, $(B(G), d_{B(G)})$ is an invariant quasi-metric meet-semilattice satisfying (DPC), that is, we have only one large-scale connected component: for every $U, V \in B(G)$, $U \cong_{d_{B(G)}} V$; in fact, the indices $|U + V : V|$ and $|U + V : U|$ are finite since both U and V are open in the compact subgroup $U + V$.

Hence, fixed $V \in B(G)$, by Theorem 5.4 and Proposition 7.8(a), we get that

$$h_{w_V}(\phi, U) = \limsup_{n \rightarrow \infty} \frac{\log|T_n(\phi, U) : U|}{n}, \quad \text{for every } U \in B(G),$$

and so

$$h_{w_V}(\phi) = \sup\{h_{w_V}(\phi, U) \mid U \in B(G)\} = h_{alg}(\phi),$$

where $h_{alg}(\phi)$ is the *algebraic entropy* of ϕ as described in [13, Proposition 2.2]

For the general definition of the algebraic entropy for continuous endomorphisms of locally compact groups, see [12, 47]. In [15], it was already obtained as a semigroup entropy in the general case.

Remark 7.12. One can carry out considerations similar to those of Example 7.11 also in other settings. For example, they can be applied to the topological entropy of continuous endomorphisms of totally disconnected locally compact groups (see [11, 13, 15, 25]), and to the algebraic and the topological entropy of continuous endomorphisms of locally linearly compact vector spaces (see [9, 10]).

7.3.2. The set-theoretic entropy

Analogously to §7.3.1 for the algebraic entropy, one can treat the set-theoretic entropy as follows.

Let S be a non-empty set and $(\mathcal{P}(S), \supseteq)$ the meet-semilattice consisting of all subsets of S ordered by inverse inclusion (see Example 4.16(a)). Every self-map $\varphi: S \rightarrow S$ induces an endomorphism of $\mathcal{P}(S)$, that we call again $\varphi: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$, such that $A \mapsto \varphi(A)$ for every $A \in \mathcal{P}(S)$. Note that

$$T_n(\varphi, A) = A \cup \varphi(A) \cup \dots \cup \varphi^{n-1}(A), \quad \text{for every } n \in \mathbb{N}_+.$$

(a) The meet-semilattice $(\mathcal{P}(S), \supseteq)$ can be equipped with the subadditive generalised norm

$$v_{set}: \mathcal{P}(S) \rightarrow \mathbb{R} \cup \{\infty\}, \quad v_{set}(A) := |A|.$$

In particular, $\varphi: \mathcal{P}(S) \rightarrow \mathcal{P}(S)$ is monotone with respect to v_{set} . By Proposition 7.5,

$$\tilde{h}_{f_{v_{set}}, \cong_{v_{set}}}(\varphi) = \sup\{h_{f_{v_{set}}}(\varphi, A) \mid A \in \mathcal{F}_{v_{set}}(\mathcal{P}(S))\}.$$

Since $\mathcal{F}_{v_{set}}(\mathcal{P}(S)) = \{A \subseteq S \mid A \text{ finite}\}$ and $f_{v_{set}}: \mathcal{P}(S) \rightarrow \mathbb{R}$ is given by $f_{v_{set}}(A) = |A|$ if A is finite and $f_{v_{set}}(A) = 0$ otherwise, we get that

$$h_{f_{v_{set}}}(\varphi, A) = \lim_{n \rightarrow \infty} \frac{|T_n(\varphi, A)|}{n}, \quad \text{and so} \quad \tilde{h}_{f_{v_{set}}, \cong_{v_{set}}}(\varphi) = h_{set}(\varphi),$$

where $h_{set}(\varphi)$ is the classical *set-theoretic entropy* of φ ([4, 15, 16]).

- (b) The pair $(\mathcal{P}(S), d_{\mathcal{P}(S)})$ is an invariant generalised quasi-metric meet-semilattice satisfying (DPC). If w is a componentwise weak weight for $d_{\mathcal{P}(S)}$, by Proposition 7.8 one obtains that, for every $(\phi, \cong_{d_{\mathcal{P}(S)}})$ -inert subset A of S ,

$$h_w(\varphi, A) = \lim_{n \rightarrow \infty} \frac{|T_n(\varphi, A) \setminus A|}{n}, \quad \text{and so} \quad \tilde{h}_{w, \cong_{d_{\mathcal{P}(S)}}}(\varphi) = \tilde{h}_{set}(\varphi),$$

where $\tilde{h}_{set}(\varphi)$ is the *intrinsic set-theoretic entropy* of φ from [16]. In the same paper, it was also proved that always $\tilde{h}_{set}(\varphi) = h_{set}(\varphi)$.

Similar considerations as those in Remark 7.10 can be done also for the set-theoretic entropy.

References

- [1] S. ABRAMSKY, A. JUNG: *Domain Theory*. Handbook of logic in computer science, Vol. 3, 1–168, Handb. Log. Comput. Sci., 3, Oxford Sci. Publ., Oxford Univ. Press, New York, 1994.
- [2] R. L. ADLER, A. G. KONHEIM, M. H. MCANDREW: *Topological entropy*. Trans. Amer. Math. Soc. 114 (1965), 309–319.
- [3] S. ASSAF, K. PAL: *Partial metric spaces with negative distances and fixed point theorems*. Topology Proc. 49 (2017), 19–40.
- [4] F. AYATOLLAH ZADEH SHIRAZI, D. DIKRANJAN: *Set-theoretical entropy: A tool to compute topological entropy*. Proceedings ICTA2011 Islamabad Pakistan July 4–10 2011, 11–32, Cambridge Scientific Publishers, 2012.
- [5] G. BIRKHOFF: *Lattice theory*. American Mathematical Society, New York, 1940.
- [6] N. BOURBAKI: *Integration. I. Chapters 1–6*, Elements of Mathematics (Berlin). Translated from the 1959, 1965 and 1967 French originals by Sterling K. Berberian. Springer-Verlag, Berlin, 2004.
- [7] M. A. BUKATIN, J. S. SCOTT: *Towards computing distances between programs via Scott domains*. In: S. Adian, A. Nerode (Eds.), Logical Foundation of Computer Science, Lecture Notes in Coimputer Science, Vol. 1234, Springer, Berlin, 1997, pp. 33–43.
- [8] M. A. BUKATIN, S. Y. SHORINA: *Partial metrics and co-continuous valuations*. Foundations of software science and computation structures (Lisbon, 1998), 125–139, Lecture Notes in Comput. Sci., 1378, Springer, Berlin, 1998.
- [9] I. CASTELLANO, A. GIORDANO BRUNO: *Algebraic entropy in locally linearly compact vector spaces*. Rings, Polynomials, and Modules, Springer (2017), 103–127.
- [10] I. CASTELLANO, A. GIORDANO BRUNO: *Topological entropy for locally linearly compact vector spaces*. Topology Appl. 252 (2019), 112–144.
- [11] I. CASTELLANO, D. DIKRANJAN, D. FRENI, A. GIORDANO BRUNO, D. TOLLER: *Intrinsic entropy for generalised quasimetric semilattices*. J. Algebra Appl. 21 (2022), no. 10, 2250244, 24 pp.
- [12] D. DIKRANJAN, A. GIORDANO BRUNO: *Topological entropy and algebraic entropy for group endomorphisms*. Proceedings ICTA2011 Islamabad Pakistan July 4–10 2011, 133–214, Cambridge Scientific Publishers, 2012.
- [13] D. DIKRANJAN, A. GIORDANO BRUNO: *The Bridge Theorem for totally disconnected LCA groups*. Topology Appl. 169 (2014), 21–32.
- [14] D. DIKRANJAN, A. GIORDANO BRUNO: *Entropy on abelian groups*. Adv. Math. 298 (2016), 612–653.
- [15] D. DIKRANJAN, I. PROTASOV, N. ZAVA: *Hyperballeans of groups*. Topology Appl. 263 (2019), 172–198.
- [16] D. DIKRANJAN, A. GIORDANO BRUNO, H.-P. KÜNZI, N. ZAVA, D. TOLLER: *Generalized quasi-metric semilattices*. Topology Appl. 309 (2022), Paper No. 107916, 35 pp.
- [17] D. DIKRANJAN, A. GIORDANO BRUNO, L. SALCE: *Adjoint algebraic entropy*. J. Algebra 324 (2010), no. 3, 442–463.
- [18] D. DIKRANJAN, A. GIORDANO BRUNO, L. SALCE, S. VIRILI: *Intrinsic algebraic entropy*. J. Pure Appl. Algebra 219 (2015), no. 7, 2933–2961.
- [19] D. DIKRANJAN, B. GOLDSMITH, L. SALCE, P. ZANARDO: *Algebraic entropy for abelian groups*. Trans. Amer. Math. Soc. 361 (2009), no. 7, 3401–3434.
- [20] D. DIKRANJAN, I. PROTASOV, N. ZAVA: *Hyperballeans of groups*. Topology Appl. 263 (2019), 172–198.
- [21] I. EIDHAMMER, I. JONASSEN, W. R. TAYLOR: *Protein bioinformatics: An algorithmic approach to sequence and structure analysis*. John Wiley & Sons Ltd, West Sussex, England, 2004.
- [22] G. GIERZ, K. H. HOFMANN, K. KEIMEL, J. D. LAWSON, M. MISLOVE, D. S. SCOTT: *Continuous lattices and domains*. Encyclopedia Math. Appl., 93 Cambridge University Press, Cambridge, 2003.
- [23] A. GIORDANO BRUNO: *Adjoint entropy vs Topological entropy*. Topology Appl. 159 (2012), no. 9, 2404–2419.
- [24] A. GIORDANO BRUNO, M. SHLOSSBERG, D. TOLLER: *Algebraic entropy on strongly compactly covered groups*. Topology Appl. 263 (2019), 117–140.
- [25] A. GIORDANO BRUNO, S. VIRILI: *Topological entropy in totally disconnected locally compact groups*. Ergodic Theory Dynam. Systems 37 (2017), no. 7, 2163–2186.
- [26] B. GOLDSMITH, K. GONG: *On adjoint entropy of abelian groups*. Comm. Algebra 40 (2012), no. 3, 972–987.
- [27] J. GOUBAULT-LARRECQ: *Non-Hausdorff Topology and Domain Theory, Selected Topics in Point-Set Topology*. New Mathematical Monographs, Cambridge University Press, Cambridge, 2013.
- [28] F. HAUSDORFF: *Grundzüge der Mengenlehre*. Leipzig, 1914.
- [29] B. M. HOOD: *Topological entropy and uniform spaces*. J. London Math. Soc. (2) 8 (1974), 633–641.
- [30] A. N. KOLMOGOROV: *New metric invariants of transitive dynamical systems and automorphisms of Lebesgue spaces*. Doklady Akad. Nauk. SSSR 119 (1958), 861–864.
- [31] H.-P. KÜNZI, V. VAJNER: *Weighted Quasi-Metrics*. Papers on general topology and applications (Flushing, NY, 1992), 64–77, Ann. New York Acad. Sci., 728, New York Acad. Sci., New York, 1994.
- [32] M. O’KEEFFE, S. ROMAGUERA, M. SCHELLEKENS: *Norm-weightable Riesz spaces and the dual complexity space*. Electronic Notes in Theoretical Computer Science 74 (2003): 105–121.
- [33] S. J. O’NEILL: *Partial metrics, valuations, and domain theory*. Papers on general topology and applications (Gorham, ME, 1995), 304–315, Ann. New York Acad. Sci., 806, New York Acad. Sci., New York, 1996.
- [34] S. G. MATTHEWS: *Partial metric topology*. Papers on general topology and applications (Flushing, NY, 1992), 183–197, Ann. New York Acad. Sci., 728, New York Acad. Sci., New York, 1994.
- [35] Y. NAKAMURA: *Entropy and semivaluations on semilattices*. Kodai Math. Sem. Rep. 22 (1970), 443–468.
- [36] S. ROMAGUERA, M. SCHELLEKENS: *Quasi-metric properties of complexity spaces*. Topology and its Applications 98, 311–322, 1999.
- [37] S. ROMAGUERA, M. SCHELLEKENS: *Quasi-metric properties of Complexity Spaces*. Topology Appl. 98, 311–322, 1999.
- [38] L. SALCE, P. ZANARDO: *A general notion of algebraic entropy and the rank entropy*. Forum Math. 21 (2009), no. 4, 579–599.
- [39] M. P. SCHELLEKENS: *The Smyth Completion: A Common Foundation for Denotational Semantics and Complexity Analysis*, in: proc. MFPS 11, Electronic Notes in Theoretical Computer Science, Vol. I, Elsevier, 211–232, 1995.
- [40] M. P. SCHELLEKENS: *On upper weighted spaces*. Proc. 11th Summer Conference on General Topology and Applications. Ann. New York Acad. Sci 806, 348–363.
- [41] M. P. SCHELLEKENS: *Extendible spaces*. Appl. Gen. Topol. 3 (2002), no. 2, 169–184.
- [42] M. P. SCHELLEKENS: *The correspondence between partial metrics and semivaluations*. Theoret. Comput. Sci. 315 (2004), no. 1, 135–149.
- [43] D. SCOTT: *A type theoretic alternative to ISWIM, CUCH, OWHY*, manuscript, University of Oxford (1969).
- [44] D. SCOTT: *Outline of a mathematical theory of computation*. In: 4th Annual Princeton Conference on Information Sciences and Systems (1970), 169–176.
- [45] C. E. SHANNON: *A Mathematical Theory of Communication*. Bell System Technical Journal, 27 (3) (1948), 379–423.
- [46] Y. G. SINAI: *On the concept of entropy of a dynamical system*. Doklady Akad. Nauk. SSSR 124 (1959), 786–781.
- [47] S. VIRILI: *Entropy for endomorphisms of LCA groups*. Topology Appl. 159 (2012), no. 9, 2546–2556.
- [48] H. WEBER: *Uniform lattices I: a generalization of topological Riesz spaces and topological Boolean rings*. Ann. Mat. Pura Appl. (4) 160 (1991), 347–370 (1992).
- [49] H. WEBER: *Uniform lattices II: order continuity and exhaustivity*. Ann. Mat. Pura Appl. (4) 165 (1993), 133–158.
- [50] M. D. WEISS: *Algebraic and other entropies of group endomorphisms*. Math. Systems Theory 8 (1974/75), no. 3, 243–248.
- [51] W. A. WILSON: *On Quasi-Metric Spaces*. Amer. J. Math. 53 (1931), no. 3, 675–684.
- [52] N. ZAVA: *Generalisations of coarse spaces*. Topology Appl. 263 (2019), 230–256.