

An Introduction to Totally Disconnected Locally Compact Groups and Their Finiteness Conditions

*Original*

An Introduction to Totally Disconnected Locally Compact Groups and Their Finiteness Conditions / Castellano, Ilaria (LONDON MATHEMATICAL SOCIETY LECTURE NOTE SERIES). - In: Modern Trends in Algebra and Representation Theory (Series Number 486) / David Jordan, Nadia Mazza, Sibylle Schroll. - [s.l.] : Cambridge University Press, 2023. - ISBN 9781009093750. [10.1017/9781009093750.012]

*Availability:*

This version is available at: 11583/3003349 since: 2025-09-25T13:38:36Z

*Publisher:*

Cambridge University Press

*Published*

DOI:10.1017/9781009093750.012

*Terms of use:*

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

Cambridge University Press postprint/Author's Accepted Manuscript

(Article begins on next page)

An introduction to  
totally disconnected locally compact groups  
and their finiteness conditions

Ilaria Castellano

**Abstract**

Here we collect the notes of the expository lectures that have been delivered for the *LMS Autumn Algebra School*, 21-25 September 2020. The aim is to introduce beginning Ph.D. students to the theory of totally disconnected locally compact groups. The exposition includes basic properties of topological groups, a proof of van Dantzig's theorem and some of the most popular (and trouble-free) examples of totally disconnected locally compact groups. The focus is on totally disconnected locally compact groups that satisfy (homological) finiteness conditions with more emphasis on *compact generation*: for compactly generated totally disconnected locally compact groups, the notion of *Cayley-Abels graph* permits us to deal with the topological group as a geometric object.



# Contents

<b>1</b>	<b>An introduction to TDLC-groups and their finiteness conditions</b>	<i>page</i> 1
1.1	Introduction	1
1.2	Preliminaries on topological groups	4
1.2.1	Warming-up	4
1.2.2	Profinite groups	8
1.2.3	Locally compact groups	11
1.3	Totally disconnected locally compact groups	14
1.3.1	van Dantzig's theorem	14
1.3.2	Consequences of van Dantzig's theorem	14
1.3.3	Examples of TDLC-groups	16
1.4	Finiteness properties for TDLC-groups	21
1.4.1	Compact generation and presentation	21
1.4.2	The Cayley-Abels graphs	22
1.4.3	Finiteness conditions in higher dimension	26
1.5	Willis' theory of TDLC-groups: a sketch	30
1.5.1	Scale function and tidy subgroups	31
1.5.2	Comments on simple TDLC-groups	32
	<i>References</i>	35



# 1

## An introduction to totally disconnected locally compact groups and their finiteness conditions

### 1.1 Introduction

The class of locally compact groups generalises discrete and Lie groups. Locally compact groups came to light in the first half of 20th century, and since then they have played a central role among topological groups. The 20th century has been characterised by intense activity on the structure theory of many algebraic objects, e.g., simple finite groups, fields, and rings with ascending chain condition (a.c.c.). What about the general structure of locally compact groups? A basic strategy to understand the structure of a locally compact group  $G$  is to split it into smaller groups: let  $G_0$  be the largest connected subset of  $G$  containing the identity element, which is a closed subgroup (see Fact 1.6), and produce the short exact sequence

$$1 \rightarrow G_0 \rightarrow G \rightarrow G/G_0 \rightarrow 1,$$

where  $G_0$  is a **connected** locally compact group and  $G/G_0$  is a **totally disconnected** locally compact group (i.e., the connected components of  $G/G_0$  are reduced to singletons). Therefore,  $G$  can be regarded as an extension of its connected component by the totally disconnected piece  $G/G_0$ . It follows that, at least in theory, questions about the structure of locally compact groups may be dealt with by treating separately the cases where  $G$  is connected and where  $G$  is totally disconnected and then combining the two answers. With the solution of Hilbert's fifth problem, our understanding of connected locally compact groups has significantly increased: they can be approximated by Lie groups (see Theorem 1.16). Therefore, the contemporary structure problem on locally compact groups concerns the class of totally disconnected locally compact groups.

## 2 An introduction to TDLC-groups and their finiteness conditions

The first part of the notes introduces the objects under investigation: topological groups that are locally compact and totally disconnected (shortly, TDLC-groups). In particular, it includes basic properties of topological groups, a proof of van Dantzig's theorem, which is the fundamental structure theorem of TDLC-groups, and several examples.

The study of this class of topological groups can be made more manageable by dividing the infinity of objects under investigation into classes of types with "similar structure": we focus on TDLC-groups satisfying some finiteness conditions. The most common finiteness condition for (totally disconnected) locally compact groups is *compact generation* - i.e., the topological group is algebraically generated by a compact subset - which naturally generalises the notion of finite generation that has been widely (and fruitfully) used in group theory to study infinite groups. Since every TDLC-group is a directed union of compactly generated open subgroups (see Fact 1.29), one can confine the investigation to the compactly generated case without losing too much information (at least from a local perspective).

All compactly generated TDLC-groups fall in the class of automorphism groups of locally finite connected graphs (see § 1.3.3). Indeed, Abels (1973/74) proved that, by using van Dantzig's theorem, we can always construct a locally finite connected graph  $\Gamma$  on which the group  $G$  acts vertex-transitively and with compact open stabilisers, the so-called *Cayley-Abels graph of  $G$*  (see § 1.4.2). To be precise, every compactly generated TDLC-group comes with a family of Cayley-Abels graphs which satisfy the striking property of being quasi-isometric each-other: the Cayley-Abels graph  $\Gamma$  of a compactly generated TDLC-group is unique up to quasi-isometry. Therefore, by analogy with the theory of finitely generated groups and their Cayley graphs, we can study a compactly generated TDLC-group as a geometric object and all the properties of  $\Gamma$  that are invariant up to quasi-isometry become *geometric group invariants*<sup>1</sup> of  $G$ . It leads to a new line of research in topological group theory, *geometric group theory for compactly generated TDLC-groups*, which can be traced back to the work of Krön and Möller (2008).

As a consequence, TDLC-groups need to be viewed as simultaneously geometric objects and topological objects. Therefore, the interaction between the local structure (i.e., the topological one) and the large-scale structure (i.e., the geometric one) becomes important. Since profinite

<sup>1</sup> The number of ends, growth and hyperbolicity are examples of geometric group invariants of a TDLC-group  $G$ .

groups are trivial as geometric groups and discrete groups are trivial as topological groups, it is not surprising that the profinite groups and the discrete groups constitute the atomic pieces in the theory of TDLC-groups and that we are curious to understand to what extent well-known results about the geometry of infinite discrete groups find an analogue in such a topological context.

The final part of the fourth chapter attempts to introduce the reader to some homological finiteness conditions for TDLC-groups that generalise compact generation in higher dimensions. Since these notes are meant to be on the non-specialist level, we only provide definitions (without details) and references (where the details are), letting the reader decide how deep to dig into the subject.

The conclusive part briefly introduces the seminal work of Willis (1994, 2001a) which was a fundamental breakthrough in the theory of TDLC-groups after several years of stillness.

Willis' theory gave start to the research interest we now benefit from.

**Pre-requisites:** The reader is supposed to have mastered linear algebra, point-set topology and basic notions from group theory. These notes aim to help beginning Ph.D. students moving the first steps towards the theory of TDLC-groups. Therefore, this text is (supposed to be) accessible to non-specialists and self-contained (most of the results are proved in these notes) but, as with any advanced topic, some of the results are stated without proofs. In such a case, references are provided.

The interested reader can also have a look at a few conference proceedings that collect part of the progress made with the general theory of TDLC-groups and include some interesting open problems; see for example Caprace and Monod (2018) and Wood et al. (2018).

**Notation:** We denote by

- $\mathbb{N}$  the set of natural numbers  $\{0, 1, 2, \dots\}$ ,
- $\mathbb{Z}$  the ring of rational integers and  $\mathbb{Q}$  the ring of rational numbers,
- $\mathbb{R}$  the field of real numbers and  $\mathbb{C}$  the field of complex numbers,
- $\mathbb{R}_+$  the subset of non-negative real numbers,
- If  $R$  is a commutative ring with unit,  $R^\times$  stands for its multiplicative group of units. For example,  $\mathbb{R}_+^\times$  is the group of positive real numbers.

## 1.2 Preliminaries on topological groups

A complete and detailed introduction to the theory of topological groups can be found in several textbooks; for example, Hewitt and Ross (2012). For convenience, all topological spaces appearing below are assumed to satisfy the Hausdorff separation axiom<sup>2</sup>.

### 1.2.1 Warming-up

**Definition 1.1** (Topological group) A **topological group** is a group  $(G, \cdot)$  which is also a topological space such that the following maps are continuous:

- the group operation

$$\cdot : G \times G \rightarrow G, \quad (x, y) \mapsto x \cdot y, \quad \forall x, y \in G,$$

where  $G \times G$  is endowed with the product topology;

- the inversion map

$$-^1 : G \rightarrow G, \quad x \mapsto x^{-1}, \quad \forall x \in G.$$

If the underlying group  $G$  is cyclic (resp., abelian, nilpotent, etc.), the topological group  $G$  is also called cyclic (resp. abelian, nilpotent, etc.). The additive notation  $(G, +)$  can describe some topological groups but only in the case when the topological group is abelian.

*Remark* Similarly, one has **topological rings** and **topological fields**<sup>3</sup>.

Every abstract group  $G$  can be viewed as a topological group if given the discrete topology. In such a case,  $G$  is called a **discrete group** and, since these notes concern topological groups, we will often refer to (abstract) groups as discrete groups.

**Exercise 1.2.1** Let  $G$  be a topological group. Prove that:

1. If a subgroup  $H \leq G$  is open, then it is also closed.
2. If a subgroup contains an open set then it is open.
3. A connected subset  $C \ni x$  is contained in the intersection of all clopen<sup>4</sup> subsets  $O \ni x$  of  $G$ .
4. If  $G$  is connected, then the only open subgroup of  $G$  is  $G$  itself.
5. Every quotient map of a topological group is open.

<sup>2</sup> Totally disconnected locally compact groups are necessarily Hausdorff.

<sup>3</sup> By “field” we always mean “commutative field”.

<sup>4</sup> A set which is both closed and open.

6. For  $H \leq G$  normal, the quotient group  $G/H$  is discrete iff  $H$  is open. In particular, the group  $G$  is discrete iff the point  $1_G$  is isolated, i.e., the singleton  $\{1_G\}$  is open.

Other (less trivial) examples of a topological group are provided by the additive group  $(\mathbb{R}, +)$  of the reals equipped with the usual topology, its subgroups  $\mathbb{Z}$  and  $\mathbb{Q}$  (with the subspace topology) and the quotient  $\mathbb{T} := \mathbb{R}/\mathbb{Z}$  (with the quotient topology). These extend to all powers  $(\mathbb{R}^d, +)$ , and so  $(\mathbb{C}, +)$ , because it can be easily proved that products of topological groups are again topological groups. Moreover, if  $R$  is a topological ring, then the ring  $M(n, R)$  of all  $n \times n$  matrices with entries in  $R$  is a topological ring if endowed with the product topology of  $R^{n \times n}$ .

**Exercise 1.2.2** Let  $R$  be a commutative topological ring such that inversion is continuous on the set of invertible elements (for example, if  $R$  is a topological field). Prove that

1. the group  $(GL(n, R), \cdot)$  of all invertible  $n \times n$  matrices with entries in  $R$  is a topological group (notice that  $GL(n, R)$  is a subset of  $M(n, R)$  but not a subgroup; in this case Cramer's rule helps with the continuity of the inversion map);
2.  $SL(n, R) = \{M \in GL(n, R) \mid \det(M) = 1\}$  is closed in  $GL(n, R)$ .

A topology  $\tau$  on the group  $G$  such that the space  $(G, \tau)$  is a topological group is called a **group topology** on  $G$ . Obviously, a topology  $\tau$  on  $G$  is a group topology if, and only if, the map  $G \times G \rightarrow G$ ,  $(x, y) \mapsto xy^{-1}$ , is continuous for all  $x, y \in G$ . Notice that, for every  $g \in G$ , the **left translation**  $x \mapsto gx$ , the **right translation**  $x \mapsto xg$ , as well as the **conjugation**  $x \mapsto gxg^{-1}$  are continuous; in other words, every topological group is a homogeneous topological space.

**Exercise 1.2.3** Prove the assertions above.

As a consequence, the topology of  $G$  is determined by a neighbourhood basis<sup>5</sup> of the identity: a family  $\{U_\alpha\}_{\alpha \in I}$  of arbitrarily small neighbourhoods of  $1_G$  determines the family  $\{gU_\alpha\}_{\alpha \in I}$  of arbitrarily small neighbourhoods of any other group element  $g$ .

**Example 1.2** One can define group topologies on  $G$  by declaring well-behaved collections of subsets to be neighbourhood basis at  $1_G$  (see Bourbaki (1998)):

<sup>5</sup> A family  $\mathcal{B}$  of neighbourhoods of the point  $x$  is said to be a *neighbourhood basis* of  $x$  if for every neighbourhood  $U$  of  $x$  there exists  $V \in \mathcal{B}$  contained in  $U$ .

6 *An introduction to TDLG-groups and their finiteness conditions*

- the **profinite topology** is determined by the family of all normal subgroups of finite index of  $G$ ;
- the **pro- $p$  topology** is determined, for a prime  $p$ , by all normal subgroups of  $G$  of finite index that is a power of  $p$ ;
- the  **$p$ -adic topology** is determined, for a prime  $p$ , by the family  $\{U_n\}_{n \in \mathbb{N}}$  of normal subgroups of  $G$ , where  $U_n$  is generated by the powers  $\{g^{p^n} \mid g \in G\}$ .

**Example 1.3** (Absolute values on fields) An **absolute value** on a field  $\mathbb{K}$  is a function  $|\_|\_ : \mathbb{K} \rightarrow \mathbb{R}$  satisfying

- |1|.  $|x| \geq 0$  for every  $x \in \mathbb{K}$ , and  $|x| = 0$  if and only if  $x = 0$ ,
- |2|.  $|xy| = |x||y|$ , for all  $x, y \in \mathbb{K}$ ,
- |3|.  $|x + y| \leq |x| + |y|$ , for all  $x, y \in \mathbb{K}$ .

If the absolute value  $|\_|\_$  satisfies the stronger condition

- |3̄|.  $|x + y| \leq \max\{|x|, |y|\}$ , for all  $x, y \in \mathbb{K}$

it is said to be **non-Archimedean**, otherwise it is **Archimedean**. For example, the usual absolute value on  $\mathbb{R}$  is Archimedean.

It is clear that  $d(x, y) = |x - y|$  gives  $\mathbb{K}$  a structure of a metric space, and the topology for which

$$B_\varepsilon(0) := \{x \in \mathbb{K} \mid |x| < \varepsilon\}, \quad \varepsilon > 0,$$

form a basis of neighbourhoods of 0 is a field topology.

**Exercise 1.2.4** Let  $\mathbb{K}$  be a field equipped with the absolute value  $|\_|\_$ . The field topology defined on  $\mathbb{K}$  by  $|\_|\_$  is discrete iff  $|x| = 1$  for all  $x \neq 0$ .

**Example 1.4** (The field of  $p$ -adic numbers) For a prime  $p$ , the  **$p$ -adic absolute value** on  $\mathbb{Q}$  is defined as  $|x|_p = p^{-n}$ , for every  $x \in \mathbb{Q}$ , where  $n$  is the unique integer such that  $x = p^n(\frac{a}{b})$  and neither of the integers  $a$  and  $b$  is divisible by  $p$  (with the convention,  $|0|_p = 0$ ). It is an example of non-archimedean absolute value. The  **$p$ -adic metric**  $d_p(x, y) = |x - y|_p$  induces a field topology on  $\mathbb{Q}$  which is called the  **$p$ -adic topology** of  $\mathbb{Q}$ ; see Example 1.2. As with the topology on  $\mathbb{Q}$  inherited from  $\mathbb{R}$ , the metric space  $(\mathbb{Q}, d_p)$  is not complete (i.e., not every Cauchy sequence converges in  $(\mathbb{Q}, d_p)$ ). Let  $\mathbb{Q}_p$  denote the completion of  $\mathbb{Q}$  with respect to the  $p$ -adic metric. The field  $\mathbb{Q}_p$  has characteristic zero and it is called the **field of  $p$ -adic numbers**. The metric  $d_p$ , and so the  $p$ -adic topology, extends from  $\mathbb{Q}$  to  $\mathbb{Q}_p$  and yields a topological field; see (Palmer, 1994, § 12.3.4).

**The identity component of  $G$ .** On a topological space  $X$  one defines the equivalence relation  $\sim$  as follows:  $x \sim y$  if there exists a connected subspace  $C \subseteq X$  such that  $x, y \in C$ . Each equivalence class is a maximal connected subspace which is called a **connected component** of  $X$ .

**Definition 1.5** Given a topological group  $G$ , the connected component containing the identity is the **identity component** of  $G$ , and it is denoted by  $G_0$ . Clearly,  $G_0$  is the union of all connected subspaces of  $G$  containing  $1_G$ , and the topological group  $G$  is connected if, and only if,  $G = G_0$ . A topological group is said to be **totally disconnected** if the identity is its own connected component, that is,  $G_0 = \{1_G\}$ .

Notice that, for every  $g \in G$ , the set  $gG_0 = G_0g$  is nothing but the connected component containing  $g$  because continuous maps preserve connectedness and translations are continuous. As a consequence, one has the following important fact.

**Fact 1.6** *Given a topological group, the identity component is a closed characteristic subgroup.*

*Proof* The subspace  $G_0$  is closed since the closure of a connected subspace is still connected.

For every  $x \in G_0$ , the translate  $x^{-1}G_0 \ni 1_G$  is connected. It follows that  $x^{-1} \in x^{-1}G_0 \subseteq G_0$ , i.e.,  $G_0$  is closed under taking inverses. On the other hand, for all  $x, y \in G_0$ , one has  $xy \in xG_0$  but  $xG_0$  is the connected component of  $x$  which is in the same connected component as  $1_G$  (because  $x \sim 1_G$ ), i.e.,  $xy \in xG_0 = G_0$  and  $G_0$  is a subgroup.

Finally,  $G_0$  is characteristic (and in particular normal) because continuous homomorphisms preserve connectedness.  $\square$

**Proposition 1.7** *Let  $G$  be a topological group and let  $G_0$  be the identity component. Then  $G/G_0$  is a totally disconnected group.*

*Proof* Let  $\pi: G \rightarrow H := G/G_0$  be the quotient map. One has to prove that  $H_0 = 1$ , i.e.,  $\pi^{-1}(H_0) = G_0$ . By the maximality of  $G_0$ , it suffices to prove that  $\pi^{-1}(H_0) \supseteq G_0$  is connected. To this end, suppose by contradiction  $\pi^{-1}(H_0)$  is a disjoint union  $C_1 \cup C_2$  of non-empty closed subsets of  $G$ . Since  $G_0$  is connected, for every  $g \in \pi^{-1}(H_0)$ , either  $gG_0 \subseteq C_1$  or  $gG_0 \subseteq C_2$ . Therefore,  $C_1$  and  $C_2$  are unions of  $G_0$ -cosets. Consequently,  $H_0$  is the disjoint union of non-empty closed subsets, contradiction.  $\square$

### 1.2.2 Profinite groups

For a complete introduction to the realm of profinite groups the reader is referred to Ribes and Zalesskii (2000); Wilson (1998) and (Caprace and Monod, 2018, Chapter 3). We recall only the definition and provide some easy examples.

A **directed poset**  $(I, \preceq)$  is a set  $I$  with a binary relation  $\preceq$  satisfying:

- (DP1)  $i \preceq i$ , for  $i \in I$ ;
- (DP2)  $i \preceq j$  and  $j \preceq k$  imply  $i \preceq k$ , for  $i, j, k \in I$ ;
- (DP3)  $i \preceq j$  and  $j \preceq i$  imply  $i = j$ , for  $i, j \in I$ ; and
- (DP4) if  $i, j \in I$ , there exists some  $k \in I$  such that  $i \preceq k$  and  $j \preceq k$ .

An **inverse system of topological groups over  $I$**  consists of a family  $\{G_i \mid i \in I\}$  of topological groups together with continuous group morphisms  $\varphi_{ij}: G_i \rightarrow G_j$ , defined whenever  $j \preceq i$ , such that the diagrams

$$\begin{array}{ccc} G_i & \xrightarrow{\varphi_{ik}} & G_k \\ & \searrow \varphi_{ij} & \nearrow \varphi_{jk} \\ & G_j & \end{array}$$

commutes whenever  $k \preceq j \preceq i$ . In addition, we assume that  $\varphi_{ii}$  is the identity morphism for every  $i \in I$ . A projective system of topological groups is said to be **surjective** if every morphism  $\varphi_{ij}$  is surjective. A family of continuous group morphisms  $\varphi_i: G \rightarrow G_i$  is said to be **compatible** with the inverse system  $(G_i, \varphi_{ij}, I)$  if, for every  $i \preceq j$ , the diagram

$$\begin{array}{ccc} G & \xrightarrow{\varphi_i} & G_i \\ & \searrow \varphi_j & \nearrow \varphi_{ji} \\ & G_j & \end{array}$$

commutes. A topological group  $G$  together with a compatible family  $\{\varphi_i: G \rightarrow G_i\}_{i \in I}$  of continuous morphisms is an **inverse limit** of the inverse system  $(G_i, \varphi_{ij}, I)$  if the following universal property is satisfied: for every topological group  $\tilde{G}$  together with a compatible family  $\{\psi_i: \tilde{G} \rightarrow G_i\}_{i \in I}$  of continuous group morphisms, there exists a unique

continuous group morphism  $\psi: \tilde{G} \rightarrow G$  such that the diagram

$$\begin{array}{ccc} \tilde{G} & \overset{\psi}{\dashrightarrow} & G \\ & \searrow \psi_i & \swarrow \varphi_i \\ & G_i & \end{array}$$

commutes for every  $i \in I$ . In such a case, we denote the inverse limit by

$$G = \varprojlim_{i \in I} (G_i, \varphi_{ij}),$$

and call the maps  $\varphi_i: G \rightarrow G_i$  *projection morphisms*. If the family  $\varphi_{ij}$  is clear from the context, we use simply  $G = \varprojlim_{i \in I} G_i$ .

**Proposition 1.8** ((Ribes and Zalesskii, 2000, Proposition 1.1.1)) *Let  $(G_i, \varphi_{ij}, I)$  be an inverse system of topological groups over a directed poset  $I$ . Then the following hold:*

- (a) *There exists an inverse limit of the inverse system  $(G_i, \varphi_{ij}, I)$ ;*
- (b) *This limit is unique in the following sense: if  $(G, \varphi_i)$  and  $(H, \psi_i)$  are two limits of  $(G_i, \varphi_{ij}, I)$ , then there is a unique topological isomorphism  $\phi: G \rightarrow H$  such that  $\psi_i \phi = \varphi_i$  for each  $i \in I$ .*

In particular, the inverse limit  $(G, \varphi_i)$  can be constructed as follows:

- $G := \{(g_i)_{i \in I} \in \prod_{i \in I} G_i \mid \varphi_{ij}(g_j) = g_i \text{ if } j \succeq i\}$ ;
- $\varphi_i: G \rightarrow G_i$  is the restriction of the projection  $\prod_{i \in I} G_i \rightarrow G_i$ ;
- the group topology on  $G$  is the subspace topology inherited by the product topology of  $\prod_{i \in I} G_i$  (and  $G$  turns out to be a closed subset).

**Fact 1.9** ((Ribes and Zalesskii, 2000, Lemma 1.1.2)) *If  $(G_i, \varphi_{ij}, I)$  is an inverse system of topological groups, then  $\varprojlim_{i \in I} G_i$  is a closed subgroup of the product  $\prod_{i \in I} G_i$ .*

**Definition 1.10** A **profinite group**  $G$  is the inverse limit  $\varprojlim_{i \in I} G_i$  of a surjective inverse system  $(G_i, \varphi_{ij}, I)$  of finite groups  $G_i$ , where each finite group  $G_i$  is assumed to have the discrete topology.

For a profinite group  $G$ , a neighbourhood basis at  $1_G$  is  $\{\ker(\varphi_i)\}_{i \in I}$ , where  $\varphi_i: G \rightarrow G_i$  are the canonical projection homomorphisms.

**Fact 1.11** *A profinite group  $G$  is compact and totally disconnected.*

*Proof* It easily follows from the fact that  $G$  is a closed subset of the product of finite groups (see Fact 1.9). To see this, one needs to recall the following basic properties:

10 *An introduction to TDLG-groups and their finiteness conditions*

- closed subspaces and products of compact (resp. totally disconnected) spaces are compact (resp. totally disconnected);
- a finite group is a compact discrete group; in particular it is totally disconnected.

□

**Example 1.12** (a) Let  $R$  be a profinite commutative ring with unit. Then the following groups (with topologies naturally induced from  $R$ ) are profinite groups:  $R^\times$ ,  $GL(n, R)$  and  $SL(n, R)$ .

- (b) Consider the natural numbers  $I = \mathbb{N}$ , with the usual partial ordering, and the group of integers  $\mathbb{Z}$ . Form the inverse system  $\{\mathbb{Z}/n\mathbb{Z}, \phi_{nm}\}$ , where the map  $\phi_{nm}: \mathbb{Z}/n\mathbb{Z} \rightarrow \mathbb{Z}/m\mathbb{Z}$  is the natural projection for  $m \leq n$ . The inverse limit produces the profinite group  $\widehat{\mathbb{Z}}$ , which can be identified with the set of equivalence classes of tuples of integers

$$\{(\overline{x_1, x_2, x_3, \dots}) \mid x_n \in \mathbb{Z}, \forall n \in \mathbb{N}, \text{ and } x_m = x_n \pmod m \text{ whenever } m|n\}.$$

Note that  $\widehat{\mathbb{Z}}$  naturally inherits a structure of profinite ring from the finite rings  $\mathbb{Z}/n\mathbb{Z}$ . The ring  $\widehat{\mathbb{Z}}$  is the **profinite completion** of  $\mathbb{Z}$ .

- (c) Let  $p$  be a prime and form the profinite group defined by the inverse limit over the system  $\{\mathbb{Z}/p^n\mathbb{Z}, \psi_{mn}\}$  given by the canonical projections. It is the **pro- $p$  completion** of  $\mathbb{Z}$  and it is topologically isomorphic to the ring of  $p$ -adic integers  $\mathbb{Z}_p$ : it suffices to prove that  $\mathbb{Z}_p$  is the inverse limit of its quotients  $\mathbb{Z}_p/p^n\mathbb{Z}_p$  (where the family  $\{p^n\mathbb{Z}_p\}_{n \in \mathbb{N}}$  is the neighbourhood basis at 0 in the group of  $p$ -adic integers) and that each  $\mathbb{Z}_p/p^n\mathbb{Z}_p$  is isomorphic to the finite group  $\mathbb{Z}/p^n\mathbb{Z}$ .

The set of the elements of  $\mathbb{Z}_p$  can be then identified with the set of all equivalence classes of sequences  $(a_1, a_2, a_3, \dots)$  of natural numbers such that  $a_m = a_n \pmod{p^m}$ , whenever  $m \leq n$ .

**Exercise 1.2.5** 1. Let  $\{G_i \mid i \in I\}$  be a collection of finite groups. Is the direct product a profinite group?

2. Consider the natural numbers  $I = \mathbb{N}$ , with the usual partial ordering, and form the constant inverse system  $\{\mathbb{Z}, id\}$ . Compute the limit.
3. Let  $G$  be a profinite group.

- (a). A closed normal subgroup  $H \leq G$  is open iff it has finite index.
- (b). Every open subgroup  $H$  of  $G$  contains a subgroup  $H_G$  that is normal and open in  $G$ . (Hint:  $H_G = \bigcap_{g \in G} gHg^{-1}$ )

### 1.2.3 Locally compact groups

An arbitrary topological space  $X$  is locally compact if every point admits a compact neighbourhood. A topological group  $G$  is **locally compact** if  $1_G$  admits a compact neighbourhood. The additive group  $(\mathbb{R}, +)$  with its usual topology is a locally compact, non-compact, abelian group. Clearly, the multiplicative group  $(\mathbb{R}^\times, \cdot)$  is also locally compact (here  $\mathbb{R}^\times$  carries the induced topology) and the same holds for the groups  $(\mathbb{C}, +)$  and  $(\mathbb{C}^\times, \cdot)$ . On the other hand,  $\mathbb{Q}$  is not locally compact with the topology inherited by  $\mathbb{R}$  (see Proposition 1.15): local compactness is not inherited by subgroups; see Exercise 1.2.6.

Different examples of locally compact groups are discrete groups and profinite groups. If  $\mathbb{T} = \mathbb{R}/\mathbb{Z}$  is the circle group<sup>6</sup>, then Tychonov's theorem yields that every power  $\mathbb{T}^I$  of  $\mathbb{T}$  is again compact and, in particular, locally compact. This is actually the most general example of a compact abelian group: every compact abelian group is isomorphic to a closed subgroup of a power of  $\mathbb{T}$ ; see (Dikranjan, 2018, Corollary 11.2.2).

**Example 1.13** (Local fields) Non-discrete locally compact fields have been classified by Van Dantzig (1931). A non-discrete locally compact field is either connected or totally disconnected. A non-discrete locally compact field is connected if and only if it is Archimedean (see Example 1.3), and then isomorphic to either  $\mathbb{R}$  or  $\mathbb{C}$ . Non-Archimedean locally compact fields are called **local fields**<sup>7</sup>. For further details see (Palmer, 1994, § 12.3.4) and references there.

- Exercise 1.2.6**
1. A closed subgroup of a locally compact group is locally compact. The closure condition is necessary, see Proposition 1.15.
  2. If  $R$  is a locally compact ring and  $n$  is a natural number, then  $R^{n \times n}$  is a locally compact ring.
  3. Every quotient of a locally compact group is locally compact.
  4. The product of a finite family of locally compact groups is locally compact (for infinite products to be locally compact is necessary the condition “all but a finite number of factors are compact”).
  5. If  $\mathbb{K}$  is a topological field, then  $GL(n, \mathbb{K})$  is open in  $\mathbb{K}^{n \times n}$ . Consequently,  $(GL(n, \mathbb{K}), \cdot)$  is locally compact exactly if  $\mathbb{K}$  is.
  6. If  $\mathbb{K}$  is a locally compact field, then  $(SL(n, \mathbb{K}), \cdot)$  is locally compact.

<sup>6</sup> The circle group is compact because it is the image of the compact subspace  $[0, 1] \subset \mathbb{R}$ .

<sup>7</sup> Some authors define a local field to be any commutative non-discrete locally compact field.

**Proposition 1.14** *A locally compact countable group is discrete.*

*Proof* Recall that a *Baire space* is a topological space with the property that for each countable collection of open dense sets  $\{U_n\}_{n \in \mathbb{N}}$  their intersection  $\bigcap_{n \in \mathbb{N}} U_n$  is dense. By the Baire category theorem - see (Hofmann, 1980, Theorem A) and references there - every locally compact group is a Baire space. Let  $G$  be a non-discrete locally compact group; in particular, each singleton  $\{g\}$  is closed but not open in  $G$ . If  $G$  is countable, then

$$G = \{g_1, \dots, g_n, \dots\} = \bigcup_{n \in \mathbb{N}} (G \setminus \{g_n\})$$

but  $\bigcap_{n \in \mathbb{N}} (G \setminus \{g_n\}) = \emptyset$ , contradiction. □

**Proposition 1.15** ((Hewitt and Ross, 2012, Theorem 5.11)) *If a subgroup  $H$  of a topological group  $G$  is locally compact, then it is closed.*

Since the identity component  $G_0$  is a closed normal subgroup of the locally compact group  $G$  (see Fact 1.6), one can form the quotient group  $G/G_0$ . By Exercise 1.2.6(3),  $G/G_0$  is locally compact. Moreover, the locally compact group  $G/G_0$  is totally disconnected by Proposition 1.7. Therefore, one has the short exact sequence

$$1 \rightarrow G_0 \rightarrow G \rightarrow G/G_0 \rightarrow 1 \tag{1.1}$$

and every locally compact group  $G$  is an extension of a TDLG-group  $G/G_0$  by the connected locally compact group  $G_0$ , which can be approximated by Lie groups.

**Theorem 1.16** *Every connected locally compact group is inverse limit of Lie groups.*

The latter description is obtained as consequence of the following fundamental result.

**Theorem 1.17** (Gleason-Yamabe Theorem) *Let  $G$  be a locally compact group. For any open neighbourhood  $U$  of the identity there exists an open subgroup  $G'$  of  $G$  and a compact normal subgroup  $K$  of  $G'$  in  $U$  such that  $G'/K$  is isomorphic to a Lie group.*

We omit the proof of Gleason-Yamabe theorem but the reader is referred to (Tao, 2014, Theorem 1.1.13).

*Proof of Theorem 1.16* Let  $G$  be connected and locally compact. In order to construct a projective system of Lie groups, let  $\mathcal{N}$  be the set of all

open neighbourhoods  $U$  of the identity  $1_G$ . By Exercise 1.2.1(4), the only open subgroup  $G'$  of  $G$  is the whole group  $G$ . Therefore, for every  $U \in \mathcal{N}$ , there exists a compact normal subgroup  $K_U$  of  $G$  such that  $G/K_U$  is isomorphic to a Lie group. Let  $\mathcal{K} = \{K_U \mid U \in \mathcal{N}\}$  be indexed by inverse inclusion. In particular,  $(\mathcal{K}, \supseteq)$  is a directed poset; see (Hofmann and Morris, 2007, Glöckner's lemma, p. 148). Whenever  $M \supseteq N$  in  $\mathcal{K}$ , one has the continuous morphism  $\varphi_{NM}: G/N \rightarrow G/M$  given by  $gN \mapsto gM$  and  $(G_N, \varphi_{NM}, \mathcal{K})$  is a surjective inverse system of Lie groups. By the universal property of the inverse limit, there exists a continuous morphism  $\gamma: G \rightarrow \varprojlim (G_N, \varphi_{NM}, \mathcal{K})$ ; namely,  $\gamma(g) = (gN)_{N \in \mathcal{K}}$ . It is clear that the kernel of  $\gamma$  is given by  $\bigcap \{N \mid N \in \mathcal{K}\}$  which is trivial since the elements of  $\mathcal{K}$  can be arbitrarily small (see the Gleason-Yamabe theorem above). Thus one needs to prove that  $\gamma$  is open and surjective but it follows by (Hofmann and Morris, 2007, Theorem 1.33) and the fact that locally compact groups are always complete (see, for example, (Hofmann and Morris, 2007, Remark 1.31)).  $\square$

As Tao (2011) writes on his blog: *this theorem asserts the “mesoscopic” structure of a locally compact group (after restricting to an open subgroup  $G'$  to remove the macroscopic structure, and quotienting out by  $K$  to remove the microscopic structure) is always of Lie type.*

*Remark* (Hilbert's fifth problem) During the International Congress of Mathematicians (Paris, 1900) D. Hilbert presented a list of 23 open problems which has turned to be very influential for the mathematics of the 20th century. The fifth of these problems asked for a *topological description of Lie groups*: does every locally euclidean topological group admits a Lie group structure? Recall that a topological group is said to be *locally euclidean* if its identity element has a neighbourhood homeomorphic to an open subspace of  $\mathbb{R}^n$ . A positive solution to this problem was achieved in the early fifties by Gleason (1952), Montgomery and Zippin (1952), and Yamabe (1953). Hilbert's fifth problem motivated an enormous volume of work on locally compact groups that shed light on the structure of connected locally compact groups. An exposition on the celebrated solution of Hilbert's fifth problem can be found in Tao (2014).

*Remark* By Theorem 1.16, Lie group techniques may be used to analyse the structure of connected groups and their automorphisms. A canonical form for automorphisms of TDLC-groups has been developed in Willis (1994, 2001a); see § 1.5.1.

### 1.3 Totally disconnected locally compact groups

#### 1.3.1 van Dantzig's theorem

By using arguments from Hewitt and Ross (2012), we prove van Dantzig's theorem - together with some consequences - which can be considered as the key theorem in the theory of TDLC-groups: the topology of a TDLC-group admits a well-behaved basis of identity neighbourhoods.

**Theorem 1.18** (van Dantzig, 1936) *Let  $G$  be a TDLC-group. Then every neighbourhood of the identity contains a compact open subgroup.*

*Proof* We follow (Dikranjan, 2018, Proof of Theorem 7.4.2.(b)). By Vedenissov's Theorem (see (Außenhofer et al., 2021, Theorem B.6.10)), there exists a neighbourhood basis  $\mathcal{B}$  at the identity consisting of compact open sets. Let  $K \in \mathcal{B}$ . For each  $x \in K$ , there are an open set  $U_x \ni 1_G$  with  $xU_x \subseteq K$  (because left translation by  $x$  is continuous at  $1_G$ ) and an open set  $V_x \ni 1_G$  with  $V_x V_x \subseteq U_x$  (because the multiplication  $\mu: G \times G \rightarrow G$  is continuous at  $(1_G, 1_G)$ ). Since also the inversion map is continuous, the open set  $V_x$  can be chosen to be symmetric. For  $K$  is compact, there are  $x_1, \dots, x_n$  such that  $K \subseteq x_1 V_{x_1} \cup \dots \cup x_n V_{x_n}$ . Set  $V = V_{x_1} \cap \dots \cap V_{x_n}$  and notice that

$$KV \subseteq \left( \bigcup_{i=1}^n x_i V_{x_i} \right) V \subseteq \bigcup_{i=1}^n x_i V_{x_i} V_{x_i} \subseteq \bigcup_{i=1}^n x_i U_{x_i} \subseteq K.$$

The inclusion  $V = 1_G V \subseteq KV \subseteq K$  implies that  $VV \subseteq U$ ,  $VVV \subseteq U$ , etc. Since  $V$  is symmetric, the subgroup  $H$  generated by  $V$  is given by

$$H = \bigcup_{n \in \mathbb{N}} \underbrace{V \dots V}_n \subseteq K.$$

By Exercise 1.2.1,  $H$  is open (and so closed) and, for  $H \subseteq K$ ,  $H$  is also compact.  $\square$

Hence, every TDLC-group contains arbitrarily small (compact) open subgroups. This is the opposite of what occurs in the connected case, where the only open subgroup is the whole group (see Exercise 1.2.1). This is also the opposite of what occurs in Lie Groups, which admit a neighbourhood of the identity that contains only the trivial subgroup.

#### 1.3.2 Consequences of van Dantzig's theorem

Here we collect a few long-known consequences of van Dantzig's theorem.

**Proposition 1.19** *Given a locally compact group  $G$ , the identity component  $G_0$  coincides with the intersection of all open subgroups of  $G$ .*

*Proof* To prove that  $G_0$  is contained in the intersection it suffices to notice that every open subgroup  $H$  of  $G$  is a clopen set (see Exercise 1.2.1(3)).

For the reverse inclusion, we show that, for every  $x \in G \setminus G_0$ , there is an open subgroup  $H_x$  not containing  $x$ . By Proposition 1.7, the group  $G/G_0$  is TDLC. Therefore, van Dantzig's theorem yields a neighbourhood basis at  $G_0$  given by compact open subgroups of  $G/G_0$ . It follows that there is a compact open subgroup  $K \subseteq G/G_0$  not containing  $xG_0$ . Given the quotient map  $\pi_0: G \rightarrow G/G_0$ , we set  $H_x = \pi_0^{-1}(K)$ .  $\square$

**Corollary 1.20** *If a topological group  $G$  admits a neighbourhood basis  $\mathcal{B}$  at  $1_G$  consisting of compact open subgroups, then  $G$  is TDLC.*

*Proof* The only part that needs some work is the total disconnectedness of  $G$ . By Proposition 1.19,  $G_0 = \bigcap \{U \mid U \in \mathcal{B}\}$ . But  $\bigcap \{U \mid U \in \mathcal{B}\} = 1$  since we assume topological groups to be Hausdorff.  $\square$

In other words, van Dantzig's theorem characterizes TDLC-groups among topological (Hausdorff) groups: they are the ones whose compact open subgroups form a neighbourhood basis at  $1_G$ . The abundance of compact open subgroups will reveal itself to be the most fruitful property of TDLC-groups.

In general, total disconnectedness is not preserved under taking quotients. Thanks to van Dantzig's theorem this is not the case for locally compact groups.

**Proposition 1.21** *The quotient of a TDLC-group by a closed normal subgroup is totally disconnected.*

*Proof* Let  $G$  be a TDLC-group and let  $N$  be a closed normal subgroup of  $G$ . It follows from van Dantzig's theorem that the collection of all compact open subgroups of  $G$  form a neighbourhood basis at  $1_G$ . Since quotient maps are open, the quotient  $G/N$  admits a neighbourhood basis at  $N$  formed by compact open subgroups, that are the quotients of all compact open subgroups of  $G$ . Thus  $G/N$  is TDLC by Corollary 1.20.  $\square$

**Exercise 1.3.1** Every locally compact group  $G$  contains an open subgroup  $H$  which is compact-by-connected.

**Proposition 1.22** *A compact totally disconnected group is a projective limit of finite groups. In particular, a topological group is profinite if, and only if, it is compact and totally disconnected.*

*Proof* Let  $G$  be a compact totally disconnected group. The set  $\mathcal{O}$  of all compact open subgroups of  $G$  form a neighbourhood basis at  $1_G$ . Since  $G$  is compact, every subgroup  $H \in \mathcal{O}$  contains a subgroup which is both open and normal in  $G$  (see Exercise 3(2)). Thus, the family  $\mathcal{NO} = \{H \in \mathcal{O} \mid H \trianglelefteq G\}$  is a neighbourhood basis at  $1_G$ . Therefore, the morphism  $G \rightarrow \prod_{H \in \mathcal{NO}} G/H$  of compact groups is injective and continuous, and provides a topological isomorphism from  $G$  to a closed subgroup of the latter product of finite groups.  $\square$

### 1.3.3 Examples of TDLC-groups

Many examples of a TDLC-group can be produced by means of van Dantzig's theorem.

- Discrete groups and profinite groups are rather trivial examples.
- (Local fields) Let  $\mathbb{K}$  be a local field; see Example 1.13. Then  $\mathbb{K}$  has a unique maximal compact subring

$$\mathfrak{o}_{\mathbb{K}} = \{x \in \mathbb{K} \mid \{x^n \mid n \geq 1\} \text{ is relatively compact}\}$$

which has a unique maximal ideal

$$\mathfrak{p}_{\mathbb{K}} = \{x \in \mathbb{K} \mid \lim_{n \rightarrow \infty} x^n = 0\}.$$

Both  $\mathfrak{o}_{\mathbb{K}}$  and  $\mathfrak{p}_{\mathbb{K}}$  are compact and open in  $\mathbb{K}$ . The ideal  $\mathfrak{p}_{\mathbb{K}}$  is principal in  $\mathfrak{o}_{\mathbb{K}}$ : there is  $\pi \in \mathbb{K}$  such that  $\mathfrak{p}_{\mathbb{K}} = (\pi)$ . The nested sequence of ideals

$$(\pi) \supset \cdots \supset (\pi^n) \supset (\pi^{n+1}) \supset \cdots$$

constitutes a basis of compact open subgroups at 0 in  $\mathbb{K}$ .

Local fields fall into two families (see (Palmer, 1994, § 12.3.4)):

1. the fields of  $p$ -adic numbers,  $\mathbb{Q}_p$  and their finite extensions, and
2. the fields of formal Laurent series,  $\mathbb{F}_q((t))$ , over some finite field  $\mathbb{F}_q$ .

Note that  $\mathbb{Q}_p$  admits  $\{p^n \mathbb{Z}_p\}_{n \in \mathbb{N}}$  as basis of compact open subgroups, where  $\mathbb{Z}_p$  is the ring of  $p$ -adic integers, and  $\mathbb{F}_q((t))$  has  $\{t^n \mathbb{F}_q((t))\}_{n \in \mathbb{N}}$  as basis of compact open subgroups, where  $\mathbb{F}_q((t))$  is the ring of formal Taylor series over  $\mathbb{F}_q$ .

- (Linear groups over local fields) Let  $\mathbb{K}$  be a local field. For instance, the group  $GL_n(\mathbb{K})$  is TDLC with the topology inherited by  $\mathbb{K}^{n^2}$ .

- (Lie groups over local fields) An exact analogue of Lie theory exists for analytic groups defined over local fields such as  $\mathbb{Q}_p$  and  $\mathbb{F}_q((t))$ ; see (Caprace and Monod, 2018, § 4) and Bode and Dupr (2021).
- (Automorphism groups of connected locally-finite graphs) A **graph**  $\Gamma$  is a pair  $(V\Gamma, E\Gamma)$  where  $V\Gamma$  is a set and  $E\Gamma$  is a collection of unordered distinct pairs of elements from  $V\Gamma$ . The elements of  $V\Gamma$  are called **vertices** and the elements of  $E\Gamma$  are called **edges**. We will need a bit of terminology for graphs: two vertices  $v$  and  $u$  are said to be adjacent, if  $\{v, u\}$  is an edge in  $\Gamma$ ; a graph is **locally finite** if each vertex  $v$  has a finite number of adjacent vertices; a **path of length  $n$**  from  $v$  to  $u$  is a sequence  $(v = v_0, v_1, \dots, v_n = u)$  of vertices, such that  $v_i$  and  $v_{i+1}$  are adjacent for  $i = 0, 1, \dots, n-1$ ; a graph is **connected** if for any two vertices  $v$  and  $u$  there is a path from  $v$  to  $u$  in the graph. An **automorphism** of a graph  $\Gamma$  is a bijection  $\varphi: V\Gamma \rightarrow V\Gamma$  such that  $\{\varphi(v), \varphi(w)\} \in E\Gamma$  if and only if  $\{v, w\} \in E\Gamma$ . The collection of automorphisms forms a group under composition, and it is denoted by  $\text{Aut}(\Gamma)$ . Let  $\Gamma$  be a connected graph and endow  $\text{Aut}(\Gamma)$  with the compact-open topology via considering  $V\Gamma$  to be a discrete space. Namely, a basis of this topology is given by the sets

$$\Sigma_{v,w} = \{g \in \text{Aut}(\Gamma) \mid g(v_i) = w_i\},$$

where  $v = (v_1, \dots, v_n)$  and  $w = (w_1, \dots, w_n)$  range over all finite<sup>8</sup> tuples of vertices of  $\Gamma$ . In particular, two automorphisms of  $\Gamma$  are “close” to each other if they agree on “many” vertices.

**Exercise 1.3.2** The compact-open topology is a group topology on  $\text{Aut}(\Gamma)$  and it coincides with the pointwise convergence topology.

*Remark* The compact-open topology on  $\text{Aut}(\Gamma)$  also coincides with the **permutation topology** (cf. Karrass and Solitar (1956)).

Let  $G = \text{Aut}(\Gamma)$ . The identity element of  $G$  belongs to an open set  $\Sigma_{a,b}$  iff  $a = b$ . Consequently, the compact-open topology on  $G$  has a neighbourhood basis at the identity formed by the pointwise stabilisers of finite sets of vertices. Recall that, given a vertex  $v \in \Gamma$ , the set  $G_{(v)} = \{g \in G \mid g(v) = v\}$  is a subgroup of  $G$  which is called the **stabiliser of  $v$** . Given a finite set of vertices  $\mathcal{V}$ , the intersection  $G_{(\mathcal{V})} = \bigcap_{v \in \mathcal{V}} G_{(v)}$  is the **pointwise stabiliser** of the set  $\mathcal{V}$ . Clearly, pointwise stabilisers of finite set of vertices are open subgroups of  $G$ .

<sup>8</sup> Notice that the length of the tuples is arbitrary.

**Proposition 1.23** *Let  $\Gamma$  be a connected locally finite graph. The pointwise stabilisers of finite sets of vertices are compact in the compact-open topology. In particular,  $G = \text{Aut}(\Gamma)$  is a TDLG-group.*

*Sketch of the proof.* It suffices to prove the claim for the stabiliser  $G_{(v)}$  of an arbitrary vertex  $v$ . For  $k > 0$ , set

$$S_k(v) = \{w \in V\Gamma \mid d_\Gamma(v, w) \leq k\},$$

where  $d_\Gamma(v, w)$  denotes the length of the shortest path from  $v$  to  $w$ . Since  $\Gamma$  is locally finite, every  $S_k(v)$  is finite. Let  $\text{Sym}(S_k(v))$  be endowed with the discrete topology. Clearly, the stabiliser  $G_{(v)}$  permutes the elements in each  $S_k(v)$  i.e., there exists a group homomorphism  $\phi_k: G_{(v)} \rightarrow \text{Sym}(S_k(v))$ . The family  $\{\phi_k\}_{k>0}$  can be then used to construct an injective group homomorphism from  $G_{(v)}$  to the profinite group  $\prod_{k>0} \text{Sym}(S_k(v))$  which is continuous and closed.  $\square$

*Remark* The latter result does not state the “non-discreteness” of  $\text{Aut}(\Gamma)$ . Indeed, it is a difficult task to determine whether a given group is discrete.

- (Neretin group of spheromorphisms of a  $d$ -regular tree) A **tree** is a connected graph without nontrivial cycles, where for nontrivial cycle we mean a path  $(v_0, \dots, v_n)$  such that  $n \geq 1$  and  $v_0 = v_n$ . A vertex  $v \in V\Gamma$  of degree 1 (i.e., has a unique adjacent vertex) is called a **leaf**. For  $d \in \mathbb{N}$ , an **infinite  $d$ -regular tree** is an infinite tree whose vertices have degree  $d + 1$ . A **finite  $d$ -regular tree** is a finite tree, whose every vertex is either a leaf, or has degree  $d + 1$ , i.e., it is an **internal vertex**. A **rooted tree** is a tree with a distinguished vertex  $o \in T$ , called its **root**. If the tree is rooted then  $d$ -regularity requires the root to have degree  $d$  instead of  $d + 1$ .

An important property of a tree  $T$  is given by the fact that there exists a unique path connecting two vertices  $v$  and  $u$ . A **ray** in  $T$  is defined to be an infinite path, i.e., a sequence  $(v_0, v_1, \dots)$  of distinct vertices of  $T$  such that the consecutive ones are adjacent. Two rays are said to be **asymptotic** if they have common tails<sup>9</sup>. Equivalence classes of asymptotic rays are called the **ends** of  $T$ . All ends of  $T$  form the set  $\partial T$  which is called the **boundary** of the tree.

<sup>9</sup> A tail of a sequence is a subsequence obtained after removing finitely many initial elements.

**The construction of  $\mathcal{N}_d$ :** Here we follow the construction and the arguments of (Caprace and Monod, 2018, Chapter 8). Let  $T$  be a  $d$ -regular tree. For every finite  $d$ -regular subtree  $F \subseteq T$ , denote by  $T \setminus F$  the (no longer connected) graph obtained by removing from  $T$  all the edges and internal vertices of  $F$ . The connected components of  $T \setminus F$  are rooted  $d$ -regular trees whose roots are the leaves of  $F$ . In particular,  $T \setminus F$  is a rooted  $d$ -regular forest such that  $\partial(T \setminus F) = \partial T$ <sup>10</sup>.

Let  $F_1, F_2 \subseteq T$  be two finite  $d$ -regular subtrees with the same number of leaves. Each forest isomorphism  $\phi: T \setminus F_1 \rightarrow T \setminus F_2$  induces a homeomorphism  $\phi_*$  of  $\partial T$ , called *spheromorphism* of  $T$ . Clearly, different choices of subtrees  $F_1, F_2$  can induce the same spheromorphism. Therefore,  $\phi$  is just a representative of  $\phi_*$ . This is important because, for each pair of spheromorphisms  $\phi_*$  and  $\psi_*$ , we find *composable* representatives: we enlarge the finite trees in the representation of the spheromorphisms in order to make the isomorphisms  $\phi$  and  $\psi$  share a target forest and a source forest. This procedure shows that the spheromorphism  $\psi_* \circ \phi_*$  is well-defined (in particular, it coincides with the composition in  $\text{Homeo}(\partial T)$ ). Hence, the set of all spheromorphisms of a  $d$ -regular tree is a group, which is called **Neretin group** and denoted by  $\mathcal{N}_d$ . The group of spheromorphisms of a  $d$ -regular tree has been introduced by Neretin (1992) by analogy with the diffeomorphism group of a circle. Roughly speaking, a spheromorphism of  $\partial T$  is a transformation induced in the boundary  $\partial T$  by a piecewise tree automorphism; indeed, spheromorphisms are also known as *almost automorphisms of  $T$* .

**Fact 1.24** *The set of all spheromorphisms is a subgroup of the homeomorphism group of the boundary  $\partial T$ . Moreover, every automorphism  $\phi$  of  $T$  induces an isomorphism  $T \setminus F \rightarrow T \setminus \phi(F)$  of forests which is independent on the finite  $d$ -regular subtree  $F$ . Thus,  $\text{Aut}(T)$  can be regarded as a subgroup of  $\mathcal{N}_d$ .*

*Remark* Let  $\phi_*$  be a spheromorphism of  $T_d$  and suppose that  $\phi_*$  admits a representative  $\phi: T_d \setminus F \rightarrow T_d \setminus F$  that leaves the trees of  $T_d \setminus F$  in place. Then  $\phi$  can be extended to an automorphism of the tree  $T_d$  which belongs to the pointwise stabiliser of the finite tree  $F$ . As a consequence,  $\phi_*$  belongs to the image of  $\text{Aut}(T_d)$  in  $\mathcal{N}_d$ .

In order to topologize the group  $\mathcal{N}_d$ , the first attempt is to endow

<sup>10</sup> Since a forest is a disjoint union of trees the notion of boundary can be easily extended to forests.

$\text{Homeo}(\partial T)$  with the compact-open topology and then give  $\mathcal{N}_d$  the subspace topology. Unfortunately, the resulting topological group is not locally compact:  $\mathcal{N}_d$  is not closed in  $\text{Homeo}(\partial T)$  with respect to the compact-open topology (see Proposition 1.15).

Instead of restricting a topology from a larger topological group, we could try to “copy and paste around” a topology coming from an abstract subgroup which is also a topological group.

**Lemma 1.25** ((Caprace and Monod, 2018, Lemma 8.4, pg. 137))  
*Suppose that an abstract group  $G$  contains a topological group  $H$  as a subgroup. If, for all open subsets  $U \subseteq H$  and  $g, g' \in G$ , the intersection  $gUg' \cap H$  is open in  $H$ , then  $G$  admits a unique group topology such that the inclusion  $H \rightarrow G$  is continuous and open.*

**Theorem 1.26** *Neretin group  $\mathcal{N}_d$  admits a unique group topology such that the natural embedding  $\text{Aut}(T_d) \rightarrow \mathcal{N}_d$  is continuous and open. With this topology,  $\mathcal{N}_d$  is a TDLC-group.*

*Proof* By the lemma above, one needs to show that for every open  $U \subseteq \text{Aut}(T_d)$  and all  $\phi_*, \psi_* \in \mathcal{N}_d$ , the subset  $\text{Aut}(T_d) \cap \phi_* U \psi_*$  is open in  $\text{Aut}(T_d)$ . A sub-basis of the compact-open topology on  $\text{Aut}(T_d)$  is given by vertex stabilisers and therefore one has to show the claim only for the sets in the sub-basis.

To this end, let  $v$  be a vertex of  $\Gamma$ . Let  $S$  be a sufficiently large sphere centred at  $v$  such that the spheromorphisms  $\phi_*$  and  $\psi_*$  admit representatives  $\phi: T_d \setminus F_1 \rightarrow T_d \setminus S$  and  $\psi: T_d \setminus S \rightarrow T_d \setminus F_2$ . Denote by  $G_{(S)}$  the pointwise stabiliser of  $S$ . Since  $G_{(S)}$  is an open subgroup of  $\text{Aut}(T_d)$  contained in  $G_{(v)}$ , there exist finitely many elements  $g_1, \dots, g_n \in G_{(v)}$  such that  $G_{(v)} = \bigsqcup_{i=1}^n g_i G_{(S)}$ . Therefore,

$$\psi_* G_{(v)} \phi_* = \bigsqcup_{i=1}^n \psi_* g_i G_{(S)} \phi_* = \bigsqcup_{i=1}^n \psi_* g_i \phi_* (\phi_*^{-1} G_{(S)} \phi_*).$$

By Remark 1.3.3,  $\phi_*^{-1} G_{(S)} \phi_*$  coincides with the pointwise stabiliser of the finite tree  $F_1$  and, therefore, it is contained in the image of  $\text{Aut}(T_d)$  in  $\mathcal{N}_d$ . It then follows that  $\psi_* G_{(v)} \phi_* \cap \text{Aut}(T_d)$  is open in  $\text{Aut}(T_d)$  (it is union of translates of the open subgroup  $G_{(F_1)}$ ).  $\square$

- (Fundamental groups of graphs of profinite groups) Bass-Serre theory carries over to the realm of TDLC-groups. Let  $(\mathcal{G}, \Lambda)$  be a finite graph of profinite groups; see Castellano and Weigel (2016). The fundamental group  $\pi_1(\mathcal{G}, \Lambda)$  can be endowed easily with a TDLC-group topology.

- (Topological semi-direct products) Let  $G$  and  $H$  be topological groups. Suppose that  $G$  acts on  $H$  continuously, i.e., there is a group action of  $G$  on  $H$  such that the map  $\alpha: G \times H \rightarrow H$  defined by the action is continuous. The **topological semi-direct product** is the abstract semi-direct product  $H \rtimes G$  endowed with the product topology.

**Exercise 1.3.3** Let  $G$  and  $H$  be TDLC-groups such that  $G$  acts continuously on  $H$ . The topological semi-direct product  $H \rtimes G$  is a TDLC-group.

- (Powers of topological groups) Let  $G$  be a locally compact group. For  $I$  infinite, the power  $G^I$  fails to be locally compact as soon as  $G$  is non-compact. To deal with this issue, given a compact open subgroup  $U$  in  $G$ , one defines the **semi-restricted power**

$$G^{I,U} = \{(g_i)_{i \in I} \in G^I \mid g_i \in U \text{ for all but finitely many } i \in I\}.$$

There is a unique group topology on  $G^{I,U}$  that makes the embedding of  $U^I$  a topological isomorphism onto an open subgroup. Moreover, such a group topology is locally compact; see (Cornulier, 2019, Proposition 2.4). TDLC-groups are full of compact open subgroups, and therefore they are amenable to this construction; in particular, semi-restricted powers of TDLC-groups are again TDLC-groups.

## 1.4 Finiteness properties for TDLC-groups

### 1.4.1 Compact generation and presentation

There are several finiteness conditions that a TDLC-group can satisfy. At this early stage, we are interested in two finiteness conditions that naturally generalise the notions of finite generation and finite presentation in the context of locally compact groups.

**Definition 1.27** A locally compact group  $G$  is said to be

- (CG) **compactly generated** if it has a compact generating set  $S$ .
- (CP) **compactly presented** if it has a presentation  $\langle S \mid R \rangle$  as an abstract group with the generating set  $S$  compact in  $G$  and the relators in  $R$  of bounded length.

It is straightforward that (CP) implies (CG). The converse is not true; see for instance (Cornulier and de la Harpe, 2016, Example 8.A.28). The notion of compact presentation was introduced in 1964 by Kneser but

it has attracted much less attention than compact generation until recently: for example, Cornulier and de la Harpe (2016) proved that compact presentation is equivalent to a metric condition and Castellano and Weigel (2016) provided an equivalent notion of compact presentation via generalised presentations which is based on van Dantzig's theorem and the notion of fundamental group of finite graphs of profinite groups.<sup>11</sup>

- Example 1.28**
1. Every profinite group is (trivially) compactly generated and compactly presented.
  2. Every compactly generated abelian TDLC-group is topologically isomorphic to  $\mathbb{Z}^n \times K$ , where  $n \in \mathbb{N}$  and  $K$  is a compact abelian group; see (Dikranjan, 2018, Theorem 12.5.5). In particular, it is compactly presented.
  3. The field of  $p$ -adic numbers  $\mathbb{Q}_p$  is not compactly generated because it is the ascending union of nested compact open subgroups, i.e.,  $\mathbb{Q}_p = \bigcup_{n \in \mathbb{Z}} p^n \mathbb{Z}_p$ .
  4. The automorphism group of a  $d$ -regular tree is compactly generated; see Corollary 1.32.
  5. The special linear group  $\mathrm{SL}_2(\mathbb{Q}_p)$  is compactly generated. Indeed, by Ihara's Theorem (Serre, 1980, p. 143, Corollary 1 to Theorem 3), we can decompose  $\mathrm{SL}_2(\mathbb{Q}_p)$  into the amalgamated free product

$$\mathrm{SL}_2(\mathbb{Z}_p) *_I \mathrm{SL}_2(\mathbb{Z}_p),$$

where  $I$  is a compact open subgroup.

6. More generally, the fundamental group of a finite graph of profinite groups satisfies (CP); see (Castellano and Weigel, 2016, § 5.8).
7. The Neretin groups are compactly presented; see Caprace and De Medts (2011) and Le Boudec (2017).

**Fact 1.29** *Every locally compact group is a directed union of compactly generated open subgroups.*

*Proof* It suffices to notice that, for any  $g \in G$  and any compact open neighbourhood  $V_g$  of  $g$ , the subgroup  $\bigcup_{n > 0} (V_g \cup V_g^{-1})^n$  is open in  $G$  and compactly generated.  $\square$

### 1.4.2 The Cayley-Abels graphs

Recall that a **group  $G$  acts on a graph  $\Gamma$**  if the set of vertices  $V\Gamma$  is a  $G$ -set and, for every  $g \in G$ ,  $\{gv, gw\} \in E\Gamma$  if and only if  $\{v, w\} \in E\Gamma$ .

<sup>11</sup> which plays a role in the geometric group theory of TDLC-groups that can be compared with the role played by free groups in the discrete case.

The group  $G$  acts **vertex-transitively** on  $\Gamma$  if  $V\Gamma$  is a transitive  $G$ -set. Given a vertex  $v \in V\Gamma$ , the set  $G_{(v)} = \{g \in G \mid gv = v\}$  is the **vertex stabiliser** of  $v$  in  $G$ .

**Definition 1.30** For a TDLC-group  $G$ , a locally finite connected graph  $\Gamma$  on which  $G$  acts vertex-transitively with compact open vertex stabilisers is called a **Cayley-Abels graph of  $G$** .

**Exercise 1.4.1** Let  $\Gamma$  be a Cayley-Abels graph of  $G$ . Let  $V\Gamma$  be endowed with the discrete topology. Prove that the map  $G \times V\Gamma \rightarrow V\Gamma$  is continuous; that is, a TDLC-group  $G$  always acts continuously on its Cayley-Abels graphs. Moreover, prove that, as far as  $G$  is non-discrete, the action of  $G$  on  $\Gamma$  is never free<sup>12</sup>.

**Proposition 1.31** *Let  $G$  be a TDLC-group. If  $G$  has a Cayley-Abels graph, then  $G$  is compactly generated.*

*Proof* Let  $\Gamma$  be a Cayley-Abels graph of  $G$  and  $v \in V\Gamma$ . Since  $\Gamma$  is locally finite, one has  $\text{star}(v) = \{v_1, \dots, v_n\}$ . Since  $G$  acts on  $\Gamma$  vertex-transitively, for every  $i = 1, \dots, n$ , there is a  $g_i \in G$  such that  $v_i = g_i v$ . We claim that, for every  $g \in G$ , there is an  $h \in \langle g_1, \dots, g_n \rangle$  such that  $gv = hv$ . This implies that  $h^{-1}g \in G_{(v)}$ , which is compact and open by hypothesis. In other words,  $G = \langle G_{(v)}, g_1, \dots, g_n \rangle$ .

Let us prove the claim: for every  $g \in G$  there is a path in  $\Gamma$  connecting  $v$  and  $gv$  because  $\Gamma$  is connected. We proceed by induction on the length of the path. For  $k = 0$  there is nothing to prove. Suppose the hypothesis for  $k$  and prove it for  $k+1$ . For  $\Gamma$  is vertex-transitive, a path of length  $k+1$  connecting  $v$  and  $gv$  is given by a  $(k+1)$ -tuple  $(v, \gamma_1 v, \dots, \gamma_k v, gv)$  with  $\gamma_1, \dots, \gamma_k \in G$ . By the inductive hypothesis, there is  $h \in \langle g_1, \dots, g_n \rangle$  such that  $\gamma_k v = hv$ . Therefore, the group element  $h^{-1}$  maps the edge  $\{\gamma_k v, gv\}$  to the edge  $\{v, h^{-1}gv\}$ . In other words,  $h^{-1}gv$  is adjacent to  $v$ , i.e.,  $h^{-1}gv = v_j = g_j v$  for some  $j \in \{1, \dots, n\}$  and the claim holds.  $\square$

**Corollary 1.32**  *$\text{Aut}(\mathcal{T}_d)$  is compactly generated for every  $d \in \mathbb{N}$ .*

Now, we do the converse: we start with a compactly generated TDLC-group  $G$  and we construct a (family of) Cayley-Abels graph(s) of  $G$ . In particular, we show that, for every compact open subgroup  $U$  of  $G$ , there is a Cayley-Abels graph admitting  $U$  as stabiliser of some vertex.

Let  $U$  be a compact open subgroup of  $G$ . For every symmetric subset

<sup>12</sup> A groups acts *freely* on a set if point stabilisers are trivial.

$S = S^{-1} \subseteq G \setminus U$  define the graph  $\Gamma_{U,S}^G$  by setting

$$V\Gamma_{U,S}^G = \{gU \mid g \in G\}, \quad \text{and} \quad E\Gamma_{U,S}^G = \{\{gU, gsU\} \mid g \in G, s \in S\}.$$

Clearly,  $G$  acts transitively on the set of vertices of  $\Gamma_{U,S}^G$ .

**Proposition 1.33** *With the above notation, the following hold:*

1. *if  $S$  is a finite set, then  $\Gamma_{U,S}^G$  is locally finite;*
2.  *$\Gamma_{U,S}^G$  is connected if, and only if,  $G = \langle S \cup U \rangle$ .*

*If  $G$  is compactly generated, there exists a Cayley-Abels graph of  $G$ .*

*Proof* 1. Since the action is vertex-transitive, it suffices to prove that the vertex  $U$  has finitely many neighbours if  $S$  is finite. Since  $U = xU$ , for every  $x \in U$ , one has that the set  $\{xU, xsU\}$  is an edge for every  $s \in S$ , and therefore the set of all neighbours of  $U$  coincides with  $\{xsU \mid x \in U, s \in S\}$ . In order to determine the cardinality of such a set, one must count the number of left cosets of  $U$  that are necessary to cover each double coset  $UsU$ . But this number is finite because  $U$  is open and the double coset  $UsU$  is compact (since  $U$  is compact). Therefore, if  $S$  is finite, the graph  $\Gamma_{U,S}^G$  is locally finite.

2. Suppose the graph  $\Gamma_{U,S}^G$  is connected. For every  $g \in G$ , there is a path  $p = (v_0, \dots, v_n)$  connecting the vertex  $U$  to the vertex  $gU$ . In particular, the vertices of the path  $p$  can be written as

$$v_0 = U, \quad v_1 = u_1s_1U, \quad \dots, \quad v_n = u_1s_1 \cdots u_ns_nU,$$

where each  $u_i \in U$  and each  $s_j \in S$ . Since  $u_1s_1 \cdots u_ns_nU = gU$ , it follows that  $g$  belongs to the subgroup generated by  $U \cup S$ .

Conversely, suppose that  $G = \langle U \cup S \rangle$ . Let  $gU$  and  $hU$  be any two vertices of the graph. The group element  $g^{-1}h$  can be then written as a word  $u_1s_1 \cdots u_ns_nu_{n+1}$  such that each  $u_i \in U$  and each  $s_j \in S$ . Thus, the sequence of vertices

$$(gU, gu_1s_1U, gu_1s_1u_2s_2U, \dots, gu_1s_1 \cdots u_ns_nu_{n+1}U = hU)$$

is a path in  $\Gamma_{U,S}^G$  connecting  $gU$  and  $hU$ . □

*Remark* The first (technical) construction of the Cayley-Abels graph is due to Abels (1973/74). A less technical approach to Cayley-Abels graphs was provided in Krön and Möller (2008), where the Cayley-Abels graphs were at the time called *rough Cayley graphs*. Today the widely accepted nomenclature is “Cayley-Abels graph”.

**Proposition 1.34** *Let  $G$  be a compactly generated TDLC-group with Cayley-Abels graph  $\Gamma$ . The group homomorphism  $\psi: G \rightarrow \text{Aut}(\Gamma)$  defined by the action of  $G$  on  $\Gamma$  is continuous, the kernel of  $\psi$  is compact and the image of  $\psi$  is closed.*

*Proof* A basis of the compact-open topology of  $\text{Aut}(\Gamma)$  is given by the family of pointwise stabilisers in  $\text{Aut}(\Gamma)$  of finite sets of vertices. The pre-image of each of these sets is the intersection of finitely many open subgroups of  $G$  (that are the stabilisers  $G_{(v)}$  of the vertices  $v$  in the finite set). Since each stabiliser  $G_{(v)}$  is open,  $\psi$  is continuous.

The kernel of  $\psi$  is closed because it is the pre-image of the closed set  $\{1\}$  and then it is compact since  $\ker(\psi) \subseteq G_{(v)}$ , for any  $v \in V\Gamma$ .

To prove that the image  $\psi(G)$  is closed, it suffices to see that  $\psi(G) \cap H$  is closed for every  $H$  in the basis of the compact-open topology of  $\text{Aut}(\Gamma)$ . Since every such  $H$  is the intersection of finitely many vertex stabilisers in  $\text{Aut}(\Gamma)$ , we only need to prove that  $\psi(G) \cap H$  is closed whenever  $H$  is the stabiliser in  $\text{Aut}(\Gamma)$  of a single vertex  $v$ . In such a case,  $\psi(G) \cap H$  coincides with  $\psi(G_{(v)})$  which is compact because  $\psi$  is continuous and  $G_{(v)}$  is compact. In particular,  $\psi(G) \cap H$  is closed.  $\square$

**Corollary 1.35** *Compactly generated TDLC-groups are second countable modulo a compact normal subgroup.*

The representation  $\psi: G \rightarrow \text{Aut}(\Gamma)$  is called the **Cayley-Abels representation of  $G$** . Hence, there is a correspondence - up to compact kernels - between compactly generated TDLC-groups and closed subgroups of automorphism groups of connected locally finite graphs.

*Remark* One can say more on the image  $\psi(G)$  of the Cayley-Abels representation:  $\psi(G)$  is a cocompact subgroup of  $\text{Aut}(\Gamma)$ , see (Wesolek, 2018, Lemma 3.12).

**Exercise 1.4.2** Suppose  $G$  is a compactly generated TDLC-group and  $\Gamma$  is a Cayley-Abels graph of  $G$ . Show that a closed subgroup  $K \leq G$  is compact if and only if, for all  $v \in V\Gamma$ ,  $Kv := \{kv \mid k \in K\}$  is finite.

### The geometric structure of compactly generated TDLC-groups:

**Definition 1.36** (Gromov (1984)) Two metric spaces  $(X, d_X)$  and  $(Y, d_Y)$  are said to be **quasi-isometric** if there is a map  $\phi: X \rightarrow Y$  and constants  $a \geq 1$  and  $b \geq 0$  such that, for all  $x_1, x_2 \in X$ , one has

$$\frac{1}{a}d_X(x_1, x_2) - \frac{b}{a} \leq d_Y(\phi(x_1), \phi(x_2)) \leq ad_X(x_1, x_2) + ab,$$

and, for all  $y \in Y$ ,

$$d_Y(y, \phi(X)) \leq b.$$

Such a map  $\phi$  is called a **quasi-isometry**. Moreover, being quasi-isometric is an equivalence relation on the class of metric spaces.

Every connected graph  $\Gamma$  can be regarded as a metric space: two vertices  $v$  and  $w$  are points at distance 1 if, and only if, there is an edge connecting  $v$  and  $w$ . In other words, we endow the set  $V\Gamma$  with the path-length metric  $d_\Gamma: V\Gamma \times V\Gamma \rightarrow \mathbb{N}$  defined as follows:

$$d_\Gamma(v, w) = \min\{\text{length of } p \mid p \text{ path connecting } v \text{ and } w\}, \quad v, w \in V\Gamma.$$

**Theorem 1.37** (Abels (1973/74), (Krön and Möller, 2008, Theorem 2))  
*Let  $G$  be a compactly generated TDLC-group. Any two Cayley-Abels graphs of  $G$  are quasi-isometric.*

The quasi-isometric invariance allows us to define *geometric invariants* of a compactly generated TDLC-group  $G$  by considering quasi-isometric invariants of a Cayley-Abels graph associated to  $G$ . For example, one can give the following definitions (that are long-known for finitely generated discrete groups) for a compactly generated TDLC-group  $G$ :

- (Hyp)  $G$  is said to be **hyperbolic** if some (and hence any) Cayley-Abels graph of  $G$  is hyperbolic.
- (Ends) The **number of ends** of  $G$  is defined to be the number of ends of some (and hence any) Cayley-Abels graph of  $G$ .

The class of hyperbolic TDLC-groups is a rich source of compactly presented TDLC-groups; see Cornulier and de la Harpe (2016). Indeed, geometric invariants often reflect structural properties of the group: for instance, an analogue of the famous Stallings' decomposition theorem is available in the context of TDLC-groups; see (Abels, 1973/74, Struktur-satz 5.7 and Korollar 5.8), (Krön and Möller, 2008, Theorem 1.3) and Castellano (2020).

### 1.4.3 Finiteness conditions in higher dimension

For discrete groups, finite generation is the first in two sequences of increasingly stronger properties: the homological finiteness conditions, the **types**  $(\mathbf{FP}_n)_{n \in \mathbb{N}}$  over a commutative unital ring  $R$ , and the homotopical finiteness conditions, the **types**  $(\mathbf{F}_n)_{n \in \mathbb{N}}$ . We recall the definitions here, but the reader is referred to (Brown, 1994, Chapter VIII) for details:

- (FP<sub>n</sub>) A discrete group  $G$  is **of type FP<sub>n</sub>** ( $0 \leq n < \infty$ ) over  $R$  if there is a projective resolution  $\{P_i\}$  of the trivial module  $R$  over  $R[G]$  such that each projective  $R[G]$ -module  $P_i$  is finitely generated for  $i \leq n$ . (If  $R = \mathbb{Z}$ , the reference to the ring  $R$  usually drops).
- (F<sub>n</sub>) A discrete group  $G$  is **of type F<sub>n</sub>** ( $0 \leq n < \infty$ ) if there exists a contractible  $G$ -CW-complex with finite cell stabilisers and such that  $G$  acts on the  $n$ -skeleta with finitely many orbits.

We will refer to the properties above as the classical finiteness properties for discrete groups. These properties are known to satisfy the following:

- A discrete group  $G$  is of type  $F_1$  if, and only if, it is finitely generated if, and only if, it is of type  $FP_1$  over  $R$ .
- A discrete group  $G$  is of type  $F_2$  if and only if it is finitely presented but being of type  $FP_2$  is strictly weaker than finite presentation; see Bestvina and Brady (1997).
- For each  $n \geq 1$ , a discrete group of type  $F_n$  is of type  $FP_n$  over  $R$  but the converse is not true (the converse becomes true if the group is suppose to be finitely presented and  $R = \mathbb{Z}$ ).
- Being of type  $FP_n$  over  $R$  (resp. of type  $F_n$ ) is a geometric property of the finitely generated group, that is, it invariant up to quasi-isometry.

A first attempt at generalising these properties to the realm of locally compact groups is due to Abels and Tiemeyer [3]. They introduced **compactness properties** for locally compact groups - we avoid here the (very technical) definitions - that are two sequences  $(\mathbf{CP}_n)_{n \geq 0}$  and  $(\mathbf{C}_n)_{n \geq 0}$  of increasingly stronger properties satisfying:

- for all  $n \geq 1$ , a discrete group is of type  $(\mathbf{CP}_n)$  (resp.  $\mathbf{C}_n$ ) if and only if it is of type  $FP_n$  (resp.  $F_n$ );
- a locally compact group is of type  $\mathbf{C}_1$  if, and only if, it is compactly generated if, and only if, it is of type  $\mathbf{CP}_1$ ;
- a locally compact group is of type  $\mathbf{C}_2$  if and only if it is compactly presented but being of type  $\mathbf{CP}_2$  is strictly weaker than compact presentation;
- for each  $n \geq 1$ , a locally compact group of type  $\mathbf{C}_n$  is also of type  $\mathbf{CP}_n$  but the converse is not true;
- being of type  $\mathbf{CP}_n$  (resp.  $\mathbf{C}_n$ ) is invariant “up to compactness”: the compactness properties remain unchanged by passing to a cocompact<sup>13</sup> subgroup or by taking the quotient by a compact normal sub-

<sup>13</sup> A closed subgroup  $H$  is *cocompact* if the quotient  $G/H$ , equivalently  $H \backslash G$ , equipped with the quotient topology is compact.

group. Such an invariance is weaker than the invariance up to quasi-isometry among the class of compactly generated locally compact groups.

For the (more amenable) class of TDLC-groups, a different approach to finiteness conditions was recently introduced and investigated in Castellano and Weigel (2016) and Castellano and Corob Cook (2020):

- (FP<sub>n</sub>) A TDLC-group  $G$  is **of type FP<sub>n</sub>** ( $0 \leq n < \infty$ ) over  $R$  if there is a resolution  $\{P_i\}$  of the trivial module  $R$  over  $R[G]$  such that each  $P_i$  is a permutation  $R[G]$ -module<sup>14</sup> with compact open stabilisers and finitely many orbits for  $i \leq n$ .
- (F<sub>n</sub>) A TDLC-group  $G$  is **of type F<sub>n</sub>** ( $0 \leq n < \infty$ ) if there exists a contractible  $G$ -CW-complex  $X$  with compact open cell stabilisers such that  $G$  acts on the  $n$ -skeleta of  $X$  with finitely many orbits.

These finiteness conditions for TDLC-groups satisfy the following:

- for  $R = \mathbb{Q}$ , permutation  $\mathbb{Q}[G]$ -modules with compact open stabilisers are projective objects in the category  $\mathbb{Q}[G]\mathbf{dis}$  of rational discrete  $\mathbb{Q}[G]$ -modules<sup>15</sup> (see Castellano and Weigel (2016)) and so we recover the homological flavour of the classical definition. In particular, whenever  $G$  is discrete, the new definition reduces to the classical notion of type FP<sub>n</sub> over  $\mathbb{Q}$ .
- for all  $n \geq 1$ , a discrete group is of type F<sub>n</sub> in the category of TDLC-groups if and only if it is of type F<sub>n</sub> in the classical sense because requiring compact open discrete stabilisers reduces to finite stabilisers;
- a TDLC-group is of type F<sub>1</sub> if, and only if, it is compactly generated if, and only if, it is of type FP<sub>1</sub> over  $R$  (see (Castellano and Corob Cook, 2020, Proposition 3.4) and (Castellano and Weigel, 2016, Proposition 5.3));
- a TDLC-group is of type F<sub>2</sub> if and only if it is compactly presented by (Castellano and Corob Cook, 2020, Proposition 3.4) but being of type FP<sub>2</sub> is strictly weaker than compact presentation;
- for each  $n \geq 1$ , a TDLC-group of type F<sub>n</sub> is also of type FP<sub>n</sub> over  $R$  (see (Castellano and Corob Cook, 2020, Fact 2.7)) but the converse is not true (the converse becomes true if the group is supposed to be compactly presented and  $R$  is replaced by  $\mathbb{Z}$ );

<sup>14</sup> A *permutation R[G]-module* is a module  $R[\Omega]$  freely  $R$ -generated by a  $G$ -set  $\Omega$ .

<sup>15</sup> An  $R[G]$ -module  $M$  is said to be *discrete* if the action  $G \times M \rightarrow M$  is continuous when  $M$  carries the discrete topology.

- being of type  $\text{FP}_n$  over  $R$  (resp.  $\text{F}_n$ ) is a geometric property (see (Castellano and Corob Cook, 2020, Corollary 5.7)).

*Remark* All the finiteness conditions above can be extended to  $n = \infty$ . Sauer and Thumann (2017) showed that Neretin groups are of type  $\text{F}_\infty$  which, in particular, implies the fact - observed already in Castellano and Weigel (2016) - that these groups are of type  $\text{FP}_\infty$  over  $\mathbb{Q}$ .

**Example 1.38** Hyperbolic TDLC-groups are of type  $\text{F}_n$  for some finite  $n$ . It is a classical result that, for a large enough constant  $d$ , the (topological realisation of the) Rips' complex  $P_d(\Gamma)$  is contractible whenever  $\Gamma$  is a hyperbolic (locally finite) graph.

Castellano and Corob Cook (2020) showed that it is possible to introduce further two sequences (**types**  $\text{KP}_n$ ) $_{n \geq 0}$  and (**types**  $\text{K}_n$ ) $_{n \geq 0}$  of increasingly stronger compactness properties. The concrete motivation comes from the fact that permutation  $R[G]$ -modules with compact open stabilisers fail to be projective over the ring  $R = \mathbb{Z}$  in the abelian category  ${}_{\mathbb{Z}[G]}\text{dis}$ . In such a case, being of type  $\text{FP}_n$  lack of a description based on the existence of partial projective resolutions of the trivial module  $\mathbb{Z}$  of finite type. The strategy to get back such a description is to embed the category  ${}_{\mathbb{Z}[G]}\text{dis}$  into a quasi-abelian category  ${}_{\mathbb{Z}[G]}\text{top}$  where the discrete permutation  $\mathbb{Z}[G]$ -modules with compact stabilisers go back to being projective again. The objects of  ${}_{\mathbb{Z}[G]}\text{top}$  are the so-called  $k$ - $\mathbb{Z}[G]$ -modules, i.e., module objects in the category of  $k$ -spaces over the  $k$ - $\mathbb{Z}$ -algebra  $\mathbb{Z}[G]$ . The reader is referred to Corob Cook (2017) for the definition and the background on this category. Since all locally compact Hausdorff spaces are  $k$ -spaces, every t.d.l.c. group is automatically a  $k$ -group and one can take advantage of the homological machinery developed in Corob Cook (2017) to define the following finiteness conditions:

- ( $\text{KP}_n$ ) A TDLC-group  $G$  is **of type**  $\text{KP}_n$  ( $0 \leq n < \infty$ ) if there is a projective resolution  $\{P_i\}$  of the trivial module  $\mathbb{Z}$  in  ${}_{\mathbb{Z}[G]}\text{top}$  such that each  $P_i$  is a free  $k$ - $\mathbb{Z}[G]$ -module on a compact space for  $i \leq n$ .
- ( $\text{K}_n$ ) A TDLC-group  $G$  is **of type**  $\text{K}_n$  ( $0 \leq n < \infty$ ) if there exists a contractible  $G$ -KW-complex<sup>16</sup>  $X$  with cocompact  $n$ -skeleta, in the compact Hausdorff model structure on  $G$ - $k$ -spaces.

*Remark* In Castellano and Corob Cook (2020) the sequence (types  $\text{KP}_n$ ) $_{n \geq 0}$  is defined over an arbitrary ring  $R$ .

The latter properties relate to types  $\text{FP}_n$  and types  $\text{F}_n$  as follows.

<sup>16</sup> The definition of such an object can be found in Corob Cook (2017).

- A t.d.l.c. group  $G$  is of type  $FP_n$  over  $\mathbb{Z}$  if and only if it is of type  $KP_n$  (see (Castellano and Corob Cook, 2020, Theorem 3.10)).
- If  $G$  has type  $F_n$  then it has type  $K_n$  (see (Castellano and Corob Cook, 2020, Theorem 3.23)).

*Open Problem* Despite the abundance of finiteness properties that are available in the TDLC context, the theory of finiteness conditions for TDLC-groups is still much less developed than the one for discrete groups. Moreover, very little is known about the relation (if there exists one) among the properties of different sequences  $CP_n, FP_n$  and  $KP_n$  (resp.  $C_n, F_n$  and  $K_n$ ). For example, is it true that type  $K_n$  implies type  $F_n$ ? At this stage, nothing is known on the relation to the sequences  $CP_n/C_n$ . A relation of the property  $C_\infty$  to the  $F_\infty$ -property has been promised by Sauer and Thumann (2017) to be discussed in forthcoming work of the authors.

*Open Problem* It would be relevant to find an example of a TDLC-group of type  $FP_2$  over  $\mathbb{Q}$  which is not compactly presented and it is “sufficiently” non-discrete (for example, it is not quasi-isometric to a discrete group). Unfortunately, the strategy developed in Bestvina and Brady (1997) does not seem to have a TDLC-analogue.

*Open Problem* For a TDLC-group  $G$  one can introduce several homological invariants. Which homological invariants are geometric? In 1991 Gromov asked whether the cohomological dimension of discrete groups is a geometric invariant. Under additional finiteness assumptions, Gersten (1993) proved that quasi-isometric groups have the same cohomological dimension and Sauer (2003) removed the finiteness assumptions in the case of rational cohomological dimension. This yields the natural question: is the rational discrete cohomological dimension defined in Castellano and Weigel (2016) a geometric invariant of a TDLC-group?

## 1.5 Willis’ theory of TDLC-groups: a sketch

As mentioned at the end of the introduction, Willis’ theory of the scale function was a fundamental breakthrough in the theory of TDLC-groups after several years of stillness and it motivated a vast research interest in the theory of TDLC-group which led to a systematic study of the class of compactly generated, topologically simple TDLC-groups that are non-discrete. In this section we confine ourselves to a brief introduction of the scale function and to some comments on (topologically) simple

TDLC-groups. Nevertheless, the student that is approaching the study of TDLC-groups is encouraged to dig further into this fundamental subject.

### 1.5.1 Scale function and tidy subgroups

Let  $\alpha \in \text{Aut}(G)$  and  $U$  be a compact open subgroup of  $G$ . Then

$$[\alpha(U) : U \cap \alpha(U)] < \infty,$$

because  $U \cap \alpha(U)$  is open in the compact set  $U$ . The **scale of**  $\alpha$  is

$$s(\alpha) = \inf\{[\alpha(U) : U \cap \alpha(U)] \mid U \text{ compact open subgroup of } G\}.$$

A subgroup  $U$  is **tidy** for  $\alpha$  if the infimum is attained at  $U$ . Tidy subgroups for  $\alpha$  always exists since  $s(\alpha)$  is the minimum of a subset of  $\mathbb{N}$ . Every tidy subgroup  $U$  can be decomposed as the product of a subgroup where  $\alpha$  expands and a subgroup where  $\alpha$  shrinks:

$$\text{if } U_{\pm} := \bigcap_{k>0} \alpha^{\pm k}(U), \text{ then } U = U_+ U_-.$$

By construction,  $U_+$  and  $U_-$  are closed subgroups,  $\alpha(U_+) \geq U_+$  and, similarly,  $\alpha(U_-) \leq U_-$ . Moreover, it can be shown that  $s(\alpha)$  represents the factor by which  $\alpha$  expands  $U_+$ , i.e.,  $s(\alpha) = [\alpha(U_+) : U_+]$ .

A striking result in the theory of the scale is the existence of an algorithm, the so-called **tidying procedure**, which produces a tidy subgroup when the input is an arbitrary compact open subgroup.

**Definition 1.39** The **scale function** of  $G$  is defined as the map

$$s : G \rightarrow \mathbb{Z}^+, \quad x \mapsto s(\alpha_x), \quad x \in G,$$

where  $\alpha_x$  denotes the inner automorphism  $y \mapsto xyx^{-1}$ .

The scale function  $s$  is known to satisfy the following properties (see Willis (1994, 2001a)):

- (s1)  $s$  is continuous if  $\mathbb{Z}^+$  carries the discrete topology;
- (s2)  $s(x) = 1 = s(x^{-1})$  if and only if there is a compact open subgroup  $U$  such that  $xUx^{-1} = U$ ;
- (s3)  $s(x^n) = s(x)^n$ , for every  $x \in G$  and  $n \geq 0$ ;
- (s4)  $\Delta(x) = s(x)/s(x^{-1})$ , where  $\Delta : G \rightarrow \mathbb{Q}^+$  is the modular function;
- (s5)  $s(\alpha(x)) = s(x)$  for every  $x \in G$  and  $\alpha \in \text{Aut}(G)$ .

The scale function encodes structural information of the group  $G$ . For a summary on the scale function (which in particular highlights the fact that tidy subgroups for automorphisms of TDLC-groups are analogues

of the Jordan canonical form of linear transformations) the reader is referred to Willis (2008) and references there.

*Remark* In the last years, Willis' theory has been investigated from different points of view bringing new approaches to the scale function of a TDLC-group. For example, Möller (2002) offers an interpretation of the fundamental ingredients of Willis' theory (that are tidy subgroups and scale function) in the setting of permutation group theory. Another example is given by the work initiated in Berlai et al. (2013), where Willis' topological dynamics of automorphisms has been reformulated in terms of the long-known theory of topological entropy.

### 1.5.2 Comments on simple TDLC-groups

Simple groups play an important role in group theory as the “indecomposable factors”. Indeed several types of simple groups have been completely classified; for instance, the simple finite groups and the simple connected Lie groups. Long-known classes of simple TDLC-groups are the class of simple Lie groups over local fields (see Bourbaki (2002)) and the class of automorphism groups of trees (see Tits (1970)).

In the realm of simple TDLC-groups, it is necessary a distinction between topological simplicity (i.e., every closed normal subgroup is trivial) and abstract simplicity (i.e., the underlying abstract group is simple). Examples show that a topologically simple TDLC-group can fail to be abstractly simple, see Willis (2007), but no example is known of topologically simple compactly generated TDLC-group that fail to be abstractly simple. Among compactly generated TDLC-groups, Smith (2017) showed that there are  $2^{\aleph_0}$  non-isomorphic compactly generated abstractly simple TDLC-groups. Nowadays, a classification of compactly generated topologically simple TDLC-groups is probably the best that can be hoped for.

**Scale function:** The theory of the scale produces invariants that could be important tools in the classification. For example,

- The set of values of the scale function: if  $G$  is compactly generated, the set of prime divisors of the values of the scale is finite (see Willis (2001b)); this set could distinguish between compactly generated simple TDLC-groups.
- The flat-rank: a notion of rank for TDLC-groups which is defined via

a notion of distance on the space of compact open subgroups (see Baumgartner et al. (2007)).

*Remark* In some cases, the flat-rank can be related to a cohomological invariant as shown in Castellano and Weigel (2016).

**Local-to-Global principle:** It relates the global properties of a compactly generated topologically simple TDLC-group with the structural properties of its compact open subgroups. This approach was initiated by

- Willis (2007) who showed that simplicity effects the local structure of the compactly generated TDLC-group: if  $G$  is compactly generated and topologically simple, then no compact open subgroup is solvable;
- Barnea et al. (2011) who addressed the question of which profinite groups can occur as compact open subgroups of compactly generated topologically simple TDLC-groups.

**Decomposition theory:** It includes methods for “breaking” a given TDLC-group into smaller (simple) pieces; see Caprace and Monod (2011). This approach has been successful for several classes of groups; for example, finite groups, profinite groups and algebraic groups. Therefore, one would hope to obtain analogous results for TDLC-groups.

General decomposition results has been obtained by Reid and Wesolek (2018b,a) who made use of the theory of *elementary groups* introduced by Wesolek (2015). Elementary groups are TDLC-groups that are built out of discrete and compact pieces.

**Geometrisation:** Cayley-Abels graphs allow the study of compactly generated TDLC-groups from a geometric perspective. Often a TDLC-group admits other types of geometric objects, e.g., *buildings*, that one can consider. For example, semi-simple Lie groups over a local field act on *affine buildings*, and also on *spherical buildings* (e.g., the affine building of  $\mathrm{SL}_2(\mathbb{Q}_p)$  is a tree and the spherical building is its boundary). Buildings are able to determine some properties of the group they are associated to. For instance, in some cases, they facilitate the computation of the Euler characteristic which turns out to have a curious connection with the value of a zeta-function in  $-1$  (see Castellano et al. (In preparation)).

Since profinite groups are trivial as geometric objects, the geometric behaviour of compactly generated TDLC-groups is often related to the

geometric behaviour of discrete groups. There is an ongoing programme of studying geometric properties of TDLC-groups by analogy with discrete groups. The aim is to understand to what extent long-known results on discrete groups find an analogue in the framework of TDLC-groups; see, for example, Castellano et al. (2022); Krön and Möller (2008); Arora et al. (2021); Cornulier and de la Harpe (2016).

## References

- Abels, H. 1973/74. Specker-Kompaktifizierungen von lokal kompakten topologischen Gruppen. *Math. Z.*, **135**, 325–361.
- Arora, S., Castellano, I., Corob Cook, G., and Martínez-Pedroza, E. 2021. Subgroups, hyperbolicity and cohomological dimension for totally disconnected locally compact groups. *Journal of Topology and Analysis*, 1–27.
- Außenhofer, L., Dikranjan, D., and Giordano Bruno, A. 2021. *Topological Groups and the Pontryagin-van Kampen Duality: An Introduction*. Vol. 83. Walter de Gruyter GmbH & Co KG.
- Barnea, Y., Ershov, M., and Weigel, Th. 2011. Abstract commensurators of profinite groups. *Trans. Amer. Math. Soc.*, 5381–5417.
- Baumgartner, U., Rémy, B., and Willis, G. A. 2007. Flat rank of automorphism groups of buildings. *Transformation Groups*, **12**(3), 413–436.
- Berlai, F., Dikranjan, D., and Giordano Bruno, A. 2013. Scale function vs topological entropy. *Topology Appl.*, 2314–2334.
- Bestvina, M., and Brady, N. 1997. Morse theory and finiteness properties of groups. *Invent. Math.*, 445–470.
- Bode, A., and Dupr, N. 2021. Locally analytic representations of  $p$ -adic groups.
- Bourbaki, N. 1998. *General topology. Chapters 5–10*. Springer-Verlag, Berlin.
- Bourbaki, N. 2002. *Lie groups and Lie algebras. Chapters 4–6*. Springer-Verlag, Berlin.
- Brown, K. S. 1994. *Cohomology of groups*. Springer-Verlag, New York.
- Caprace, P.-E., and De Medts, T. 2011. Simple locally compact groups acting on trees and their germs of automorphisms. *Transform. Groups*, 375–411.
- Caprace, P.-E., and Monod, N. 2011. Decomposing locally compact groups into simple pieces. *Math. Proc. Cambridge Philos. Soc.*, 97–128.
- Caprace, P.-E., and Monod, N. 2018. Future directions in locally compact groups: a tentative problem list. In: *New directions in locally compact groups*. Cambridge Univ. Press, Cambridge.
- Castellano, I. 2020. Rational discrete first degree cohomology for totally disconnected locally compact groups. *Math. Proc. Cambridge Philos. Soc.*, 361–377.

- Castellano, I., and Corob Cook, G. 2020. Finiteness properties of totally disconnected locally compact groups. *J. Algebra*, 54–97.
- Castellano, I., and Weigel, Th. 2016. Rational discrete cohomology for totally disconnected locally compact groups. *J. Algebra*, 101–159.
- Castellano, I., Marchionna, B., and Weigel, Th. 2022. A Stallings-Swan theorem for unimodular totally disconnected locally compact groups. *arXiv preprint arXiv:2201.10847*.
- Castellano, I., Chinello, G., and Weigel, Th. In preparation. The Hattori-Stallings rank, the Euler characteristic and the  $\zeta$ -functions of a totally disconnected locally compact group.
- Cornulier, Y. 2019. Locally compact wreath products. *Journal of the Australian Mathematical Society*, **107**(1), 2652.
- Cornulier, Y., and de la Harpe, P. 2016. *Metric geometry of locally compact groups*. EMS Tracts in Mathematics, vol. 25. European Mathematical Society (EMS), Zürich. Winner of the 2016 EMS Monograph Award.
- Corob Cook, G. 2017. Homotopical algebra in categories with enough projectives. *arXiv preprint arXiv:1703.00569*.
- Dikranjan, D. 2018. Introduction to topological groups. <http://users.dimi.uniud.it/~dikran.dikranjan/ITG.pdf>.
- Gersten, S. 1993. Quasi-isometry invariance of cohomological dimension.
- Gleason, A. M. 1952. Groups without small subgroups. *Annals of mathematics*, 193–212.
- Gromov, M. 1984. Infinite groups as geometric objects. Pages 385–392 of: *Proceedings of the International Congress of Mathematicians*, vol. 1.
- Hewitt, E., and Ross, K. A. 2012. *Abstract Harmonic Analysis: Volume I Structure of Topological Groups Integration Theory Group Representations*. Springer Science and Business Media.
- Hofmann, K. H. 1980. A note on Baire spaces and continuous lattices. *Bulletin of the Australian Mathematical Society*, **21**(2), 265–279.
- Hofmann, K. H., and Morris, S. A. 2007. *The Lie theory of connected pro-Lie groups: a structure theory for pro-Lie algebras, pro-Lie groups, and connected locally compact groups*. Vol. 2. European Mathematical Society.
- Karrass, A., and Solitar, D. 1956. Some remarks on the infinite symmetric groups. *Math. Z.*
- Krön, B., and Möller, R. G. 2008. Analogues of Cayley graphs for topological groups. *Math. Z.*, 637–675.
- Le Boudec, A. 2017. Compact presentability of tree almost automorphism groups. *Annales de l'Institut Fourier*.
- Möller, R. G. 2002. Structure Theory of Totally Disconnected Locally Compact Groups via Graphs and Permutations. *Canadian Journal of Mathematics*.
- Montgomery, D., and Zippin, L. 1952. Small subgroups of finite-dimensional groups. *Proceedings of the National Academy of Sciences of the United States of America*, **38**(5), 440.
- Neretin, Y. A. 1992. On combinatorial analogs of the group of diffeomorphisms of the circle. *Izvestiya Rossiiskoi Akademii Nauk*.

- Palmer, T. W. 1994. *Banach Algebras and the General Theory of \*-Algebras: Volume 2, \*-Algebras*. Cambridge university press.
- Reid, C. D., and Wesolek, P. R. 2018a. Dense normal subgroups and chief factors in locally compact groups. *Proceedings of the London Mathematical Society*.
- Reid, C. D., and Wesolek, P. R. 2018b. The essentially chief series of a compactly generated locally compact group. *Mathematische Annalen*.
- Ribes, L., and Zalesskii, P. 2000. Profinite groups. Pages 19–77 of: *Profinite Groups*. Springer.
- Sauer, R. 2003. Homological Invariants and Quasi-Isometry. *Geometric & Functional Analysis GAFA*, **16**, 476–515.
- Sauer, R., and Thumann, W. 2017. Topological models of finite type for tree almost automorphism groups. *International Mathematics Research Notices*, **2017**(23), 7292–7320.
- Serre, Jean-Pierre. 1980. SL 2. Pages 69–136 of: *Trees*. Springer.
- Smith, S. M. 2017. A product for permutation groups and topological groups. *Duke Mathematical Journal*.
- Tao, T. 2011. What’s new. <https://terrytao.wordpress.com/2011/10/08/254a-notes-5-the-structure-of-locally-compact-groups-and-hilberts-fifth-problem/>.
- Tao, T. 2014. *Hilbert’s fifth problem and related topics*. American Mathematical Soc.
- Tits, J. 1970. Sur le groupe des automorphismes d’un arbre. In: *Essays on topology and related topics*. Springer.
- Van Dantzig, D. 1931. *Studien over topologische algebra*. Ph.D. thesis, HJ Paris Amsterdam.
- Wesolek, P. R. 2015. Elementary totally disconnected locally compact groups. *Proceedings of the London Mathematical Society*.
- Wesolek, P. R. 2018. *An introduction to totally disconnected locally compact groups*.
- Willis, G. A. 1994. The structure of totally disconnected, locally compact groups. *Mathematische Annalen*.
- Willis, G. A. 2001a. Further properties of the scale function on a totally disconnected group. *Journal of Algebra*.
- Willis, G. A. 2001b. The number of prime factors of the scale function on a compactly generated group is finite. *Bulletin of the London Mathematical Society*, **33**(2), 168–174.
- Willis, G. A. 2007. Compact open subgroups in simple totally disconnected groups. *Journal of Algebra*.
- Willis, G. A. 2008. *A canonical form for automorphisms of totally disconnected locally compact groups*. De Gruyter.
- Wilson, J. S. 1998. *Profinite groups*. Clarendon Press.
- Wood, D. R., De Gier, J., Praeger, C. E., and Tao, T. 2018. *2016 MATRIX Annals*. Springer.
- Yamabe, H. 1953. A generalization of a theorem of Gleason. *Annals of Mathematics*, 351–365.

