

A large multi-agent system with noise both in position and control

Original

A large multi-agent system with noise both in position and control / D'Onofrio, G., Melchor Hernandez, A.. - In: ESAIM-CONTROL OPTIMISATION AND CALCULUS OF VARIATIONS. - ISSN 1262-3377. - 32:(2026), pp. 1-33.
[10.1051/cocv/2026003]

Availability:

This version is available at: 11583/3008417 since: 2026-03-09T11:45:31Z

Publisher:

Edp Sciences

Published

DOI:10.1051/cocv/2026003

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

A LARGE MULTI-AGENT SYSTEM WITH NOISE BOTH IN POSITION AND CONTROL

GIUSEPPE D'ONOFRIO¹  AND ANDERSON MELCHOR HERNANDEZ^{2,*}

Abstract. In this work, we consider a multi-population system where the dynamics of each agent evolve according to a system of stochastic differential equations in a general functional setup, determined by the global state of the system. Each agent is associated with a probability measure, that assigns the label accounting for the population to which the agent belongs. We do not assume any prior knowledge of the label of a single agent, and we allow that it can change as a consequence of the interaction among the agents. Furthermore, the system is affected by noise both in the agent's position and labels. First, we study the well-posedness of such a system and then a mean-field limit, as the number of agents diverges, is investigated together with the analysis of the properties of the limit distribution both with Eulerian and Lagrangian perspectives. As an application, we consider a large network of interacting neurons with random synaptic weights, introducing resets in the dynamics.

Mathematics Subject Classification. 60B10, 60H10, 93E03, 49N80, 60J70.

Received July 19, 2025. Accepted January 2, 2026.

1. INTRODUCTION

Mean-field analysis stands as a formidable mathematical tool employed to investigate large-scale systems characterized by a multitude of interacting components. These components may take the form of particles, agents, or individuals, and the methodology behind mean-field analysis allows the study of emergent behaviors at a collective level. Over the years, mean-field analysis has evolved into a versatile tool, finding applications across a diverse range of scientific disciplines. In physics, it has been instrumental in understanding phase transitions and collective phenomena in self-driven particle systems, as exemplified by the work of Vicsek *et al.* [1]. In the domain of economics, mean-field game theory, as developed by Lasry and Lions [2], provides a framework for analyzing strategic interactions among a large number of agents. Moreover, the insights garnered from mean-field analysis have proven invaluable in the field of neuroscience, enabling the study of emergent properties arising from the complex interactions of a large number of neurons (see for instance [3–7] just to name a few). More recently, these tools are being used in the context of machine learning and artificial neural networks [8, 9]. In economics, it is customary to refer to mean field games, where the term denotes the analysis of the average behavior of a large number of players or agents. The evolution of such models is typically studied

Keywords and phrases: Mean-field limit, Wasserstein distance, well-posedness of stochastic differential equations, Eulerian and Lagrangian solutions, neuronal modeling, random synaptic weights.

¹ Dipartimento di Scienze Matematiche “G. L. Lagrange”, Politecnico di Torino, Corso Duca degli Abruzzi 24, 10129 Torino, Italy.

² Dipartimento di Matematica, Università di Bologna, Via Zamboni 33, 40126, Bologna, Italy.

* Corresponding author: anderson.melchor@unibo.it

using the so-called Hamilton–Jacobi–Bellman equations, often coupled with a Fokker–Planck type equation [10, 11]. In all these models, the interacting agents or individuals are assumed to belong to different species without explicit spatial dependency. In most cases, this assumption arises due to the low abundances of agents or individuals, which can disrupt the convergence towards a limit model or introduce stochastic properties in the final model, as discussed in [12], Remark 2.7.

In this work, we consider a stochastic population model where an agent located at position $x \in \mathbb{R}^d$ is associated with a probability measure λ supported in some compact metric space (U, d_U) . Usually, we represent U as a set of labels accounting for the population to which the agent belongs. In particular, to describe the time evolution of the system, we consider for each agent i a stochastic differential equation (SDE) of the following form:

$$dX^i(t) = v_{\Lambda_t^N}(X^i(t), \lambda^i(t))dt + \sqrt{2\sigma}dB^i(t) \quad (1.1)$$

where the position X^i at time t is driven by a velocity field $v_{\Lambda_t^N}$, $t \in [0, T]$, where Λ_t^N is the empirical measure $\frac{1}{N} \sum_{i=1}^N \delta_{(X^i(t), \lambda^i(t))}$ and λ^i is the probability distribution of labels for the agent i . Furthermore, the dynamics of λ^i are given by the differential equation

$$d\lambda^i(t) = \mathcal{J}_{\Lambda_t^N}(X^i(t), \lambda^i(t))dt, \quad (1.2)$$

where $\mathcal{J}_{\Lambda_t^N}$ is a driving operator (the precise setting of the involved functional spaces is given in Section 2). The term λ^i expresses the probability that the agent i , at a certain moment in time, has a certain label, *i.e.* belongs to a subset of U . Indeed, we do not assume any prior knowledge of the label of a single agent, and we allow that it can change as a consequence of the interaction among the agents, like in [13]. Recently, in [14], the authors studied a model similar to (1.1). However, in their study, $\lambda^i(t)$ is given as the solution of non-local continuity equation. In such a situation, the pair $(\text{law}(X^i(t)), \lambda^i(t))$ represents a system of agents divided into two categories. The probability law of $X^i(t)$ represents an uncontrolled population called followers, while $\lambda^i(t)$ represents the controlled population called leaders. It is important to notice that in our case, the effect of the noise in the variable $X^i(t)$ affects the behavior of $\lambda^i(t)$ by means of the velocity field \mathcal{J} , and thus we need to study the well-posedness of the system given by (1.1)–(1.2). For a similar model without diffusion, we refer to [15]. The idea of a multi-agent model, where individuals are equipped with strategies and influenced by diffusive effects, was previously explored in [16] (see also [17]) as an extension of [18]. Here, we develop the study more in the spirit of [13]. In model (1.1)–(1.2), we consider the sequence $(B^i(t))_{i \in \mathbb{N}}$, $t \in [0, T]$, as a set of independent Brownian motions. The characteristic feature of this model is that, in the limit, we obtain a continuity equation where a diffusive part is developed. Here, we assume $\sigma > 0$, as the case $\sigma = 0$ has already been studied in some previous works, such as [13, 18, 19]. Furthermore, for the sake of generality, we are mainly concerned with local regularity properties of both $v_{\Lambda_t^N}$, $\mathcal{J}_{\Lambda_t^N}$, that is, both fields do not admit a global Lipschitz constant, and thus we need to enforce our hypotheses by considering a further sublinearity property of the fields in the involved arguments.

In the mean-field limit, we study also the time-regularity of the limit of the sequence of the empirical measures as a solution of an evolution equation. In doing that, we make use of the concept of Eulerian and Lagrangian solutions.

The model (1.1)–(1.2) straightforwardly can be used for various applications just reinterpreting the role of the agents and the controls (labels). As an applications of the previous model we consider a system of interacting neurons whose synaptic weights, that describe the influence that a neuron has onto the others, follow a certain distribution in the spirit of neuronal networks with random synaptic weights [20–22]. In this context, that belongs to the class of noisy integrate-and-fire models, the stochastic process X^i describes the time evolution of the membrane potential of a neuron receiving the inputs from the other neurons in the network it is embedded in. These contributions are summarized in the drift part of (1.1) according to the velocity field $v_{\Lambda_t^N}$ until the membrane potential reaches a physiological value X_F triggering a spike. As a result of this spike, which is assumed to occur instantaneously, the membrane potential of all the other connected neurons receives

a contribution. Afterward, the dynamics of the spiking neuron restart from the resting state X_R . The strength of the interaction/contribution between neurons and its evolution is described through a general operator $\mathcal{T}_{\Lambda_t^N}$ that is perturbed by a random effect of the environment, being it dependent on X^i .

It is important to note that model (1.1) assumes that the global dynamics of X^i does not exhibit resets. Therefore, for neuronal modeling, here we include the feature of the spike generation; that is, we consider the system

$$\left\{ \begin{array}{l} X^i(t) = X_0^i + \int_0^t v_{\Lambda_s^N}(X^i(s), \lambda^i(s)) ds + \sqrt{2\sigma} B_t^i, \text{ if } X^i(t^-) < X_F, \\ X^i(t) = X_R, \\ \lambda^i(t) = \lambda_0^i + \int_0^t \mathcal{T}_{\Lambda_s^N}(X^i(s), \lambda^i(s)) ds, \end{array} \right. \quad \text{if } X^i(t^-) = X_F, \quad (1.3)$$

where X_F is an additional parameter describing the spiking threshold and X_R is the reset value (resting potential).

Here, we are particularly interested in the following case. Let us consider a measurable function $\alpha : U \times \mathcal{Y} \rightarrow \mathbb{R}$, and define $\mathcal{T} : \mathcal{Y} \times \mathcal{P}_1(\mathcal{Y}) \rightarrow \mathcal{F}(U)$ as follows:

$$\begin{aligned} \mathcal{T}(y, \Psi) &:= \int_{\mathcal{Y}} \alpha(\cdot, z) d\Psi(z), \quad \text{such that } \int_U \int_{\mathcal{Y}} \alpha(u, z) d\Psi(z) d\mu(u) = 0, \\ \mu(u) &= \sum_{k \geq 1} \beta_k \delta_{u_k}, \quad u_k \in U, \quad \beta_k \in \mathbb{R}, \end{aligned}$$

where $\mu \in \mathcal{M}(U)$ defines a signed measure. Furthermore, we consider a Lipschitz velocity field $b : \mathbb{R}^d \rightarrow \mathbb{R}^d$, and set $v_{\mathcal{T}} : \mathcal{Y} \rightarrow \mathbb{R}^d$, $(x, \lambda) \mapsto b(x)$. The functional form of \mathcal{T} is of great generality and contains, as special cases, the random synaptic weights seen as constants, Bernoulli or Gaussian random variables considered in previous works [20, 22–24].

Moreover, since weight distributions in real cortex have been observed to be broad, neurons can belong to different subgroups with different behaviors. The variance of the temporal fluctuations of the input currents may be large enough to endow the network with its own source of variability (see [25] for an exhaustive review). Therefore, in Section 4.1, we consider a heterogeneous population of neurons. In mathematical terms, this translates in an additional source of noise in equation (1.2), namely

$$\lambda^i(t) = \lambda_0^i + \int_0^t \mathcal{T}_{\Lambda_s^N}(X^i(s), \lambda^i(s) + R(s)) ds,$$

with $R(t) := \sum_{h \in \mathbb{N}} a_h W_h(t) e_h$, where $(W_h(t))_{h \in \mathbb{N}}$ is a sequence of independent Brownian motions, $(a_h)_{h \in \mathbb{N}}$ is a sequence of non-negative numbers such that $\sum_h a_h^2 < +\infty$, and $(e_h)_h$ is a proper sequence of signed measures.

For all these models, we prove the well-posedness of the coupled SDEs and discuss the link between Eulerian and Lagrangian solutions of (1.1)–(1.2). Finally, for (1.3) we show the qualitative behavior in the one-dimensional case using simulations.

Classical models of large neuronal network can be studied also using a diffusion approximation, which, however, requires that the frequency of the inputs is high, but their amplitude is small (see [26]). Mean-field theory is a device to analyze the collective behavior of such a dynamical system without these requests as long as the number of interacting particles is high. The theory allows to reduce the behavior of the system to the properties of a handful of parameters. In neural circuits, these parameters are typically the firing rates. In this context, we are often interested in the properties averaged across the distribution of possible weights [27–29] and

in the limit of infinite network size [30, 31]. Some important properties, such as the existence of a sharp phase transition or synchronization, are only obtainable in this limit. However, fluctuations are sometimes so high that care must be taken in the averaging process, as in the case of multiplicative noise; see [32]. For this reason we introduce sources of noise in the network itself. This goes also in the direction of addressing the modeling of synaptic plasticity, where the synaptic weight itself is a random process evolving in time, as suggested in [33].

The paper is organized as follows: in Section 2 we introduce our notation and we recall some preliminary results. We describe the multi-agent system described by coupled SDEs with noise both in position and labels along with the relative assumptions and a description of the spaces in which we are working. The main results on the well-posedness of the considered system, along with comments on the properties of the mean-field limit, are stated and proven in Section 3. In particular, we develop the mean-field description using both the Eulerian and Lagrangian approaches to study the time regularity of the limit distribution. Finally, in Section 4, we consider a large network of interacting neurons whose activity is described by the aforementioned system, both for homogeneous and heterogeneous populations of neurons. This model exhibits jumps, and the corresponding well-posedness results are provided.

2. THE MODEL

In this section, we will provide a detailed overview of the model, including a comprehensive description of all the tools used and the key assumptions it is based on. In details, Subsection 2.1 introduces some preliminaries and notation, while in Subsection 2.2 we provide the functional spaces needed for our purpose, and in Subsection 2.3 our assumptions. Furthermore, in Subsection 2.4, we recall the notion of Eulerian and Lagrangian solutions. Subsection 2.5 is dedicated to the particle system and to the notion of strong solution for SDEs.

2.1. Preliminaries and notation

Let (X, d_X) be a metric space, we denote by $\mathcal{M}(X)$ the space of signed Borel measures with finite total variation, by $\mathcal{M}_+(X)$ and $\mathcal{P}(X)$ the convex subsets of nonnegative measures and probability measures, respectively. For $\sigma \in \mathcal{M}(X)$, $|\sigma| \in \mathcal{M}_+(X)$ denotes the total variation measure of σ . We shall also use the notation $\mathcal{M}_0(X)$ for the subset of measures with zero mean. Given a metric space (X, d_X) , we consider the Lipschitz space

$$\text{Lip}(X, d_X) := \{\phi : X \rightarrow \mathbb{R} \mid \exists L > 0 : \forall x, y \in X \ |\phi(x) - \phi(y)| \leq L d_X(x, y)\}.$$

For a continuous function $\phi \in C(X)$ we denote by

$$\text{Lip}(\phi) := \sup_{\substack{x, y \in X \\ x \neq y}} \frac{|\phi(x) - \phi(y)|}{d_X(x, y)}$$

its Lipschitz constant. In the next, we endow $\text{Lip}(X, d_X)$ with the Lipschitz norm $\|\phi\|_{\text{Lip}} := \sup_{x \in X} |\phi(x)| + \text{Lip}(\phi)$. Given a measure $\mu \in \mathcal{M}_+(X)$, (E, d_E) another metric space, and $f : X \rightarrow E$ a μ -measurable function, we shall denote by $f_{\#} \in \mathcal{M}_+(E)$ the push-forward measure, having the same mass as μ and defined by $f_{\#}\mu(B) = \mu(f^{-1}(B))$ for any Borel set $B \subset E$. We often use the change of variable formula

$$\int_E g d f_{\#}\mu = \int_X g \circ f d\mu$$

whenever either one of the integrals makes sense. In a complete and separable metric space (X, d_X) , we shall use the Kantorovich–Rubinstein distance $\mathcal{W}_1(\cdot, \cdot)$ in the class $\mathcal{P}(X)$. For $\mu_1, \mu_2 \in \mathcal{M}_1(X)$, the 1-Wasserstein

distance $\mathcal{W}_1(\mu_1, \mu_2)$ is defined by

$$\mathcal{W}_1(\mu_1, \mu_2) := \inf \left\{ \int_{X \times X} d_X(x_1, x_2) \gamma(dx_1, dx_2) \mid \gamma \in \Gamma(\mu_1, \mu_2) \right\}$$

where $\Gamma(\mu_1, \mu_2)$ is the set of admissible coupling between μ_1 and μ_2 . It is worth to recall that due to Kantorovich duality, one can also consider the following definition

$$\mathcal{W}_1(\mu_1, \mu_2) := \sup \left\{ \int_X \phi d(\mu_1 - \mu_2) \mid \phi \in \text{Lip}(X, d_X), \text{Lip}(\phi) \leq 1 \right\}.$$

Notice that $\mathcal{W}_1(\mu_1, \nu_1)$ is finite if μ_1, ν_1 belong to the space

$$\mathcal{P}_1(X) := \left\{ \mu \in \mathcal{P}(X) \mid \int_X d_X(x, \bar{x}) d\mu(x) < \infty \text{ for some } \bar{x} \in X \right\}.$$

Note that $(\mathcal{P}_1(X), \mathcal{W}_1)$ is complete if (X, d_X) is complete. Moreover, by [34], Theorem 2.2.1 the following holds true: a sequence $(\mu_n) \subset \mathcal{P}_1(X)$ converges to $\mu \in \mathcal{P}_1(X)$ with respect to the Wasserstein distance \mathcal{W}_1 if and only if, for all $\phi \in \text{Lip}(X, d_X)$,

$$\int_X \phi d\mu_n \xrightarrow{n \rightarrow \infty} \int_X \phi d\mu, \quad \int_X d_X(\cdot, \bar{x}) d\mu_n \xrightarrow{n \rightarrow \infty} \int_X d_X(\cdot, \bar{x}) d\mu.$$

Suppose that E is a Banach space, the notation $C_b^1(E)$ will be used to denote the subspace of $C_b(E)$ of functions having bounded continuous Fréchet differential at every point. Hence, we make use of the notation $D\varphi$ to denote such a Fréchet differential. Let $T > 0$, in the case $\varphi : [0, T] \times E \rightarrow \mathbb{R}$, we make use of the symbol $\frac{d}{dt}$ to denote the partial derivative with respect to the variable $t \in [0, T]$, and D will only stand for the derivative with respect to the variables in E .

2.2. The framework

In what follows, we are interested in studying SDEs on \mathcal{Y} , where $\mathcal{Y} := \mathbb{R}^d \times \mathcal{P}(U)$. The state space of our system is given by pairs $(x, \lambda) = y \in \mathcal{Y}$. Here $x \in \mathbb{R}^d$ denotes the spatial component of an agent, whereas the element $\lambda \in \mathcal{P}(U)$ denotes a probability distribution over the space U , which we assume to be a compact metric space. It can be interpreted as a space of strategies [13, 18].

We now consider a compact metric space (U, d_U) , and consider the set $\mathcal{F}(U)$ defined as

$$\mathcal{F}(U) := \overline{\text{span}(\mathcal{P}(U))}^{\|\cdot\|_{\text{BL}}}$$

as the closure in the dual space $(\text{Lip}(U, d_U))^*$ with respect to the dual norm

$$\|\ell\|_{\text{BL}} := \sup \left\{ \langle \ell, \phi \rangle : \phi \in \text{Lip}(U, d_U), \|\phi\|_{\text{Lip}} \leq 1 \right\}.$$

The space $\mathcal{F}(U)$ is the so-called Arens–Eells space introduced in [35]. This is a separable Banach space containing $\mathcal{M}(U)$. Furthermore, for a measure $\nu \in \mathcal{M}_0$, the $\|\cdot\|_{\text{BL}}$ -norm is equivalent to the norm induced by the dual formulation of the 1-Wasserstein distance.

On the other hand, note that $\mathcal{M}(U)$, when endowed with the total variation norm

$$\|\sigma\|_{\text{TV}} := \sup \left\{ \int_U \phi d\sigma : \phi \in C(U), |\phi| \leq 1 \right\}$$

has the structure of Banach space, isometrically isomorphic to the dual of $C(U)$. We will also use the representation formulas

$$\|\sigma\|_{\text{TV}} = |\sigma|(U) = \sup \left\{ \int_U \phi d\sigma : \phi \in B_b(U), |\phi| \leq 1 \right\}$$

where $B_b(U)$ denotes the class of bounded Borel functions $\phi : U \rightarrow \mathbb{R}$. Furthermore, we have by Kantorovich duality that

$$\|\mu_1 - \mu_2\|_{\text{BL}} \leq \mathcal{W}_1(\mu_1, \mu_2) \leq (1 + D_U) \|\mu_1 - \mu_2\|_{\text{BL}} \quad (2.1)$$

where $D_U := \min_{x_0 \in U} \max_{x_1 \in U} d_U(x_0, x_1) \leq \text{diam}(U)$, and $\text{diam}(U)$ denotes the diameter of U .

In the next we need to consider the space $\mathcal{E} := \mathbb{R}^d \times \mathcal{F}(U)$ endowed with the norm $\|y\|_{\mathcal{E}} = \|(x, \sigma)\|_{\mathcal{E}} := |x| + \|\sigma\|_{\text{BL}}$, which is a separable Banach space. For $y \in \mathcal{Y}$ and $\Psi \in \mathcal{P}_1(\mathcal{Y})$, we consider the vector field $b_\Psi : \mathcal{Y} \rightarrow \mathcal{E}$ defined as

$$b_\Psi(y) := \begin{pmatrix} v_\Psi(y) \\ \mathcal{T}_\Psi(y) \end{pmatrix}. \quad (2.2)$$

The first component of b_Ψ is a velocity field in \mathbb{R}^d determined by the global state of the system Ψ ; the second component is expressed in terms of an operator $\mathcal{T}_\Psi : \mathcal{Y} \rightarrow \mathcal{E}$ which sees the location of agents around Ψ .

For a given $R > 0$, we denote by B_R the closed ball of radius R in \mathbb{R}^d and by $B_R^{\mathcal{Y}}$ the ball of radius R in \mathcal{Y} , that is, $B_R^{\mathcal{Y}} := \{y \in \mathcal{Y} : \|y\|_{\mathcal{E}} \leq R\}$.

Remark 2.1. Since U is assumed compact, by [34], Corollary 2.2.5 $\mathcal{P}(U)$ is compact and thus \mathcal{Y} is a locally compact space. Therefore $B_R^{\mathcal{Y}}$ is a compact set. Since $\mathcal{Y} \subset \mathcal{E}$, for any $\Psi \in \mathcal{P}(\mathcal{Y})$ we may define

$$m_1(\Psi) := \int_{\mathcal{Y}} \|y\|_{\mathcal{E}} d\Psi.$$

Hence,

$$\mathcal{P}_1(\mathcal{Y}) = \{\Psi \in \mathcal{P}(\mathcal{Y}) : m_1(\Psi) < +\infty\}.$$

2.3. Assumptions

We assume that the velocity field $v_\Psi : \mathcal{Y} \rightarrow \mathbb{R}^d$ satisfies the following structural assumptions:

(A1) For every $R > 0$, for every $\Psi \in \mathcal{P}(B_R^{\mathcal{Y}})$, $v_\Psi \in \text{Lip}(B_R^{\mathcal{Y}}; \mathbb{R}^d)$ uniformly with respect to Ψ , that is, there exists a positive constant $L_{v,R}$ such that

$$|v_\Psi(y^1) - v_\Psi(y^2)| \leq L_{v,R} \|y^1 - y^2\|_{\mathcal{E}};$$

(A2) for every $R > 0$, for every $\Psi \in \mathcal{P}(B_R^{\mathcal{Y}})$, there exists a positive constant $L_{v,R}$ such that for every $y \in B_R^{\mathcal{Y}}$ and for every $\Psi^1, \Psi^2 \in \mathcal{P}(B_R^{\mathcal{Y}})$

$$|v_{\Psi^1}(y) - v_{\Psi^2}(y)| \leq L_{v,R} \mathcal{W}_1(\Psi^1, \Psi^2);$$

(A3) there exists $M_v > 0$ such that for every $y \in \mathcal{Y}$ and for every $\Psi \in \mathcal{P}_1(\mathcal{Y})$ there holds

$$|v_\Psi(y)| \leq M_v (1 + \|y\|_{\mathcal{E}} + m_1(\Psi)).$$

The assumptions on the operator \mathcal{T} are the following. For $\Psi \in \mathcal{P}_1(\mathcal{Y})$, the operator $\mathcal{T}_\Psi : \mathcal{Y} \rightarrow \mathcal{F}(U)$ is such that

(B1) for every $(y, \Psi) \in \mathcal{Y} \times \mathcal{P}_1(\mathcal{Y})$, constants belong to the kernel of $\mathcal{T}_\Psi(y)$, that is

$$\langle \mathcal{T}_\Psi(y), 1 \rangle_{\mathcal{F}(U), \text{Lip}(U)} = 0;$$

(B2) for every $(y, \Psi) \in \mathcal{Y} \times \mathcal{P}_1(\mathcal{Y})$, there exists a positive constant $M_{\mathcal{T}}$ such that

$$\|\mathcal{T}_\Psi(y)\|_{\text{BL}} \leq M_{\mathcal{T}}(1 + |x| + m_1(\Psi)), \text{ where } y = (x, \lambda);$$

(B3) for every $R > 0$ there exists a positive constant $L_{\mathcal{T}, R}$ such that for every $(y^1, \Psi^1), (y^2, \Psi^2) \in B_R^{\mathcal{Y}} \times \mathcal{P}(B_R^{\mathcal{Y}})$,

$$\|\mathcal{T}_{\Psi^1}(y^1) - \mathcal{T}_{\Psi^2}(y^2)\|_{\text{BL}} \leq L_{\mathcal{T}, R} (\|y^1 - y^2\|_{\mathcal{E}} + \mathcal{W}_1(\Psi^1, \Psi^2));$$

(B4) for every $R > 0$ there exists $\delta_R > 0$ such that for every $(y, \Psi) \in B_R^{\mathcal{Y}} \times \mathcal{P}_1(\mathcal{Y})$ we have

$$\mathcal{T}_\Psi(y) + \delta_R \lambda \geq 0, \text{ where } y = (x, \lambda).$$

Remark 2.2. Let us give some comments about our assumptions. Let us observe that by (A3) and (B2) $\mathcal{T}_\Psi(y)$ is continuous with respect to the BL-norm, and thus by [18], Section 2.2 the integral form in (1.3) for $\lambda^i(t)$ makes sense. More precisely, let us consider $0 \leq s < t$. Notice that for each $i = 1, \dots, N$, we have by (B2), and Jensen inequality that

$$\mathbb{E} \left\| \int_s^t \mathcal{T}_{\Psi_t^N}(X^i(r), \lambda^i(r)) dr \right\|_{\text{BL}}^2 \leq (t-s) \int_s^t \mathbb{E} (M_{\mathcal{T}} (1 + |X^i(r)| + m_1(\Lambda_r^N)))^2 dr.$$

Applying (A3) we get that $\mathbb{E} \int_0^r |X^i(s)|^2 ds \leq CT$ for some positive constant C (this follows from (3.7) below), and $m_1(\Lambda_r^N) < +\infty$ for all $0 \leq r \leq T$ (see (3.6) below), then

$$\mathbb{E} \left\| \int_s^t \mathcal{T}_{\Psi_t^N}(X^i(r), \lambda^i(r)) dr \right\|_{\text{BL}}^2 \leq CT(t-s)^2$$

for some positive constant C .

Let us observe that assumptions (B1), and (B4) are needed to ensure that the displacement $\lambda + \frac{1}{\delta_R} \mathcal{T}_\Psi(x, \lambda)$ defines a probability measure. Furthermore, note that $\mathcal{P}(U)$ is a convex weak-star closed subset of $\mathcal{M}(U)$, corresponding, via the dual representation of $\mathcal{M}(U)$, to the class of non-decreasing functionals $G : C(U) \rightarrow \mathbb{R}$ with $G(1) = 1$. Hence, we can extend $\lambda + \frac{1}{\delta_R} \mathcal{T}_\Psi(x, \lambda) : \text{Lip}(U) \rightarrow \mathbb{R}$ to a positive linear functional $\lambda + \frac{1}{\delta_R} \mathcal{T}_\Psi(x, \lambda) : C(U) \rightarrow \mathbb{R}$. More precisely, observe that for all $\psi \in (\text{Lip}(U))'$, the dual of $\text{Lip}(U)$, the duality pairing $\langle \cdot, \cdot \rangle$ satisfies

$$|\langle \psi, f \rangle| \leq \|f\|_\infty \langle \psi, 1 \rangle \text{ for all } f \in \text{Lip}(U).$$

Since $\text{Lip}(U)$ is dense in $C(U)$, we may take $\psi = \lambda + \frac{1}{\delta_R} \mathcal{T}_\Psi(x, \lambda)$, and extend it to $C(U)$ by the Riesz representation theorem; in particular, it belongs to $(C(U))'$. Assumption (B4) implies that $\langle \mathcal{T}_\Psi, 1 \rangle = 0$. Therefore $\psi(1) = \langle \lambda, 1 \rangle = 1$, and thus $\psi \in \mathcal{M}_+(U)$ is a probability measure. Therefore, based on this observation, we can obtain the well-posedness of (2.9) as a consequence of Proposition 2.3 below. It might seem that introducing the Banach space $\mathcal{F}(U)$ is unnecessary, since the trajectories of $\lambda(t)$ never leave $\mathcal{P}(U)$. However, the well-posedness of (2.9) relies on a theoretical result that is established only in Banach spaces; while $(\mathcal{P}_1(\mathcal{Y}), \mathcal{W}_1)$ is merely a metric space. For this reason, we extend the dynamics to \mathcal{E} in order to exploit this Banach structure, and we regard \mathcal{Y} as a convex closed subset of \mathcal{E} .

Proposition 2.3 ([13], Prop. 3.1). *For $y \in \mathcal{Y}$ and $\Psi \in \mathcal{P}_1(\mathcal{Y})$, define $b_\Psi(y)$ as in (2.2). Assume that $v_\Psi : \mathcal{Y} \rightarrow \mathbb{R}^d$ satisfies (A1)–(A3) and $\mathcal{T}_\Psi : \mathcal{Y} \rightarrow \mathcal{F}(U)$ satisfies (B1)–(B4). Then the following hold true:*

(i) *for every $R > 0$, for every $\Psi \in \mathcal{P}(B_R^\mathcal{Y})$, and for every $y^1, y^2 \in B_R^\mathcal{Y}$, there exists a positive constant L_R such that*

$$\|b_\Psi(y^1) - b_\Psi(y^2)\|_{\mathcal{E}} \leq L_R \|y^1 - y^2\|_{\mathcal{E}};$$

(ii) *for every $R > 0$; for every $\Psi^1, \Psi^2 \in \mathcal{P}(B_R^\mathcal{Y})$, and for every $y \in B_R^\mathcal{Y}$, there exists a positive constant L_R such that*

$$\|b_{\Psi^1}(y) - b_{\Psi^2}(y)\|_{\mathcal{E}} \leq L_R \mathcal{W}_1(\Psi^1, \Psi^2);$$

(iii) *for every $R > 0$, there exists $\theta > 0$ such that for every $y \in B_R^\mathcal{Y}$ and for every $\Psi \in \mathcal{P}(B_R^\mathcal{Y})$*

$$y + \theta b_\Psi(y) \in \mathcal{Y};$$

(iv) *there exists $M > 0$ such that for every $y \in \mathcal{Y}$ and for every $\Psi \in \mathcal{P}_1(\mathcal{Y})$ there holds*

$$\|b_\Psi(y)\|_{\mathcal{E}} \leq M(1 + \|y\|_{\mathcal{E}} + m_1(\Psi)).$$

2.4. Eulerian and Lagrangian solutions

Let us discuss the concepts of Eulerian and Lagrangian solutions in the context of studying the asymptotic behavior of a system like (1.1)–(1.2) as $N \rightarrow +\infty$. Let us first recall the notion of Eulerian solution. In the next, given a sufficiently smooth function $\varphi : \mathcal{Y} \rightarrow \mathbb{R}$, and generic $\Psi \in \mathcal{P}_1(\mathcal{Y})$ we define the operator

$$\mathcal{L}_\Psi \varphi(y) := \sigma \text{Tr}(D_{yy} \varphi(y)) + \langle b_\Psi, D_y \varphi(y) \rangle \quad (2.3)$$

where b_Ψ is defined according to (2.2). In the next, we make the following important assumption about \mathcal{L}_Ψ .

(C1) Let us suppose that there exists $r > 0$ such that for each $\Psi \in \mathcal{P}_1(B_r^\mathcal{Y})$, $\mathcal{L}_\Psi : \text{Dom}(\mathcal{L}_\Psi) \subset C_0(\mathcal{Y}) \rightarrow C_0(\mathcal{Y})$ is densely defined, a dissipative linear operator such that for some $\eta > 0$

$$\text{Range}(\eta I - \mathcal{L}_\Psi) = C_0(\mathcal{Y}).$$

Here, we denote by $C_0(\mathcal{Y})$ the space of continuous real-valued functions on \mathcal{Y} vanishing at infinity. That is, $\varphi \in C_0(\mathcal{Y})$ if it is continuous and for all $\varepsilon > 0$ there exists a compact set $K_\varepsilon \subset \mathcal{Y}$ such that $|\varphi(y)| < \varepsilon$ for all $y \notin K_\varepsilon$.

Definition 2.4. Let $\Lambda \in C^0([0, T]; (\mathcal{P}_1(\mathcal{Y}), \mathcal{W}_1))$ and let $\bar{\Lambda} \in \mathcal{P}(\mathcal{Y})$ be a given compactly supported initial condition. We say that Λ is an Eulerian solution to the parabolic equation

$$\frac{d}{dt} \Lambda_t + \mathcal{L}_{\Lambda_t}^*(\Lambda_t) = 0, \quad (2.4)$$

starting from $\bar{\Lambda}$ if and only if for every $\varphi \in C_b^1([0, T] \times \mathcal{E})$,

$$\int_{\mathcal{Y}} \varphi(t, y) d\Lambda_t(y) - \int_{\mathcal{Y}} \varphi(0, y) d\bar{\Lambda}(y) = \int_0^t \int_{\mathcal{Y}} \left(\frac{d}{ds} \varphi(s, y) + \mathcal{L}_{\Lambda_s} \varphi(s, y) \right) d\Lambda_s(y) ds, \quad (2.5)$$

where $\mathcal{L}_{\Lambda_t}^*$ denotes the dual operator of \mathcal{L}_{Λ_t} . Notice that for us φ will be always a function which does not depend on t , and then its time derivative is not needed in (2.5). Notice that our operator \mathcal{L}_{Ψ} can be related to an SDE.

Remark 2.5. Suppose that we have a generic Brownian motion $(\bar{B}(t))_{t \in [0, T]}$ defined in a complete probability space $(\Omega, \{\mathcal{F}_t\}_{t \in [0, T]}, \mathbb{P})$. Let us consider the equations

$$\begin{aligned} X(t) &= \bar{X}_0 + \int_0^t v_{\Lambda}(X(s), \lambda(s)) ds + \sqrt{2\sigma} \bar{B}(t), \\ \lambda(t) &= \bar{\lambda}_0 + \int_0^t \mathcal{T}_{\Lambda}(X(s), \lambda(s)) ds, \end{aligned} \tag{2.6}$$

where $X(0) = \bar{X}_0 \in L^2(\Omega; \mathbb{R}^d)$, $\lambda_0 \in L^2(\Omega, \mathcal{P}(U))$ are the initial conditions for (2.6). Since \mathcal{E} is a separable Banach space, thanks to Itô calculus, we can express \mathcal{L}_{Ψ} in a more specific manner. Indeed, suppose that $\varphi : \mathcal{Y} \rightarrow \mathbb{R}$ is a $C^2(\mathcal{Y})$ function. Then by the Itô formula for $d\varphi(X(t), \lambda(t))$ one has:

$$d\varphi(X(t), \lambda(t)) = \nabla_x \varphi(X(t), \lambda(t)) dX(t) + D_{\lambda} \varphi(X(t), \lambda(t)) d\lambda(t) + \sigma \Delta_x \varphi(X(t), \lambda(t)) dt$$

where D_{λ} denote the Frechét derivative with respect to the variable λ , and $(X(t), \lambda(t))$ is the solution of (2.6) above. By taking the integral with respect to the time variable, and the expected valued with respect to \mathbb{P} , we obtain that

$$\begin{aligned} \int_{\mathcal{Y}} \varphi(y) d\mu_t(y) - \int_{\mathcal{Y}} \varphi(y) d\mu_0(y) &= \int_0^t \int_{\mathcal{Y}} \langle D_y \varphi(y), b_{\Psi}(y) \rangle \mu_s(y) ds \\ &\quad + \sigma \int_0^t \int_{\mathcal{Y}} \Delta_x \varphi(y) d\mu_s(s) ds. \end{aligned}$$

Hence, we obtain

$$\mathcal{L}_{\Psi} \varphi(y) = \sigma \Delta_x \varphi(y) + \langle b_{\Psi}, D_y \varphi(y) \rangle.$$

Our assumption C1 guarantees the existence of a strongly continuous semigroup of contractions on $C_0(\mathcal{Y})$. In general, this can be proven when the first differential equation of (2.6), has the following structure

$$dX(t) = (AX(t) + v_{\Lambda}(X(t), \lambda(t))) dt + \sqrt{2\sigma} d\bar{B}(t),$$

where $A : \mathbb{R}^d \rightarrow \mathbb{R}^d$ is the infinitesimal generator of a strongly continuous semigroup. However, in our case $A = 0$, and thus results such as [36], Proposition 2.8, which establish that right processes and strongly continuous semigroups with generator \mathcal{L}_{Ψ} are associated in the sense of Dirichlet forms, cannot be applied. Moreover, results such as [37], Theorem 7.7, which establish the existence of solutions to (2.6), cannot be applied either.

Before recalling the definition of Lagrangian solution, we recall a further important concept.

Definition 2.6. Let us fix $T > 0$, $\bar{y} \in \mathcal{Y}$, and $(\bar{B}(t))_{t \in [0, T]}$ be a \mathbb{R}^d -valued Brownian motion. The transition map $\bar{F}_{\Lambda}(t, s, \bar{y})$ associated with the equation (2.6) above, where the initial condition is replaced by $(X(s), \lambda(s)) = \bar{y}$ is defined through

$$\bar{F}_{\Lambda}(t, s, \bar{y}) := (X(t), \lambda(t)), \tag{2.7}$$

where $(X(t), \lambda(t))_{t \in [0, T]}$ is the unique solution to equation (2.6) driven by $(\bar{B}(t))_{t \in [0, T]}$.

We now recall what a Lagrangian solution is for an equation of the form (2.4).

Definition 2.7. Suppose that $\Lambda \in C^0([0, T]; (\mathcal{P}_1(\mathcal{Y}), \mathcal{W}_1))$ and let $\bar{\Lambda}$ be any given initial condition for (2.4). We say that Λ is a Lagrangian solution to (2.4) starting from $\bar{\Lambda}$ if and only if it satisfies that

$$\Lambda_t = \bar{F}_\Lambda(t, 0, \cdot) \# \bar{\Lambda} \quad \text{for every } 0 \leq t \leq T, \mathbb{P}\text{-a.s.},$$

where $\bar{F}_\Lambda(t, s, y)$ are the transition maps as defined in (2.7).

Remark 2.8. Observe that the measure Λ_t can be seen as the action of the evaluation map ev_t which can be defined as follows. We set for each $t \in [0, T]$ the evolution map $\text{ev}_t : C^0([0, T]; (\mathcal{P}_1(\mathcal{Y}), \mathcal{W}_1)) \rightarrow \mathcal{P}_1(\mathcal{Y})$ as

$$\text{ev}_t(\Lambda) := \Lambda_t, \quad \text{for all } \Lambda \in C^0([0, T]; (\mathcal{P}_1(\mathcal{Y}), \mathcal{W}_1)). \quad (2.8)$$

In general, notice that the definition of the evolution map ev_t can be defined in $C^0([0, T]; E)$, where E is a generic Banach space, see [18].

2.5. The multi-agent system: Strong solutions for SDEs

In this part, we introduce the model and we recall the notion of strong solutions for SDEs. Let us fix $T > 0$, $\sigma > 0$, and for $N \in \mathbb{N}$ we consider N independent and identically distributed \mathbb{R}^d -valued Brownian motions $(B_t^i)_{t \in [0, T]}$. In what follows, we consider a particle system of N agents evolving according to

$$\begin{cases} X^i(t) = X_0^i + \int_0^t v_{\Lambda_s^N}(X^i(s), \lambda^i(s)) ds + \sqrt{2\sigma} B_t^i, \\ \lambda^i(t) = \lambda_0^i + \int_0^t \mathcal{T}_{\Lambda_s^N}(X^i(s), \lambda^i(s)) ds, \end{cases} \quad (2.9)$$

where $X^i(t) \in \mathbb{R}^d$, $\lambda^i(t) \in \mathcal{P}(U)$ for each $i = 1, \dots, N$, and

$$\Lambda_t^N := \frac{1}{N} \sum_{i=1}^N \delta_{(X^i(t), \lambda^i(t))} \quad (2.10)$$

is the empirical measure associated to the system. Since in this part N is fixed, we use Λ_t rather than Λ_t^N if confusion does not occur. We introduce the vector-valued variable $\mathbf{y} := (y^1, \dots, y^N) \in \mathcal{Y}^N \subset \mathcal{E}^N$, which we endow with the norm

$$\|\mathbf{y}\|_{\mathcal{E}^N} := \frac{1}{N} \sum_{i=1}^N \|y^i\|_{\mathcal{E}}.$$

The natural space to study the well posedness of the system described through (2.9) is \mathcal{E}^N . However, notice that $\hat{\mathcal{Y}}^N := (\mathbb{R}^d)^N \times (\mathcal{P}(U))^N$, and $\hat{\mathcal{E}}^N := (\mathbb{R}^d)^N \times (\mathcal{F}(U))^N$ can be also endowed with the same norm $\|\mathbf{y}\|_{\mathcal{E}^N}$, making them more convenient spaces to analyze such a system. In what follows, we consider $\boldsymbol{\lambda} := (\lambda^1, \dots, \lambda^N) \in (\mathcal{P}(U))^N$, and $\mathbf{X} := (X^1, \dots, X^N) \in (\mathbb{R}^d)^N$. For each $\Psi \in \mathcal{P}_1(\mathcal{Y})$, we consider the map $\mathbf{v}_\Psi^N : \hat{\mathcal{Y}}^N \rightarrow (\mathbb{R}^d)^N$ which is defined through

$$\mathbf{v}_\Psi^N(\mathbf{X}, \boldsymbol{\lambda}) := (v_\Psi(X^1, \lambda^1), \dots, v_\Psi(X^N, \lambda^N)).$$

Furthermore, we define the map $\mathcal{J}_\Psi^N : \widehat{\mathcal{Y}}^N \rightarrow (\mathcal{F}(U))^N$ as

$$\mathcal{J}_\Psi^N(\mathbf{X}, \boldsymbol{\lambda}) := (\mathcal{J}_\Psi(X^1, \lambda^1), \dots, \mathcal{J}_\Psi(X^N, \lambda^N)),$$

and we set $\widehat{\mathbf{B}}(t) := (B^1(t), \dots, B^N(t))$ as the $d \times N$ -valued Brownian motion. Then we write (2.9) in the compact form

$$\begin{cases} \mathbf{X}(t) = \mathbf{X}_0 + \int_0^t \mathbf{v}_{\Lambda_s}^N(\mathbf{X}(s), \boldsymbol{\lambda}(s)) ds + \sqrt{2\sigma} \mathbf{B}(t), \\ \boldsymbol{\lambda}(t) = \boldsymbol{\lambda}_0 + \int_0^t \mathcal{J}_{\Lambda_s}^N(\mathbf{X}(s), \boldsymbol{\lambda}(s)) ds. \end{cases} \quad (2.11)$$

In the next, we suppose that $\mathbf{X}_0 \in L^2(\Omega; (\mathbb{R}^d)^N)$, and $\lambda_0 \in (\mathcal{P}_1(U))^N$ for all $N \in \mathbb{N}$. Since the above is an SDE, we now recall what a strong solution is for (2.11).

Definition 2.9. Suppose that $(\Omega, \mathcal{B}, \mathbb{P})$ is a complete probability space endowed with the σ -algebra generated by $(\mathbf{B}(t))_{t \in [0, T]}$ and that we denote as $(\mathcal{F}_t)_{t \in [0, T]}$. We say that an \mathcal{E}^N -valued predictable process $\mathbf{Y}(t) = (\mathbf{X}(t), \boldsymbol{\lambda}(t))$, $t \in [0, T]$ is a strong solution of (2.11) if $\mathbf{Y}(t)$ satisfies \mathbb{P} -a.s. (2.11), and it has a continuous version.

Let (X, d_X) be a metric space. We denote by $L^2(X) := L^2(\Omega \times [0, T]; X)$, and we endow it with the norm

$$\|Y\|_{L^2(X)} := \mathbb{E} \left(\int_0^T \|Y(s)\|_X^2 ds \right), \quad Y \in L^2(X). \quad (2.12)$$

Further, we set $\mathcal{M}_{\text{ad}}^2(X) := L_{\text{ad}}^2(\Omega \times [0, T]; X)$ as the space of adapted processes with respect to the σ -algebra $(\mathcal{F}_t)_{t \in [0, T]}$ with values in X , and such that the norm (2.12) is finite.

Remark 2.10. In this work, we discuss the well-posedness of system (2.11) for every choice of initial condition $(\mathbf{X}_0, \boldsymbol{\lambda}_0) \in \widehat{\mathcal{Y}}^N$. Furthermore, we are interested to looking forward the existence and uniqueness of solutions $\mathbf{Y} \in \mathcal{M}_{\text{ad}}^2(\widehat{\mathcal{Y}}^N)$.

3. MAIN RESULTS

In this section, we collect our main results. In the next Theorem, we study the well-posedness of (2.9).

Theorem 3.1. *Let us fix a filtered probability space $(\Omega, \mathcal{B}, \mathbb{P})$ endowed with a complete filtration $(\mathcal{F}_t)_{t \in [0, T]}$ generated by the Brownian motion $(\mathbf{B}(t))_{t \in [0, T]}$. Assume that for every $y \in \mathcal{Y}$ and $\Psi \in \mathcal{P}_1(\mathcal{Y})$ the velocity field $v_\Psi : \mathcal{Y} \rightarrow \mathbb{R}^d$ satisfies (A1)–(A3), and the operator \mathcal{J}_Ψ satisfies (B1)–(B4). Then, for every choice $\overline{\mathbf{Y}}_0 := (\overline{\mathbf{X}}_0, \overline{\boldsymbol{\lambda}}_0)$ as initial condition of (2.11) such that $\overline{\mathbf{Y}}_0 \in L^2(\widehat{\mathcal{Y}}^N)$, the system (2.11) has a unique solution \mathbf{Y} that belongs to $\mathcal{M}_{\text{ad}}^2(\widehat{\mathcal{Y}}^N)$.*

In the next theorem, we prove that Eulerian solutions are also Lagrangian solutions.

Theorem 3.2. *Let $r > 0$ such that condition (C1) holds true. Furthermore, let us suppose that $\Lambda \in C^0([0, T]; (\mathcal{P}_1(\mathcal{Y}), \mathcal{W}_1))$ and let $\overline{\Lambda}$ be any given initial condition for (2.4) with $\overline{\Lambda} \in \mathcal{P}(B_r^{\mathcal{Y}})$. Moreover, suppose that Λ is an Eulerian solution to (2.4). Then $\Lambda \in C^0([0, T]; (\mathcal{P}(B_r^{\mathcal{Y}}), \mathcal{W}_1))$, and it is a Lagrangian solution for (2.4).*

Our next result states the existence of a Eulerian solution for equation in (2.4), and it can be related to the mean-field limit of system (2.9).

Theorem 3.3. *Let $r > 0$ such that condition (C1) holds true, and let $\bar{\Lambda} \in \mathcal{P}(B_r^y)$ be a fixed initial condition. The following hold true:*

- (i1) *there exists a unique Eulerian solution $\{\Lambda_t : t \in [0, T]\}$ in the sense of Definition 2.4 to (2.4) starting from $\bar{\Lambda}$.*
(i2) *Suppose that $\bar{\Lambda}^N := \frac{1}{N} \sum_{i=1}^N \delta_{\bar{y}^{N,i}}$ is a sequence belonging to $\mathcal{P}(B_r^y)$ such that*

$$\lim_{N \rightarrow +\infty} \mathcal{W}_1(\bar{\Lambda}^N, \bar{\Lambda}) = 0.$$

Then for all $t \in [0, T]$

$$\lim_{N \rightarrow +\infty} \mathcal{W}_1(\Lambda_t^N, \Lambda_t) = 0 \quad \mathbb{P} - a.s.,$$

and where Λ_t^N are the empirical measures given by (2.10) related to the unique solution of the system given by (2.11) with initial datum $\{\bar{y}^{N,i}\}$.

We start by proving Theorem 3.1. Let us remark that we cannot guarantee *a priori* that Λ_t for $t > 0$ is composed of independent and identically distributed random variables due to assumption (C1). Thus, approximating the system (2.11) using the propagation of chaos, as proposed by Sznitman (see [38]), might not be possible. Indeed, one needs a system of couples $(\bar{X}^1(t), \bar{\lambda}^1(t)), \dots, (\bar{X}^N(t), \bar{\lambda}^N(t))$ of independent and identically distributed random variables sufficiently close to (2.11) in the sense of [39], Definition 4.1. This, in fact, requires further knowledge on the structure of \mathbf{v} , and \mathcal{J} satisfying (A1)–(A3), and (B1)–(B4), respectively. For a self-contained review of the propagation of chaos approach, see also [40]. Hence, our proof relies on Corollary A.2, which helps us overcome the local properties of \mathbf{v} , and \mathcal{J} . Nevertheless, we need to construct probability measures $\lambda_n(t)$, $n \in \mathbb{N}$, which are necessary to obtain the solution of (2.11) by means of the Picard iteration method. This convergence issue, in the case of an equation like (2.11) with additional infinite-dimensional white noise in the second equation, requires a deeper analysis, which we postpone to future work.

Proof of Theorem 3.1. We prove this result by making use of a Picard iteration, and Corollary A.2. For each $n \in \mathbb{N}$, let us consider for any $\theta > 0$ but fixed, we set $g_{\Lambda_t}(\mathbf{X}_n(t), \lambda_n(t)) := \lambda_n(t) + \theta \mathcal{J}_{\Lambda_t}^N(\mathbf{X}_n(t), \lambda_n(t))$ where

$$\begin{cases} \mathbf{X}_{n+1}(t) = \mathbf{X}_0 + \int_0^t \mathbf{v}_{\Lambda_s}^N(\mathbf{X}_n(s), \lambda_n(s)) ds + \sqrt{2\sigma} \mathbf{B}(t), \\ \lambda_{n+1}(t) = e^{-\frac{t}{\theta}} \lambda_0 + \frac{1}{\theta} \int_0^t e^{\frac{s-t}{\theta}} g_{\Lambda_s}(\mathbf{X}_n(s), \lambda_n(s)) ds, \\ \mathbf{X}_0 := \bar{\mathbf{X}}_0, \\ \lambda_0 := \bar{\lambda}_0. \end{cases} \quad (3.1)$$

Notice that for each $n \in \mathbb{N}$, $\lambda_n \in (\mathcal{P}(U))^N$. Indeed, since

$$g_{\Lambda_t}(\mathbf{X}_n(t), \lambda_n(t)) \in (\mathcal{P}(U))^N, \quad \forall t \in [0, T],$$

then by the convexity of $(\mathcal{P}(U))^N$, it holds that for all $t \in [0, T]$

$$\frac{\frac{1}{\theta} \int_0^t e^{\frac{s-t}{\theta}} g_{\Lambda_s}(\mathbf{X}_n(s), \lambda_n(s)) ds}{\frac{1}{\theta} \int_0^t e^{\frac{s-t}{\theta}} ds} \in (\mathcal{P}(U))^N.$$

Since $\frac{1}{\theta} \int_0^t e^{\frac{s-t}{\theta}} ds = 1 - e^{-\frac{t}{\theta}}$, we get that

$$\frac{1}{\theta} \int_0^t e^{\frac{s-t}{\theta}} g_{\Lambda_s}(\mathbf{X}_n(s), \boldsymbol{\lambda}_n(s)) ds \in (1 - e^{-\frac{t}{\theta}})(\mathcal{P}(U))^N,$$

and then

$$\boldsymbol{\lambda}_{n+1}(t) = e^{-\frac{t}{\theta}} \Lambda_0 + \frac{1}{\theta} \int_0^t e^{\frac{s-t}{\theta}} g_{\Lambda_s}(\mathbf{X}_n(s), \boldsymbol{\lambda}_n(s)) ds \in (\mathcal{P}(U))^N.$$

We now use Corollary A.2 to show that (3.1) admits a solution. That is, we show that the differential equation

$$\begin{cases} d\mathbf{X}_{n+1}(t) = \mathbf{v}_{\Lambda_t}^N(\mathbf{X}_n(t), \boldsymbol{\lambda}_n(t)) + \sqrt{2\sigma} d\mathbf{B}(t), \\ d\boldsymbol{\lambda}_{n+1}(t) = -\frac{1}{\theta} \boldsymbol{\lambda}_{n+1}(t) + \frac{1}{\theta} g_{\Lambda_t}^N(\mathbf{X}_n(t), \boldsymbol{\lambda}_n(t)), \\ \mathbf{X}_0 := \bar{\mathbf{X}}_0, \\ \boldsymbol{\lambda}_0 := \bar{\boldsymbol{\lambda}}_0. \end{cases} \quad (3.2)$$

admits a unique strong solution. Indeed, we can write such a system as

$$d\mathbf{Y}(t) = \widehat{b}_{\Lambda_t}^n(\mathbf{Y}(t)), \quad \mathbf{Y} = (\mathbf{Y}_1, \mathbf{Y}_2)$$

where

$$\widehat{b}_{\Lambda_t}^n(\mathbf{Y}(t)) := \begin{pmatrix} \mathbf{v}_{\Lambda_t}^N(\mathbf{X}_n(t), \boldsymbol{\lambda}_n(t)) + \sqrt{2\sigma} d\mathbf{B}(t) \\ -\frac{1}{\theta} \mathbf{Y}_2(t) + \frac{1}{\theta} g_{\Lambda_t}(\mathbf{X}_n(t), \boldsymbol{\lambda}_n(t)) \end{pmatrix}.$$

Notice that $\widehat{b}_{\Lambda_t}^n(c)$ is time-continuous for all $c \in \widehat{\mathcal{Y}}^N$. On the other hand, for every $R > 0$, we need to verify that there exists $\theta_1 > 0$ such that for $c \in \widehat{\mathcal{Y}}^N$, $\|c\|_{\widehat{\mathcal{Y}}^N} \leq R$ then $c + \theta_1 \widehat{b}_{\Lambda_t}^n(c(t)) \in \widehat{\mathcal{Y}}^N$. Indeed, we may choose $\theta_1 = \theta$, and thus

$$c_2 + \theta_1 \left(-\frac{1}{\theta} c_2 + \frac{1}{\theta} g_{\Lambda_t}(\mathbf{X}_n(t), \boldsymbol{\lambda}_n(t)) \right) = g_{\Lambda_t}(\mathbf{X}_n(t), \boldsymbol{\lambda}_n(t)) \in (\mathcal{P}(U))^N.$$

Let us now notice that condition (i') and (ii') in Corollary A.2 are satisfied. Indeed, by definition of \widehat{b} we have that

$$\left\| \widehat{b}_{\Lambda_t}^n(\mathbf{Y}_1) - \widehat{b}_{\Lambda_t}^n(\mathbf{Y}_2) \right\|_{\widehat{\mathcal{Y}}^N} \leq \frac{1}{\theta} \|\mathbf{Y}_1 - \mathbf{Y}_2\|_{\widehat{\mathcal{Y}}^N}.$$

In the same way, we can verify that

$$\left\| \widehat{b}_{\Lambda_t}^n(\mathbf{Y}_1) \right\|_{\widehat{\mathcal{Y}}^N} \leq \frac{1}{\theta} (1 + \|\mathbf{Y}_1\|_{\widehat{\mathcal{Y}}^N}).$$

Therefore, for each fixed $n \in \mathbb{N}$, we have that there exists a unique curve $t \mapsto (\mathbf{X}_{n+1}(t), \boldsymbol{\lambda}_{n+1}(t))$ solving the differential equation (3.2). We now show that it is possible to construct the solution of (2.11) as the limit of $(\mathbf{X}_{n+1}(t), \boldsymbol{\lambda}_{n+1}(t))$, as $n \rightarrow +\infty$. First, let us now prove that $(\mathbf{X}_1, \boldsymbol{\lambda}_1) =: \mathbf{Y}_1 \in \mathcal{M}_{\text{ad}}^2(\widehat{\mathcal{Y}}^N)$, and it has a

continuous version. Let $t \in [0, T]$, and $s, t \in [0, T]$ with $s < t$. By assumption (A3), one gets

$$\begin{aligned}
|\mathbf{X}_1(t) - \mathbf{X}_1(s)| &\leq \int_s^t |\mathbf{v}_{\Lambda_r}^N(\mathbf{X}_0(r), \boldsymbol{\lambda}_0(r))| dr + \sqrt{2\sigma} |\mathbf{B}(t) - \mathbf{B}(s)| \\
&= \frac{1}{N} \sum_{i=1}^N \int_s^t |v_{\Lambda_r}(\mathbf{X}_0^i(r), \boldsymbol{\lambda}_0^i(r))| dr + \sqrt{2\sigma} |\mathbf{B}(t) - \mathbf{B}(s)| \\
&\leq M_v \int_s^t (1 + \|\bar{Y}_0\|_{\widehat{\mathcal{E}}^N} + m_1(\Lambda_r^N)) dr + \sqrt{2\sigma} |\mathbf{B}(t) - \mathbf{B}(s)| \\
&\leq M_v (2 + \|\bar{Y}_0\|_{\widehat{\mathcal{E}}^N}) |t - s| + \sqrt{2\sigma} |\mathbf{B}(t) - \mathbf{B}(s)|.
\end{aligned}$$

Hence, by taking the square and expectation with respect to \mathbb{P} , we get

$$\mathbb{E} |\mathbf{X}_1(t) - \mathbf{X}_1(s)|^2 \leq 2(M_v(2 + \|\bar{Y}_0\|_{\widehat{\mathcal{E}}^N}))^2 |t - s|^2 + 4\sigma |t - s|.$$

Therefore, By the Kolmogorov continuity theorem, $(\mathbf{X}_1(t))_{t \in [0, T]}$ has a continuous version. On the other hand, notice that

$$\begin{aligned}
|\mathbf{X}_1(t)| &\leq |\bar{\mathbf{X}}_0| + \int_0^t |\mathbf{v}_{\Lambda_s}^N(\mathbf{X}_0(s), \boldsymbol{\lambda}_0(s))| ds + \sqrt{2\sigma} |\mathbf{B}(t)| \\
&= |\bar{\mathbf{X}}_0| + \frac{1}{N} \sum_{i=1}^N \int_0^t |v_{\Lambda_s}(\mathbf{X}^i(s), \boldsymbol{\lambda}^i(s))| ds + \sqrt{2\sigma} |\mathbf{B}(t)| \\
&\leq |\bar{\mathbf{X}}_0| + M_v \int_0^t (1 + \|\bar{Y}_0\|_{\widehat{\mathcal{E}}^N} + m_1(\Lambda_s^N)) ds + \sqrt{2\sigma} |\mathbf{B}(t)|.
\end{aligned}$$

From here, we conclude that there exists a further positive constant still denoted as C such that

$$\mathbb{E} \int_0^T |\mathbf{X}_1(t)|^2 dt \leq CT.$$

Let us now proceed with $\boldsymbol{\lambda}_1(t)$. Notice that for all $s, t \in [0, T]$ with $s \leq t$

$$\begin{aligned}
\|\boldsymbol{\lambda}_1(t) - \boldsymbol{\lambda}_1(s)\|_{(\mathcal{F}(U))^N} &\leq |e^{-\frac{t}{\theta}} - e^{-\frac{s}{\theta}}| \|\boldsymbol{\lambda}_0\|_{(\mathcal{F}(U))^N} \\
&\quad + \widetilde{M}_{\mathcal{T}} \int_s^t e^{-\frac{r-t}{\theta}} (1 + |\mathbf{X}_0(r)| + m_1(\Lambda_r^N)) dr,
\end{aligned}$$

where $\widetilde{M}_{\mathcal{T}}$ is a positive constant depending on $M_{\mathcal{T}}$, and θ . By taking the square and expectation with respect to \mathbb{P} , one gets

$$\begin{aligned}
\mathbb{E} \|\boldsymbol{\lambda}_1(t) - \boldsymbol{\lambda}_1(s)\|_{(\mathcal{F}(U))^N}^2 &\leq 2|e^{-\frac{t}{\theta}} - e^{-\frac{s}{\theta}}|^2 \|\boldsymbol{\lambda}_0\|_{(\mathcal{F}(U))^N}^2 \\
&\quad + 2\widetilde{M}_{\mathcal{T}}^2 \mathbb{E} \left(\int_s^t (2 + |\mathbf{X}_0(r)|) dr \right)^2 \\
&\leq 2|e^{-\frac{t}{\theta}} - e^{-\frac{s}{\theta}}|^2 \|\boldsymbol{\lambda}_0\|_{(\mathcal{F}(U))^N}^2 \\
&\quad + 4\widetilde{M}_{\mathcal{T}}^2 \left(4|t - s|^2 + |t - s| \mathbb{E} \int_s^t |\mathbf{X}_0(r)|^2 dr \right),
\end{aligned}$$

and thus by the Kolmogorov continuity theorem, $(\boldsymbol{\lambda}_1(t))_{t \in [0, T]}$ has a continuous version. Furthermore, we can find a positive constant $C := C(\theta, M_{\mathcal{T}}, \lambda_0, \mathbf{X}_0, T)$ such that

$$\mathbb{E} \left\| \int_0^T \boldsymbol{\lambda}_1^2(t) dt \right\|_{(\mathcal{F}(U))^N} \leq CT.$$

Therefore $(\mathbf{Y}_1(t))_{t \in [0, T]} \in \mathcal{M}_{\text{ad}}^2(\widehat{\mathcal{Y}}^N)$. Let us now proceed by induction. That is, we suppose that $(\mathbf{X}_n, \boldsymbol{\lambda}_n) =: \mathbf{Y}_n \in \mathcal{M}_{\text{ad}}^2(\widehat{\mathcal{Y}}^N)$, and we prove that $\mathbf{Y}_{n+1} \in \mathcal{M}_{\text{ad}}^2(\widehat{\mathcal{Y}}^N)$. As before, let us take $s, t \in [0, T]$ and $s < t$. Then

$$\begin{aligned} |\mathbf{X}_{n+1}(t) - \mathbf{X}_{n+1}(s)| &\leq \int_s^t |v_{\Lambda_s^N}(\mathbf{X}_n(s), \boldsymbol{\lambda}_n(s))| ds + \sqrt{2\sigma} |\mathbf{B}(t) - \mathbf{B}(s)| \\ &= \frac{1}{N} \sum_{i=1}^N \int_s^t |v_{\Lambda_s}(\mathbf{X}_n^i(s), \boldsymbol{\lambda}_n^i(s))| ds + \sqrt{2\sigma} |\mathbf{B}(t) - \mathbf{B}(s)| \\ &\leq M_v \int_s^t (1 + \|\mathbf{Y}_n\|_{\widehat{\mathcal{E}}^N} + m_1(\Lambda_s^N)) ds + \sqrt{2\sigma} |\mathbf{B}(t) - \mathbf{B}(s)|. \end{aligned}$$

Hence, by taking the square and Jensen inequality, we get

$$\begin{aligned} |\mathbf{X}_{n+1}(t) - \mathbf{X}_{n+1}(s)|^2 &\leq 2M_v^2 |t - s| \int_s^t (1 + \|\mathbf{Y}_n\|_{\widehat{\mathcal{E}}^N} + m_1(\Lambda_s^N))^2 ds + 4\sigma |\mathbf{B}(t) - \mathbf{B}(s)|^2 \\ &\leq 4M_v^2 |t - s| \int_s^t (4 + \|\mathbf{Y}_n\|_{\widehat{\mathcal{E}}^N}^2) ds + 4\sigma |\mathbf{B}(t) - \mathbf{B}(s)|^2, \end{aligned}$$

and thus from this inequality and the inductive hypothesis, and the Kolmogorov continuity theorem, $(\mathbf{X}_{n+1}(t))_{t \in [0, T]}$ has a continuous version. On the other hand, notice that

$$\mathbb{E} \int_0^T |\mathbf{X}_{n+1}(t)|^2 dt \leq CT,$$

for some positive constant C . Further, the same conclusion can be obtained for λ_{n+1} . Therefore $\mathbf{Y}_n \in \mathcal{M}_{\text{ad}}^2(\widehat{\mathcal{Y}}^N)$ for all $n \in \mathbb{N}$. To conclude, We now need to prove that $(\mathbf{Y}_n)_n$ is a Cauchy sequence. Let us prove that there exists a positive constant $M > 0$ such that

$$\mathbb{E} \left(\sup_{s \in [0, t]} \|\mathbf{Y}_{n+1}(s) - \mathbf{Y}_n(s)\|_{\widehat{\mathcal{Y}}^N}^2 \right) \leq \frac{(Mt)^{n+1}}{(n+1)!} \quad (3.3)$$

for all $n \in \mathbb{N}$, and all $t \in [0, T]$. By assumption (A1), let us take R large enough such that $\Lambda_t^N \in \mathcal{P}(B_R^{\mathcal{Y}^N})$. We proceed by induction. Let us underline that the involved constants may vary from line to line. Notice that by assumption (A3), we can find a positive constant C depending on $\|\mathbf{Y}_0\|_{\widehat{\mathcal{E}}^N}$ such that

$$|\mathbf{X}_1(t) - \mathbf{X}_0(t)| \leq Ct + \sqrt{2\sigma} |\mathbf{B}(t)|,$$

and thus by Doob's inequality one gets

$$\mathbb{E} \sup_{s \in [0, t]} |\mathbf{X}_1(s) - \mathbf{X}_0(s)|^2 \leq Ct^2,$$

for some positive constant C . On the other hand, by assumption (B2) we have that

$$\begin{aligned} |\boldsymbol{\lambda}_1(t) - \boldsymbol{\lambda}_0| &\leq |\boldsymbol{\lambda}_0|(1 - \exp(-\frac{t}{\theta})) + \widetilde{M}_{\mathcal{T}} \int_0^t \exp\left(\frac{s-t}{\theta}\right) (2 + |\mathbf{X}_0|) ds \\ &\leq \frac{t}{\theta} |\boldsymbol{\lambda}_0| + \widetilde{M}_{\mathcal{T}} (2 + |\mathbf{X}_0|) t, \end{aligned}$$

and thus

$$\mathbb{E} \sup_{s \in [0, t]} |\boldsymbol{\lambda}_1(s) - \boldsymbol{\lambda}_0|^2 \leq Ct^2$$

where C is a positive constant depending on $M_{\mathcal{T}}, \sigma, \theta, T, \mathbb{E} \|\mathbf{Y}_0\|^2$. Therefore

$$\mathbb{E} \sup_{s \in [0, t]} \|\mathbf{Y}_1(s) - \mathbf{Y}_0(s)\|_{\widehat{\mathfrak{y}}_N}^2 \leq Ct^2$$

where C is a positive constant depending on $M_{\mathcal{T}}, \sigma, \theta, T, \mathbb{E} \boldsymbol{\lambda}_0^2$. We now suppose that (3.3) holds true for $n-1 \in \mathbb{N}$, and then we prove it for $n \in \mathbb{N}$. Since assumptions (A1)–(A3), and (B1)–(B4) hold true, By Proposition 2.3 item (i), there exists a positive constant L_R such that

$$\begin{aligned} |\mathbf{X}_{n+1}(t) - \mathbf{X}_n(t)| &\leq L_R \int_0^t \|\mathbf{Y}_n(s) - \mathbf{Y}_{n-1}(s)\|_{\widehat{\mathfrak{y}}_N} ds, \\ |\boldsymbol{\lambda}_{n+1}(t) - \boldsymbol{\lambda}_n(t)| &\leq L_R \int_0^t \|\mathbf{Y}_n(s) - \mathbf{Y}_{n-1}(s)\|_{\widehat{\mathfrak{y}}_N} ds, \end{aligned}$$

and thus

$$\|\mathbf{Y}_{n+1}(t) - \mathbf{Y}_n(t)\|_{\widehat{\mathfrak{y}}_N} \leq L_R \int_0^t \|\mathbf{Y}_n(s) - \mathbf{Y}_{n-1}(s)\|_{\widehat{\mathfrak{y}}_N} ds.$$

From this inequality, we get that for each $s \in [0, t]$

$$\|\mathbf{Y}_{n+1}(s) - \mathbf{Y}_n(s)\|_{\widehat{\mathfrak{y}}_N} \leq L_R \int_0^s \|\mathbf{Y}_n(r) - \mathbf{Y}_{n-1}(r)\|_{\widehat{\mathfrak{y}}_N} dr.$$

Hence, by Jensen inequality one gets

$$\|\mathbf{Y}_{n+1}(s) - \mathbf{Y}_n(s)\|_{\widehat{\mathfrak{y}}_N}^2 \leq sL_R^2 \int_0^s \|\mathbf{Y}_n(r) - \mathbf{Y}_{n-1}(r)\|_{\widehat{\mathfrak{y}}_N}^2 dr.$$

and by taking the expected value, we obtain that for each $s \in [0, t]$

$$\begin{aligned} \mathbb{E} \|\mathbf{Y}_{n+1}(s) - \mathbf{Y}_n(s)\|_{\widehat{\mathfrak{y}}_N}^2 &\leq tL_R^2 \mathbb{E} \int_0^t \|\mathbf{Y}_n(s) - \mathbf{Y}_{n-1}(s)\|_{\widehat{\mathfrak{y}}_N}^2 ds \\ &\leq (tL_R)^2 \int_0^t \frac{(Ms)^n}{n!} ds \end{aligned}$$

where in the last inequality we have used the inductive hypothesis. Therefore, we conclude that

$$\mathbb{E} \sup_{s \in [0, t]} \|\mathbf{Y}_{n+1}(s) - \mathbf{Y}_n(s)\|_{\widehat{\mathfrak{y}}_N}^2 \leq (TL_R)^2 \frac{(Mt)^{n+1}}{(n+1)!}$$

By redefining the constant M , we obtain the desired inequality (3.3). A direct consequence of (3.3) is that by Markov inequality

$$\begin{aligned} \mathbb{P}\left(\sup_{t \in [0, T]} \|\mathbf{Y}_{n+1}(t) - \mathbf{Y}_n(t)\|_{\hat{\mathcal{Y}}^N}^2 > 2^{-n}\right) &\leq 4^n \mathbb{E}\left(\sup_{t \in [0, T]} \|\mathbf{Y}_n(t) - \mathbf{Y}_{n-1}(t)\|_{\hat{\mathcal{Y}}^N}^2\right) \\ &\leq 4^n \frac{(MT)^{n+1}}{(n+1)!} \end{aligned}$$

Since the series $\sum_{n \in \mathbb{N}} 4^n \frac{(MT)^{n+1}}{(n+1)!}$ converges, then by Borel–Cantelli

$$\mathbb{P}\left(\limsup_{n \in \mathbb{N}} \{\omega \in \Omega : \sup_{t \in [0, T]} \|\mathbf{Y}_{n+1}(t) - \mathbf{Y}_n(t)\|_{\hat{\mathcal{Y}}^N}^2 > 2^{-n}\}\right) = 0.$$

Then, we may define $\mathbf{Y} := \lim_{n \rightarrow +\infty} \mathbf{Y}_n \in \mathcal{M}_{\text{ad}}^2(\hat{\mathcal{Y}}^N)$. Notice that this process solves

$$\begin{cases} \mathbf{X}(t) = \mathbf{X}_0 + \int_0^t \mathbf{v}_{\Lambda_s}(\mathbf{X}(s), \boldsymbol{\lambda}(s)) ds + \sqrt{2\sigma} \mathbf{B}(t), \\ \boldsymbol{\lambda}(t) = e^{-\frac{t}{\theta}} \boldsymbol{\lambda}_0 + \frac{1}{\theta} \int_0^t e^{\frac{s-t}{\theta}} g_{\Lambda_s}(\mathbf{X}(s), \boldsymbol{\lambda}(s)) ds, \\ \mathbf{X}_0 := \bar{\mathbf{X}}_0, \\ \boldsymbol{\lambda}_0 := \bar{\boldsymbol{\lambda}}_0. \end{cases}$$

It is true that, since $\hat{\mathcal{Y}}^N$ is a closed convex subset of $\hat{\mathcal{E}}^N$, then by Corollary A.2, we have that $\boldsymbol{\lambda}(t) \in (\mathcal{P}(U))^N$. However, this implication is not entirely straightforward, and therefore we provide further details in the following. Since $\{\boldsymbol{\lambda}_n(t)\}_{n \in \mathbb{N}}$ is a Cauchy sequence with respect to the BL-norm $\|\boldsymbol{\lambda}_n(t)\|_{\text{BL}} = \frac{1}{N} \sum_{i=1}^N \|\lambda_n^i\|_{\text{BL}}$, by applying (2.1), for any $\varepsilon > 0$ there exists $N \in \mathbb{N}$ such that

$$\mathcal{W}_1(\boldsymbol{\lambda}_n(t), \boldsymbol{\lambda}_m(t)) \leq (1 + D_U) \|\boldsymbol{\lambda}_n(t) - \boldsymbol{\lambda}_m(t)\|_{\text{BL}} \leq (1 + D_U) \varepsilon$$

for any $n, m \geq N$ positive integers, and all $t \in [0, T]$. On the other hand, for some $\bar{x} \in U$,

$$\begin{aligned} m_1(\boldsymbol{\lambda}_n(t)) &= \frac{1}{N} \sum_{i=1}^N \int_U d_U(x, \bar{x}) \lambda_n^i(dx) \leq \sup_{x \in U} d_U(x, \bar{x}) \\ &\leq \text{diam}(U) < +\infty, \end{aligned}$$

where $\text{diam}(U)$ is the diameter of U . Then, we conclude that $\{\boldsymbol{\lambda}_n(t)\}_{n \in \mathbb{N}}$ is a Cauchy sequence with respect to the Wasserstein-1 distance. Applying [34], Proposition 2.2.8, $((\mathcal{P}_1(U))^N, \mathcal{W}_1)$ is a complete metric space, then $\boldsymbol{\lambda}(t) \in (\mathcal{P}(U))^N$.

We now need to come back to the original problem. Indeed, we now aim to show that \mathbf{Y} solves (2.11). The only thing that we need to prove is that

$$\boldsymbol{\lambda}(t) = e^{-\frac{t}{\theta}} \boldsymbol{\lambda}_0 + \frac{1}{\theta} \int_0^t e^{\frac{s-t}{\theta}} g_{\Lambda_s}(\mathbf{X}(s), \boldsymbol{\lambda}(s)) ds, \quad \bar{\boldsymbol{\lambda}}(t) = \boldsymbol{\lambda}_0 + \int_0^t \mathcal{J}_{\Lambda_s}^N(\mathbf{X}(s), \boldsymbol{\lambda}(s)) ds,$$

coincide. Let us define for each $t \in [0, T]$ the function $\phi(t) := \boldsymbol{\lambda}(t) - \bar{\boldsymbol{\lambda}}(t)$. Notice that

$$\frac{d}{dt} \boldsymbol{\lambda}(t) = -\frac{1}{\theta} \boldsymbol{\lambda}(t) + \frac{1}{\theta} (\boldsymbol{\lambda}(t) + \theta \mathcal{J}_{\Lambda_t}(\mathbf{X}(t), \boldsymbol{\lambda}(t))) = \mathcal{J}_{\Lambda_t}(\mathbf{X}(t), \boldsymbol{\lambda}(t)),$$

and thus by noting that $\frac{d}{dt}\bar{\lambda}(t) = \mathcal{J}_{\Lambda_t}(\mathbf{X}(t), \boldsymbol{\lambda}(t))$ then $\frac{d}{dt}\phi(t) = 0$ for all $t \in [0, T]$. Thus $\phi(t) = c$ for all $t \in [0, T]$. So that, by the initial condition, we conclude that $c = 0$, and we are done. \square

3.1. Mean-field limit

We now analyze the asymptotic behavior of (2.11) as $N \rightarrow +\infty$. Specifically, we want to demonstrate the existence of an Eulerian solution as $N \rightarrow +\infty$ as stated in Theorem 3.3. To this aim, we first derive the Theorem 3.2, and some preliminary results.

Proof of Theorem 3.2. In order to prove this result, let us notice that $\mathcal{L}_{\bar{\Lambda}}$ defines a strongly continuous semigroup. Indeed, by condition (C1) one has that

$$C_0(\mathcal{Y}) = \overline{\text{range}(\eta I - \mathcal{L}_{\bar{\Lambda}})} = \text{range}(\eta I - \bar{\mathcal{L}}_{\bar{\Lambda}}).$$

Then by the Lummer–Phillips theorem, we can conclude that $\bar{\mathcal{L}}_{\bar{\Lambda}}$ generates a strongly continuous semigroup of contractions on $C_0(\mathcal{Y})$ that we denoted as $\{S_t\}_{t \geq 0}$. By Theorem B.2, $S_t^* \bar{\Lambda}$ is well-defined and defines a semigroup of linear and continuous operators on $(C_b(\mathcal{Y}))^*$. Moreover, since $\bar{\Lambda} \in \mathcal{P}(B_r^{\mathcal{Y}})$ and S_t is a contraction, then $S_t \bar{\Lambda} \in \mathcal{P}(B_r^{\mathcal{Y}})$. On the other hand, since (B.3) admits a unique solution, and Λ satisfies (2.4) then $\Lambda_t = S_t^* \bar{\Lambda}$. Moreover, by remark 2.5, we conclude that $S_t^* \bar{\Lambda} = (X(t), \lambda(t))_{\#} \bar{\Lambda}$, and we are done. \square

We now aim to prove Theorem 3.3. We start by showing that the sequences $\{\Lambda_t^N\}_{N \in \mathbb{N}}$, $\{X^N(t)\}_{N \in \mathbb{N}}$, and $\{\lambda^N(t)\}_N$ are tight. To this aim, we use the so-called Aldous criteria, see [41], Theorem 16.10, see also Appendix C below.

Proposition 3.4. *Let us consider the same setting as in Theorem 3.1. and suppose that for each $i = 1, \dots, N$, X_0^i is distributed according to some fixed probability measure $\bar{\mu}$. Then the following holds true.*

1. *There exists a positive constant C independent of N such that*

$$\sup_{i=1, \dots, N} \left\{ \sup_{t \in [0, T]} \left\{ \mathbb{E}|X^i(t)|^2 + \mathbb{E}\|\lambda^i(t)\|^2 \right\} \right\} < C; \quad (3.4)$$

2. *For all $\varepsilon > 0, \eta > 0$ there exists $\delta_0 > 0$ and $N_0 \in \mathbb{N}$ such that for all $N \geq N_0$, and for all discrete valued $\sigma(X^N(s) : s \in [0, T])$ -stopping times β with $0 \leq \beta \leq \beta + \delta_0 \leq T$, it is true that*

$$\sup_{0 \leq \delta \leq \delta_0} \mathbb{P}(|X^N(\beta + \delta) - X^N(\beta)| \geq \eta) \leq \varepsilon; \quad (3.5)$$

and the same inequality holds true for the sequence $\{\text{Law}(\lambda_t^N)\}_{N \in \mathbb{N}}$.

3. *For all $t \in [0, T]$, the sequence of probability distributions $\{\text{Law}(X_t^N)\}_{N \in \mathbb{N}}$ is tight as a sequence of probability measures on \mathbb{R}^d , and the same holds true for the sequence of probability distributions $\{\text{Law}(\lambda_t^N)\}_{N \in \mathbb{N}}$ as a sequence of probability measures on $\mathcal{P}(U)$.*

Remark 3.5. Let us recall that $\mathcal{M}(\mathcal{Y})$ is a locally convex Hausdorff topological space, and its topological dual is $C_b(\mathcal{Y})$. Furthermore, $\mathcal{P}(\mathcal{Y})$ is a closed subset of $\mathcal{M}(\mathcal{Y})$. This fact is crucial when considering the asymptotic behavior of Λ_t^N , the empirical measure of $(X^i(t), \lambda^i(t))_{i=1}^N$, especially in the case of identically distributed random variables. Despite the dependencies introduced by the interaction terms $v_{\Lambda_t^N}$, and $\mathcal{J}_{\Lambda_t^N}$, we will show that Λ_t^N converges to some $\bar{\Lambda}_t$ \mathbb{P} -a.s.. A strategy to establish this convergence is to prove that the sequence $(\Lambda_t^N)_{N \in \mathbb{N}}$ satisfies a large deviation principle (LDP). This is feasible because the closed convex hull of any compact set $K \subset \mathcal{P}(\mathcal{Y})$ is compact with respect to the relative topology induced by $\mathcal{M}(\mathcal{Y})$. To demonstrate this LDP, we can

apply [42], Theorem 6.1.3, which establishes that the sequence $(\Lambda_t^N)_N$ satisfies a weak large deviation principle in $\mathcal{P}(\mathcal{Y})$, and the convex rate function is given by

$$\Phi_t^*(\nu) := \sup_{\varphi \in C_b(\mathcal{Y})} \{\langle \varphi, \nu \rangle - \Phi_t(\nu)\}, \quad \nu \in \mathcal{P}(\mathcal{Y}),$$

where, for $\varphi \in C_b(\mathcal{Y})$,

$$\Phi_t(\nu) := \log \mathbb{E} \left(\exp \left\langle \varphi, \delta_{y_t^1} \right\rangle \right) = \log \mathbb{E}(\exp(\varphi(y_t^1))),$$

at least in the case of independent, and identically distributed random variables. Subsequently, by [42], Lemma 6.2.6, we have that $(\Lambda_t^N)_{N \in \mathbb{N}}$ is exponentially tight, and thus, a full large deviation principle follows for all $t \in [0, T]$. Therefore, $\Lambda_t^N \rightarrow \bar{\Lambda}_t$ \mathbb{P} -a.s., for all $t \in [0, T]$.

Notice that Remark 3.5 provides an interesting description of the convergence for the empirical measure Λ_t^N . However, it does not guarantee time regularity properties for the limit $\bar{\Lambda}_t$. Therefore, it motivates the study of (2.9) by considering the Eulerian and Lagrangian solutions.

Proof of Proposition 3.4. Let us fix $N \in \mathbb{N}$, and for each $i = 1, \dots, N$ consider the system (2.9). We start by proving that (3.4) holds true. Indeed, notice that

$$\begin{aligned} |X^i(t)| &\leq |X^i(0)| + \int_0^t |v_{\lambda_s^N}(X^i(s), \lambda^i(s))| ds + \sqrt{2\sigma} |B_t^i| \\ &\leq |X^i(0)| + M_v \int_0^t (1 + \|(X^i(s), \lambda^i(s))\| + m_1(\Lambda_s^N)) ds + \sqrt{2\sigma} |B_t^i|. \end{aligned}$$

On the other hand, we have that

$$\begin{aligned} \|\lambda^i(t)\| &\leq \|\lambda^i(0)\| + \int_0^t |\mathcal{T}_{\lambda_s^N}(X^i(s), \lambda^i(s))| ds \\ &\leq \|\lambda^i(0)\| + M_{\mathcal{T}} \int_0^t (1 + \|X^i(s)\| + m_1(\Lambda_s^N)) ds \\ &\leq \|\lambda^i(0)\| + M_{\mathcal{T}} \int_0^t (1 + \|(X^i(s), \lambda^i(s))\| + m_1(\Lambda_s^N)) ds. \end{aligned}$$

Therefore, one obtains that

$$\begin{aligned} \frac{1}{N} \sum_{i=1}^N \|(X^i(t), \lambda^i(t))\|_{\mathcal{Y}} &\leq \frac{1}{N} \sum_{i=1}^N \|(X^i(0), \lambda^i(0))\|_{\mathcal{Y}} + C_{v, \mathcal{T}} t + \sqrt{2\sigma} |B_t^i| \\ &\quad + C_{v, \mathcal{T}} \int_0^t \left(\frac{1}{N} \sum_{i=1}^N \|(X^i(s), \lambda^i(s))\|_{\mathcal{Y}} + m_1(\Lambda_s^N) \right) ds \\ &\leq \frac{1}{N} \sum_{i=1}^N \|(X^i(0), \lambda^i(0))\|_{\mathcal{Y}} + C_{v, \mathcal{T}} t + \sqrt{2\sigma} |B_t^i| \\ &\quad + C_{v, \mathcal{T}} \int_0^t \left(\frac{2}{N} \sum_{i=1}^N \|(X^i(s), \lambda^i(s))\|_{\mathcal{Y}} \right) ds, \end{aligned}$$

where $C_{v,\mathcal{T}} := \max\{M_v, M_{\mathcal{T}}\}$. By taking the expected value, we derive that

$$\begin{aligned} \frac{1}{N} \sum_{i=1}^N \mathbb{E} \|(X^i(t), \lambda^i(t))\|_{\mathbf{y}} &\leq \frac{1}{N} \sum_{i=1}^N \mathbb{E} \|(X^i(0), \lambda^i(0))\|_{\mathbf{y}} + C_{v,\mathcal{T}}t \\ &\quad + C_{v,\mathcal{T}} \int_0^t \left(\frac{1}{N} \sum_{i=1}^N \mathbb{E} \|(X^i(s), \lambda^i(s))\|_{\mathbf{y}} + \mathbb{E}m_1(\Lambda_s^N) \right) ds. \end{aligned}$$

By applying Grönwall inequality, it implies that

$$\begin{aligned} \frac{1}{N} \sum_{i=1}^N \mathbb{E} \|(X^i(t), \lambda^i(t))\|_{\mathbf{y}} &\leq \left(\frac{1}{N} \sum_{i=1}^N \mathbb{E} \|(X^i(0), \lambda^i(0))\|_{\mathbf{y}} \right) \times \exp(C_{v,\mathcal{T}}T) \\ &\quad + \left(C_{v,\mathcal{T}}t + \int_0^t \mathbb{E}m_1(\Lambda_s^N) ds \right) \times \exp(C_{v,\mathcal{T}}T), \end{aligned}$$

and thus

$$\begin{aligned} \mathbb{E}m_1(\Lambda_t^N) &\leq \frac{1}{N} \sum_{i=1}^N \mathbb{E} \|(X^i(t), \lambda^i(t))\|_{\mathbf{y}} \\ &\leq \left(\frac{1}{N} \sum_{i=1}^N \mathbb{E} \|(X^i(0), \lambda^i(0))\|_{\mathbf{y}} \right) \times \exp(C_{v,\mathcal{T}}T) \\ &\quad + \left(C_{v,\mathcal{T}}t + \int_0^t \mathbb{E}m_1(\Lambda_s^N) ds \right) \times \exp(C_{v,\mathcal{T}}T). \end{aligned}$$

Therefore, we obtain by Grönwall inequality that

$$\mathbb{E}m_1(\Lambda_t^N) \leq \exp(C_{v,\mathcal{T}}) \exp(\exp(C_{v,\mathcal{T}})t) \left(C_{v,\mathcal{T}}t + \frac{1}{N} \sum_{i=1}^N \mathbb{E} \|(X^i(0), \lambda^i(0))\|_{\mathbf{y}} \right) =: C_{v,\mathcal{T},t}, \quad (3.6)$$

for all $t \in [0, T]$. From here, we obtain by Jensen inequality, and the Itô identity that

$$\begin{aligned} \mathbb{E}|X^i(t)|^2 &\leq \mathbb{E}|X^i(0)|^2 + C_v t \int_0^t \left(1 + \mathbb{E} \|(X^i(s), \lambda^i(s))\|^2 + C_{v,\mathcal{T},T}^2 \right) ds + 2\sigma t, \\ \mathbb{E} \|\lambda^i(t)\|^2 &\leq \mathbb{E} \|\lambda^i(0)\|^2 + M_{\mathcal{T}} t \int_0^t \left(1 + \|(X^i(s), \lambda^i(s))\| + C_{v,\mathcal{T},T}^2 \right) ds. \end{aligned} \quad (3.7)$$

Therefore, by Grönwall inequality, we can find a positive constant $C := C(v, \mathcal{T}, T)$ independent of N , and depending on T, v, \mathcal{T} such that (3.5) holds true. Let us proceed with the second item. Let β be a generic stopping time, and let $\delta > 0$. We have that for each $i = 1, \dots, N$,

$$\begin{aligned} |X^i(\beta + \delta) - X^i(\beta)| &\leq \int_{\beta}^{\beta+\delta} v_{\Lambda_s^N}(X^i(s), \lambda^i(s)) ds + \sqrt{2\sigma} |B_{\delta}^i| \\ &\leq M_v \int_{\beta}^{\beta+\delta} \left(1 + \|(X^i(s), \lambda^i(s))\| + m_1(\Lambda_s^N) \right) ds + \sqrt{2\sigma} |B_{\delta}^i|. \end{aligned}$$

Then by taking the expectation in both sides of the previous inequality, we get

$$\mathbb{E}|X^i(\beta + \delta) - X^i(\beta)| \leq C\delta$$

where we have used that $\mathbb{E}\|X^i(t)\| < +\infty$, and $\mathbb{E}m_1(\Lambda_t^N) < +\infty$ for all $t \in [0, T]$, and thus C is a positive constant independent of N , and t . Hence, by letting

$$U_\varepsilon := \left\{ \omega \in \Omega : |X^i(\beta + \delta, \omega) - X^i(\beta, \omega)| \leq \frac{C}{\varepsilon} \right\},$$

we have that by Markov inequality that

$$\mathbb{P}\left(U_\varepsilon^c > \frac{C}{\varepsilon}\right) \leq \frac{\varepsilon \mathbb{E}|X^i(\beta + \delta) - X^i(\beta)|}{C} \leq \varepsilon,$$

and this completes the proof of (3.5). The same reasoning can be applied for $\{\text{Law}(\lambda_t^N)\}_{N \in \mathbb{N}}$. Let us proceed with the third item. Notice that by Markov inequality, one has that

$$\mathbb{P}\left(\sup_{t \in [0, T]} \|(X^i(t), \lambda^i(t))\| \geq a\right) \leq \frac{\mathbb{E}\left(\sup_{t \in [0, T]} \|(X^i(t), \lambda^i(t))\|\right)^2}{a^2},$$

for all $a > 0$. Then by letting $a \rightarrow +\infty$, we have by the previous estimates that

$$\lim_{a \rightarrow +\infty} \mathbb{P}\left(\sup_{t \in [0, T]} \|(X^i(t), \lambda^i(t))\| \geq a\right) = 0.$$

Therefore, we can conclude that the sequences $\{X^i(t) : t \in [0, T]\}_{i \in \mathbb{N}}$, $\{\text{Law}(\lambda_t^N)\}_{N \in \mathbb{N}}$ are tight. \square

3.2. Stability

In the next, we prove that starting from a probability measure $\bar{\Lambda} \in \mathcal{P}(B_r^{\mathcal{Y}})$ for some $r > 0$, Lagrangian solutions to (2.4) belongs to $\mathcal{P}_1(B_r^{\mathcal{Y}})$.

Lemma 3.6. *Let us take $r > 0$, and $\bar{\Lambda} \in \mathcal{P}_c(B_r^{\mathcal{Y}})$. Furthermore, let us suppose that the velocity field $v_\Psi : \mathcal{Y} \rightarrow \mathbb{R}^d$ satisfies (A1)–(A3), and the operator \mathcal{T}_Ψ satisfies (B1)–(B4). Suppose that $\Lambda \in C^0([0, T]; (\mathcal{P}_1(\mathcal{Y}), \mathcal{W}_1))$ is a Lagrangian solution for (2.4) with initial condition $\bar{\Lambda}$. Then $\Lambda_t \in \mathcal{P}_1(B_r^{\mathcal{Y}})$ for all $t \in [0, T]$.*

Proof. We want to prove that $\Lambda_t \in \mathcal{P}_1(B_r^{\mathcal{Y}})$. To do that, we prove that $m_1(\Lambda_t) \leq m_1(\bar{\Lambda})$. Indeed, since by hypothesis Λ is a Lagrangian solution, and by using (2.8) $\Lambda_t = (\text{ev}_t)_\# \bar{\Lambda}$ the following inequalities hold true. Let $X := C([0, T]; \mathcal{Y})$ so that

$$\begin{aligned} m_1(\Lambda_t) &= \int_{\mathcal{Y}} \|y\|_{\mathcal{Y}} \, d\Lambda_t(y) \\ &= \int_X \|\text{ev}_t(\varphi)\|_{\mathcal{Y}} \, d\bar{\Lambda}(\varphi) = \int_X \|\varphi(t)\|_{\mathcal{Y}} \, d\bar{\Lambda}(\varphi) \\ &\leq \int_X \sup_{t \in [0, T]} \|\varphi(t)\| \, d\bar{\Lambda}(\varphi) \\ &= \int_X \|\varphi\|_\infty \, d\bar{\Lambda}(\varphi) = m_1(\bar{\Lambda}), \end{aligned}$$

and thus $m_1(\Lambda_t) \leq m_1(\bar{\Lambda}) \leq r$. Therefore, we obtain that $\Lambda_t \in \mathcal{P}_1(B_r^{\mathcal{Y}})$ for all $t \in [0, T]$. \square

Remark 3.7. Notice that if we directly use the system (3.9), by Grönwall inequality we can prove that $\Lambda_t \in \mathcal{P}_1(B_R^y)$ where

$$m_1(\Lambda_t) \leq (r + Mt) \exp(2Mt) =: R.$$

Lemma 3.8. Suppose that Λ^1, Λ^2 are two Lagrangian solutions for (2.4) starting from $\bar{\Lambda}^1, \bar{\Lambda}^2 \in \mathcal{P}_1(B_r^y)$ for some fixed $r > 0$. Then

$$\mathcal{W}_1(\Lambda_t^1, \Lambda_t^2) \leq C \exp(tL_R) \mathcal{W}_1(\bar{\Lambda}^1, \bar{\Lambda}^2),$$

where $C := 1 + \exp(\exp(TL_R)) \exp(TL_R)$.

Proof. Let us fix $r > 0$, and consider two Lagrangian solutions Λ^1, Λ^2 for (2.4) starting from $\bar{\Lambda}^1, \bar{\Lambda}^2 \in \mathcal{P}_1(B_r^y)$. By using (3.9), we get

$$\begin{aligned} \|y^1(t) - y^2(t)\|_y &\leq \|y_0^1 - y_0^2\|_y + \int_0^t \|b_{\Lambda_s^1}(y^1(s)) - b_{\Lambda_s^2}(y^2(s))\|_y ds \\ &\leq \|y_0^1 - y_0^2\|_y + \int_0^t \|b_{\Lambda_s^1}(y^1(s)) - b_{\Lambda_s^1}(y^2(s))\|_y ds \\ &\quad + \int_0^t \|b_{\Lambda_s^1}(y^2(s)) - b_{\Lambda_s^2}(y^2(s))\|_y ds \\ &\leq \|y_0^1 - y_0^2\|_y + L_R \int_0^t \|y^1(s) - y^2(s)\|_y ds + L_R \int_0^t \mathcal{W}_1(\Lambda_s^1, \Lambda_s^2) ds. \end{aligned}$$

Let us set $\alpha(t) := \|y_0^1 - y_0^2\|_y + L_R \int_0^t \|y^1(s) - y^2(s)\|_y ds$. By Grönwall inequality, we obtain

$$\begin{aligned} \|y^1(t) - y^2(t)\|_y &\leq \alpha(t) + L_R \int_0^t \alpha(s) \exp(L_R(t-s)) ds \\ &\leq \alpha(t) \left(1 + L_R \int_0^t \exp(L_R(t-s)) ds \right) \\ &= \alpha(t) \exp(tL_R). \end{aligned}$$

From the previous inequality, we then get that

$$\mathcal{W}_1(\Lambda_t^1, \Lambda_t^2) \leq \exp(tL_R) \left(\mathcal{W}_1(\Lambda_0^1, \Lambda_0^2) + L_R \int_0^t \mathcal{W}_1(\Lambda_s^1, \Lambda_s^2) ds \right).$$

By applying Grönwall inequality, we obtain

$$\begin{aligned} \mathcal{W}_1(\Lambda_t^1, \Lambda_t^2) &\leq \exp(tL_R) \mathcal{W}_1(\Lambda_0^1, \Lambda_0^2) \\ &\quad + \mathcal{W}_1(\Lambda_0^1, \Lambda_0^2) \int_0^t L_R \exp(2L_R s) \exp\left(L_R \int_s^t \exp(L_R r) dr\right) ds \\ &\leq \exp(tL_R) \mathcal{W}_1(\Lambda_0^1, \Lambda_0^2) \\ &\quad + \exp(\exp(L_R T)) \exp(TL_R) \mathcal{W}_1(\Lambda_0^1, \Lambda_0^2) L_R \int_0^t \exp(sL_R) ds \\ &\leq \exp(tL_R) \mathcal{W}_1(\Lambda_0^1, \Lambda_0^2) \\ &\quad + \exp(\exp(L_R T)) \exp(TL_R) \mathcal{W}_1(\Lambda_0^1, \Lambda_0^2) \exp(tL_R). \end{aligned}$$

□

We let for all $i = 1, \dots, N$,

$$b_{\Psi}(y(t)) = \begin{pmatrix} v_{\Psi}(y(t)) \\ \mathfrak{T}_{\Psi}(y(t)) \end{pmatrix}, \quad W^i(t) := \begin{pmatrix} -\sqrt{2\sigma} \frac{d}{dt} B^i(t) \\ 0 \end{pmatrix},$$

and we write equation (2.9) as

$$\frac{d}{dt} y^i(t) = b_{\Lambda_t^N}(y^i(t)) - W^i(t). \quad (3.8)$$

Let us notice that by (3.6), we have that our sequence $\{\Lambda_t^N\}_{t \in [0, T]}$ belongs to some ball $B_R^{\mathfrak{y}}$ for some positive radius R independent of N . Since U is assumed compact, we have that $B_R^{\mathfrak{y}}$ is compact, and then we have that Λ_t converges weakly to some $\Lambda_t \in B_R^{\mathfrak{y}}$ as $N \rightarrow +\infty$. Then as $N \rightarrow +\infty$, we obtain the SDE

$$\begin{aligned} X(t) &= \bar{X}_0 + \int_0^t v_{\Lambda_t}(X(s), \lambda(s)) ds + \sqrt{2\sigma} \bar{B}(t), \\ \lambda(t) &= \bar{\lambda}_0 + \int_0^t \mathfrak{T}_{\Lambda_t}(X(s), \lambda(s)) ds, \end{aligned} \quad (3.9)$$

where $X(0) = \bar{X}_0 \in L^2(\Omega; \mathbb{R}^d)$, and $\lambda(0) = \bar{\lambda}_0 \in L^2(\Omega; \mathcal{P}(U))$, \bar{B} is a suitable Brownian motions. We are now ready to prove Theorem 3.3.

Proof of Theorem 3.3. Let us take $r > 0$ such that condition (C1) holds true, and let $\bar{\Lambda} \in \mathcal{P}(B_r^{\mathfrak{y}})$ be a fixed initial condition for (2.4). In what follows, we start by proving item (ii). Consider a sequence $\bar{\Lambda}^N$ of atomic measures such that

$$\lim_{N \rightarrow +\infty} \mathcal{W}_1(\bar{\Lambda}^N, \bar{\Lambda}) = 0.$$

Notice that by letting $\bar{\Lambda}^N := \frac{1}{N} \sum_{i=1}^N \delta_{y_0^i}$, where y_0^i are the initial datum for

$$\dot{y}^i(t) = \widehat{b}_{\Lambda_t^i}(y^i(t)) - W(t),$$

with

$$W(t) := \begin{pmatrix} -\sqrt{2\sigma} \dot{B}^i(t) \\ 0 \end{pmatrix},$$

by Theorem 3.2, we obtain that $\Lambda_t^N = (\bar{F}(t, 0, \cdot))_{\#} \bar{\Lambda}^N$ is a Lagrangian and Eulerian solution to (2.4) starting from $\bar{\Lambda}^N$. Then by Lemma 3.8, we have that there exists a positive constant C such that

$$\mathcal{W}_1(\Lambda_t^N, \Lambda_t^M) \leq C \exp(tL_R) \mathcal{W}_1(\bar{\Lambda}^N, \bar{\Lambda}^M) \quad (3.10)$$

for all $t \in [0, T]$, and all $N, M \in \mathbb{N}$. Moreover, by Lemma 3.6, we get that $\Lambda^N \in C([0, T]; \mathcal{P}_1(B_r^{\mathfrak{y}}, \mathcal{W}_1))$. Furthermore, since by assumption $\lim_{N \rightarrow +\infty} \mathcal{W}_1(\bar{\Lambda}^N, \bar{\Lambda}) = 0$, and by Remark 2.1 $\mathcal{P}(B_r^{\mathfrak{y}})$ is compact, by (3.10) one obtains that the sequence $\{\Lambda_t^N\}_{N \in \mathbb{N}}$ \mathcal{W}_1 -converges to some limit $\Lambda_t \in \mathcal{P}(B_r^{\mathfrak{y}})$ for each $t \in [0, T]$. We now prove that $\Lambda := \lim_{N \rightarrow +\infty} \Lambda^N$ is a Lagrangian solution. Let us now consider $\dot{y}^N(t) = \widehat{b}_{\Lambda_t^N}(y(t)) - \bar{W}(t)$, and

$\dot{y}(t) = \widehat{b}_\Lambda(y(t)) - \overline{W}(t)$ starting from some $\bar{y} \in B_r^y$, and where

$$\overline{W}(t) := \begin{pmatrix} -\sqrt{2\sigma} \frac{d}{dt} \overline{B}(t) \\ 0 \end{pmatrix},$$

for some suitable Brownian motion \overline{B} . Notice that

$$\begin{aligned} \|y^N(t) - y(t)\|_y &\leq \int_0^t \|b_{\Lambda_s^N}(y^N(s)) - b_{\Lambda_s}(y(s))\|_y ds \\ &\leq \int_0^t \|b_{\Lambda_s}(y^N(s)) - b_{\Lambda_s}(y(s))\|_y ds \\ &\quad + \int_0^t \|b_{\Lambda_s^N}(y^N(s)) - b_{\Lambda_s}(y^N(s))\|_y ds \\ &\leq L_R \int_0^t \|y^N(s) - y(s)\|_y ds + L_R \int_0^t \mathcal{W}_1(\Lambda_s^N, \Lambda_s) ds. \end{aligned}$$

Then by Lemma 3.8, there exists a positive constant C such that

$$\|y^N(t) - y(t)\|_y \leq L_R \int_0^t \|y^N(s) - y(s)\|_y ds + L_R C \int_0^t \exp(sL_R) \mathcal{W}_1(\overline{\Lambda}^N, \overline{\Lambda}) ds,$$

and thus

$$\begin{aligned} \|y^N(t) - y(t)\|_y &\leq C \exp(tL_R) \mathcal{W}_1(\overline{\Lambda}^N, \overline{\Lambda}) \\ &\quad + L_R \mathcal{W}_1(\overline{\Lambda}^N, \overline{\Lambda}) \exp(tL_R) \int_0^t \exp(L_R(t-s)) ds \\ &\leq \exp(2tL_R) (C+1) \mathcal{W}_1(\overline{\Lambda}^N, \overline{\Lambda}) \end{aligned}$$

and from here, we get by a comparison argument that

$$\|\overline{F}_{\Lambda^N}(t, 0, \bar{y}) - \overline{F}_\Lambda(t, 0, \bar{y})\| \leq \exp(2tL_R) (C+1) \mathcal{W}_1(\overline{\Lambda}^N, \overline{\Lambda}),$$

and thus

$$\Lambda_t^N = (\overline{F}_{\Lambda^N}(t, 0, \bar{y}))_{\#} \overline{\Lambda}^N \rightarrow (\overline{F}_\Lambda(t, 0, \bar{y}))_{\#} \overline{\Lambda}.$$

Hence by the uniqueness of Lagrangian solutions we get that $(\overline{F}_\Lambda(t, 0, \bar{y}))_{\#} \overline{\Lambda} = \Lambda_t$. By Theorem 3.2, we conclude that Λ is also a Eulerian solution, and we are done. \square

4. A MEAN-FIELD MODEL FOR A LARGE NETWORK OF INTERACTING NEURONS

In this part, we consider a mean-field approach to study networks of spiking neurons. The time evolution, between two consecutive spikes, of the membrane potential $X^i(t)$ of a neuron i embedded in a network, is described by a system of type (2.11). This model is of integrate-and-fire type, since the drift part accounts (*integrate*) for the inputs the neuron receives from its neighboring neurons. At each spiking time \bar{t} (*fire*), that is when $X^i(\bar{t}) = X_F$ (firing threshold), the potential is reset to X_R (resting potential) and all other neurons

connected receive an additional potential depending on λ^i . These inputs affect the membrane potential through the field v that depends both on the actual state X^i and on the coupling strength with the connected neurons. The dependence of v on the state X^i follows the principles of integrate-and-fire neuronal models with reversal potentials (see [43]). The dependence of v on λ^i , on the other hand, accounts for the different synaptic connection among neurons and could also include both inhibitory and excitatory contributions. Finally the dependence of the whole dynamics on the empirical measure Λ , which in this case represents the distribution of membrane potential values, can be used to model phenomena such as collective behavior, bursting, and avalanches, to name a few.

More precisely, we consider the system

$$\begin{cases} X^i(t) = X_0^i + \int_0^t v_{\Lambda_s^N}(X^i(s), \lambda^i(s)) ds + \sqrt{2\sigma} B_t^i, & \text{if } X^i(t^-) < X_F, \\ X^i(t) = X_R, & \text{if } X^i(t^-) = X_F, \\ \lambda^i(t) = \lambda_0^i + \int_0^t \mathcal{T}_{\Lambda_s^N}(X^i(s), \lambda^i(s)) ds. \end{cases} \quad (4.1)$$

This model includes the feature that the interactions between the neurons are also subject to random synaptic weights [20, 21]. Different scenarios can arise according to the form of \mathcal{T} .

In what follows, let us consider a function $\alpha : U \times \mathcal{Y} \rightarrow \mathbb{R}$, and define $\mathcal{T} : \mathcal{Y} \times \mathcal{P}_1(\mathcal{Y}) \rightarrow \mathcal{F}(U)$ as

$$\begin{aligned} \mathcal{T}(y, \Psi) &:= \int_{\mathcal{Y}} \alpha(\cdot, z) d\Psi(z), \quad \text{such that } \int_U \int_{\mathcal{Y}} \alpha(u, z) d\Psi(z) d\mu(u) = 0, \\ \mu(u) &= \sum_{k \geq 1} \beta_k \delta_{u_k}, \quad u_k \in U, \beta_k \in \mathbb{R}, \end{aligned} \quad (4.2)$$

where $\mu \in \mathcal{M}(U)$ defines a signed measure. Furthermore, we consider a Lipschitz velocity field $b : \mathbb{R}^d \rightarrow \mathbb{R}^d$, and we define $v_{\mathcal{T}} : \mathcal{Y} \rightarrow \mathbb{R}^d$, $(x, \lambda) \mapsto b(x)$. Since the third equation of model (4.1) serves as a recipient to model the strength of the connection between neurons, we can think of U as a discrete set that counts the number of neurons present in the brain. For the sake of simplicity, we may take $U = \mathbb{Z}^d$, endowed with a suitable metric that makes it compact.. By taking $\Psi = \Lambda_t^N$, $N \in \mathbb{N}$, we have that $\mathcal{T}(y, \Lambda_t^N)$ in (4.2) can be written as

$$\mathcal{T}(y, \Psi) = \frac{1}{N} \sum_{k \geq 1} \sum_{i=1}^N \beta_k \alpha(u_k, X^i(t), \lambda^i(t)). \quad (4.3)$$

Remark 4.1. Expression (4.3) is of great generality and contains, as special cases, the random synaptic weights seen as constants, Bernoulli or Gaussian random variables considered in previous works [20, 22–24]. To support this claim, we can compare system (4.1) and (4.3) with, for instance, the equation presented in [20]

$$X^i(t) = X_0^i + \int_0^t b(X^i(s)) ds + \sum_{j=1}^N \sum_{k \geq 1} J_N^{j \rightarrow i} 1_{\{\tau_k^j \leq t\}} + W_t \quad (4.4)$$

that combines the two equations for X and λ and where τ_k^j denotes the k -th spike time of neuron j and the synaptic weights $J_N^{j \rightarrow i}$ are defined in two representative cases:

- $J_N^{j \rightarrow i} = \alpha/N$, with α constant
- $J_N^{j \rightarrow i} = \beta \alpha^{i,j}/N$, with β constant and $\alpha^{i,j} \sim \text{Bern}(p)$.

Unlike (2.11), the dynamics of model (4.1) exhibit resets (jumps) corresponding to the spikes. Therefore, comments on the well-posedness of (4.1) are necessary and are provided in the following theorem. In what follows, we state the main result regarding the existence and uniqueness of solutions for the system (4.1). The hypotheses are essentially the same as those in the previous sections; however, we present the next result under the assumption that the fields v, \mathcal{T} are globally Lipschitz.

Theorem 4.2. *Let us fix a filtered probability space $(\Omega, \mathcal{B}, \mathbb{P})$ endowed with a complete filtration $(\mathcal{F}_t)_{t \in [0, T]}$ generated by the Brownian motion $(\mathbf{B}(t))_{t \in [0, T]}$. Assume that $v_{\Psi} : \mathcal{Y} \rightarrow \mathbb{R}^d$, $(x, \lambda) \mapsto b(x)$, where $b : \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a Lipschitz velocity field with Lipschitz constant L_b . Assume that \mathcal{T} is defined as in (4.2) where α is a measurable function with respect to $\mathcal{B}(U) \times \mathcal{B}(\mathcal{Y})$, and it is Lipschitz in the second component with Lipschitz constant L_{α} . Furthermore, suppose that X_F is a uniformly bounded random variable, that is $\sup_{\omega \in \Omega} X_F^{\omega} < +\infty$, and that for all $i \in \mathbb{N}$, $(X_R, \Lambda^i(0)) \in L^2(\Omega, \mathcal{B}, \mathcal{P}; \mathcal{E})$. Then the system (4.1) admits a unique strong solution.*

Proof. In what follows, we use the same notation, and follow the argument used during the proof of Theorem 3.1. Since by hypothesis $\mathbb{P}(X_F = +\infty) = 0$, then we can take $\omega \in \Omega$ such that $X_F^{\omega} < +\infty$. We then propose the following Picard iteration. For $n \in \mathbb{N} \setminus \{0\}$, with an abuse of notation, we set

$$\left\{ \begin{array}{l} \mathbf{X}_{n+1}(t) = \mathbf{X}_0 + \int_0^t \mathbf{v}_{\Lambda_s}^N(\mathbf{X}_n(s), \boldsymbol{\lambda}_n(s)) ds + \sqrt{2\sigma} \mathbf{B}(t), \text{ if } \mathbf{X}_n(t^-) < \mathbf{X}_F, \\ \mathbf{X}_{n+1}(t) = \mathbf{X}_R + \int_0^t \mathbf{v}_{\Lambda_s}^N(\mathbf{X}_R(s), \boldsymbol{\lambda}_n(s)) ds + \sqrt{2\sigma} \mathbf{B}(t), \text{ if } \mathbf{X}_n(t^-) = \mathbf{X}_F, \\ \boldsymbol{\lambda}_{n+1}(t) = e^{-\frac{t}{\theta}} \boldsymbol{\lambda}_0 + \frac{1}{\theta} \int_0^t e^{-\frac{s-t}{\theta}} g_{\Lambda_s}(\mathbf{X}_n(s), \boldsymbol{\lambda}_n(s)) ds, \\ \mathbf{X}_0 := \overline{\mathbf{X}}_0, \\ \boldsymbol{\lambda}_0 := \overline{\boldsymbol{\lambda}}_0, \end{array} \right.$$

where $\mathbf{X}_R := \underbrace{(X_R, \dots, X_R)}_{N\text{-times}}$, and $\mathbf{X}_F := \underbrace{(X_F, \dots, X_F)}_{N\text{-times}}$, and we have borrowed the notation used in (3.1).


Notice that we have indicated $\mathbf{X}_n(t^-) < \mathbf{X}_F$ meaning that each component satisfies such an inequality. We then proceed as in proof of Theorem 3.1. Nevertheless, we need to consider two cases. First, we need to consider when $\mathbf{X}_n(t^-) < \mathbf{X}_F$, for fixed $n \in \mathbb{N}$. In that cases, we need to show that there exists a constant $R > 0$ such that

$$\mathbb{E} \left(\sup_{s \in [0, t]} \|\mathbf{Y}_{n+1}(s) - \mathbf{Y}_n(s)\|^2 \right) \leq \frac{(Rt)^n}{n!} \quad (4.5)$$

for all $n \in \mathbb{N}$, and all $t \in [0, T]$. Second, suppose that $\mathbf{X}_n(t^-) = \mathbf{X}_F$. Hence, we possibly find a further positive constant \widehat{R} such that (4.5) holds true. Then, we optimize between R and \widehat{R} to find a new constant R for which (4.5) holds true. From here, we may proceed with the proof of our desired result by following the same reasoning as in the proof of Theorem 3.1. \square

Remark 4.3. We proved Theorem 4.2 within a general framework in which X_F is a bounded random variable. However, in most neuronal models, the spiking threshold is a physiological parameter and is typically treated as a constant.

Having shown the existence of the solution, let us now show two pictures to illustrate the qualitative behavior of (4.1) in a simple case. In Figure 1, we observe the dynamics of $X^i(t)$ in the one-dimensional case for $N = 1000$, and where we have chosen $X_F = 0.7$, $X_R = 0.01$. To provide insights into the behavior of (4.1), we considered a linear system of the form $v_{\Lambda_t}^N(X^i(t), \lambda^i(t)) = a \cdot X^i(t) + b \cdot \lambda^i(t) + c \cdot \Lambda_t^i$, $\mathcal{T}_{\Lambda_t^N}(X^i(t), \lambda^i(t)) =$

FIGURE 1. Trajectories for X , and λ with $N = 1000$.

$d \cdot X^i(t) + e \cdot \lambda^i(t) + f \cdot \Lambda_t^N - \mathbb{E}(d \cdot X^i(t) + e \cdot \lambda^i(t) + f \cdot \Lambda_t^N)$, with $a = 0.5$, $b = 0.3$, $c = 0.2$; $d = 0.4$, $e = 0.2$, $f = 0.1$ (we omit the units since this is just an illustrative example). Notice that whenever $X^i(t) = X_R$, $\lambda^i(t)$ exhibits singularities in its trajectories.

4.1. Heterogeneous population of interacting neurons

In this section, we consider a more general version of the model studied in the previous sections. In particular, we include in the second equation of (2.9) an additional noise.

This network can be interpreted as a collection of neurons with different characteristics (resulting for instance in a different firing rate) and therefore, it is a model of an heterogeneous network. The weights distributions in cortex have been observed to be broad and neurons can belong to different subgroups with different behaviors. Moreover, the variance of the temporal fluctuations of the input currents may be large enough to endow the network with its own source of variability (Sect. 6.4.2 in [25]).

For these reasons we consider the system

$$\begin{cases} X^i(t) = X_0^i + \int_0^t v_{\Lambda_s^N}(X^i(s), \lambda^i(s)) ds + \sqrt{2\sigma} B^i(t), \\ \lambda^i(t) = \lambda_0^i + \int_0^t \mathcal{T}_{\Lambda_s^N}(X^i(s), \lambda^i(s) + R(s)) ds, \\ R(t) := \sum_{h \in \mathbb{N}} a_h W_h(t) e_h, \end{cases} \quad (4.6)$$

where $(W_h(t))_{h \in \mathbb{N}}$ is a sequence of independent Brownian motions, which are also independent of B^i for all $i \in \mathbb{N}$. Here, $(a_h)_{h \in \mathbb{N}}$ is a sequence of non-negative numbers such that $\sum_h a_h^2 < +\infty$, and $(e_h)_h$ is a sequence of signed measures supported in U such that $\int_U de_h(u) = 0$ for all $h \in \mathbb{N}$. The existence of strong solutions can be treated by following the same approach of Theorem 3.1. Indeed, we have the following result.

Proposition 4.4. *Let us fix a filtered probability space $(\Omega, \mathcal{B}, \mathbb{P})$ endowed with a complete filtration $(\mathcal{F}_t)_{t \in [0, T]}$ generated by the Brownian motion $(B(t))_{t \in [0, T]}$. Assume that for every $y \in \mathcal{Y}$ and $\Psi \in \mathcal{P}_1(\mathcal{Y})$ the velocity field $v_\Psi : \mathcal{Y} \rightarrow \mathbb{R}^d$ satisfies (A1)–(A3), and the operator \mathcal{T}_Ψ as defined in (4.2) satisfies (B1)–(B4). Moreover, let us consider $(\lambda_h)_{h \in \mathbb{N}}$ be a sequence of non-negative numbers such that $\sum_h \lambda_h^2 < +\infty$, and let $(W_h(t))_{h \in \mathbb{N}}$, $t \in [0, T]$ a sequence of independent real-valued Brownian motions, such that for all $t \in [0, T]$, $R(t) \in \mathcal{M}_0(U)$. Then, for every choice $Y_0 := (\bar{X}_0, \bar{\lambda}_0)$ as initial condition of (4.6) such that $Y_0 \in L^2(\hat{\mathcal{Y}})$, the system (4.6) has a unique solution Y that belongs to $\mathcal{M}_{\text{ad}}^2(\hat{\mathcal{Y}})$.*

Proof. Notice that for almost all $\omega \in \Omega$, $\mathcal{J}_{\Lambda_t^N}(X^i(t), \lambda^i(t) + R(t))$ satisfies our assumptions (B1)–(B4). Therefore, we may apply Theorem 3.1 by replacing $\mathcal{J}_{\Lambda_t^N}$ with $\bar{\mathcal{J}}_{\Lambda_t^N}(X^i(t), \lambda^i(t)) := \mathcal{J}_{\Lambda_t^N}(X^i(t), \lambda^i(t) + R(t))$, and where (4.6) can be written as

$$\begin{cases} X^i(t) = X_0^i + \int_0^t v_{\Lambda_s^N}(X^i(s), \lambda^i(s)) ds + \sqrt{2\sigma} B^i(t), \\ \lambda^i(t) = \lambda_0^i + \int_0^t \bar{\mathcal{J}}_{\Lambda_s^N}(X^i(s), \lambda^i(s)) ds. \end{cases}$$

Moreover, in order to apply the Picard iteration used in Theorem 3.1, we define $\bar{g}_{\Lambda_t}(X_n^i(t), \lambda_n^i(t)) := \lambda_n^i(t) + \theta \bar{\mathcal{J}}_{\Lambda_t^N}(X_n^i(t), \lambda_n^i(t))$ for some suitable $\theta > 0$. Then, we consider the Picard iteration (3.1) with g_{Λ_t} replaced by \bar{g}_{Λ_t} . That is, we have

$$\begin{cases} X_{n+1}^i(t) = X_0^i + \int_0^t \mathbf{v}_{\Lambda_s^N}^N(X_n^i(s), \lambda_n^i(s)) ds + \sqrt{2\sigma} B^i(t), \\ \lambda_{n+1}^i(t) = e^{-\frac{t}{\theta}} \lambda_0^i + \frac{1}{\theta} \int_0^t e^{\frac{s-t}{\theta}} \bar{g}_{\Lambda_s}(X_n^i(s), \lambda_n^i(s)) ds, \end{cases}$$

and thus we proceed by following the reasoning of Theorem 3.1, and our conclusion follows. \square

5. POSSIBLE FUTURE DIRECTIONS

Notice that for equations of the form (2.11), we have assumed that hypothesis C1 holds true to derive the existence and uniqueness of Lagrangian solutions. In the case of equations of the form (4.1), we are unable to formulate such a condition in a manner that takes into account the effect of the random variable X_F , which controls the spike times of X . However, since simulations show that the trajectories of λ are C^0 and the trajectories of X are almost the same as those we can simulate through (2.9), we conjecture that it is possible to obtain the existence of both Lagrangian and Eulerian solutions. Nonetheless, this remains a point for further investigation. Additionally, we would like to reformulate all the results by relaxing assumption (C1). Moreover, we seek to analyze the asymptotic behavior of a system of particles as time diverges. More precisely, we are interested in whether the continuity equation governed by the operator in assumption (C1) admits a stationary solution. In this aspect we intend to compare our analysis with results such as those in [44]. Specifically, we aim to determine a decay rate for the family of probability measures Λ_t^N using a Doeblin-type condition; see, for instance, [45], Theorem 2.3.

ACKNOWLEDGMENTS

The authors would like to thank Alessandro Baldi and Marco Morandotti for inspiring discussions that greatly influenced this work. The author G.D. acknowledges financial support under the MIUR-PRIN 2022 project “Non-Markovian dynamics and non-local equations”, No. 202277N5H9 – CUP: D53D23005670006. G.D. participates also in the INdAM – GNAMPA Project, *Deterministic Control of Stochastic Dynamics* CUP E53C23001670001. AMH has been supported by project PRIN 2022 “understanding the LEarning process of QUantum Neural networks (LeQun)”, proposal code 2022WHZ5XH – CUP J53D23003890006.

DATA AVAILABILITY STATEMENT

The research data associated with this article are included in the article.

REFERENCES

- [1] T. Vicsek, A. Czirók, E. Ben-Jacob, I. Cohen and O. Shochet, Novel type of phase transition in a system of self-driven particles. *Phys. Rev. Lett.* **75** (1995) 1226–1229.
- [2] J.-M. Lasry and P.-L. Lions, Mean field games. *Jpn. J. Math.* **2** (2007) 229–260.

- [3] F. Delarue, J. Inglis, S. Rubenthaler and E. Tanré, Particle systems with a singular mean-field self-excitation. application to neuronal networks. *Stoch. Processes Appl.* **125** (2015) 2451–2492.
- [4] B. Duchet, C. Bick and Á. Byrne, Mean-field approximations with adaptive coupling for networks with spike-timing-dependent plasticity. *Neural Computat.* **35** (2023) 1481–1528.
- [5] G. Dumont and P. Gabriel, The mean-field equation of a leaky integrate-and-fire neural network: measure solutions and steady states. *Nonlinearity* **33** (2020) 6381.
- [6] P. Robert and J. Touboul, On the dynamics of random neuronal networks. *J. Statist. Phys.* **165** (2016) 545–584.
- [7] A. Treves, Mean-field analysis of neuronal spike dynamics. *Network Computat. Neural Syst.* **4** (1983) 259–284.
- [8] J. Han, Q. Li, *et al.*, A mean-field optimal control formulation of deep learning. *Res. Math. Sci.* **6** (2019) 1–41.
- [9] J. Sirignano and K. Spiliopoulos, Mean field analysis of deep neural networks. *Math. Oper. Res.* **47** (2022) 120–152.
- [10] P. Cardaliaguet and S. Hadikhhanloo, Learning in mean field games: the fictitious play. *ESAIM: Control Optim. Calc. Var.* **23** (2017) 569–591.
- [11] H. Pham and X. Wei, Bellman equation and viscosity solutions for mean-field stochastic control problem. *ESAIM: Control Optim. Calc. Var.* **24** (2018) 437–461.
- [12] L. Popovic and A. Véber, A spatial measure-valued model for chemical reaction networks in heterogeneous systems. *Ann. Appl. Probab.* **33** (2023) 3706–3754.
- [13] M. Morandotti and F. Solombrino, Mean-field analysis of multipopulation dynamics with label switching. *SIAM J. Math. Anal.* **52** (2020) 1427–1462.
- [14] S. Almi, M. Morandotti and F. Solombrino, Optimal control problems in transport dynamics with additive noise. *J. Differ. Equ.* **373** (2023) 1–47.
- [15] M. Fornasier, S. Lisini, C. Orrieri and G. Savaré, Mean-field optimal control as gamma-limit of finite agent controls. *Eur. J. Appl. Math.* **30** (2019) 1153–1186 .
- [16] A. Baldi, *Well-posedness and propagation of chaos for multi-agent models with strategies and diffusive effects*. Master’s thesis, Politecnico di Torino (2023).
- [17] A. Baldi and M. Morandotti, Well-posedness and propagation of chaos for multi-agent models with strategies and diffusive effects. arXiv preprint arXiv:2507.14058 (2025).
- [18] L. Ambrosio, M. Fornasier, M. Morandotti and G. Savaré, Spatially inhomogeneous evolutionary games. *Commun. Pure Appl. Math.* **71** (2018) 1353–1402.
- [19] G. Ascione, D. Castorina and F. Solombrino, Mean-field sparse optimal control of systems with additive white noise. *SIAM J. Math. Anal.* **55** (2023) 6965–6990.
- [20] P. Grazieschi, M. Leocata, C. Mascart, J. Chevallier, F. Delarue and E. Tanré, Network of interacting neurons with random synaptic weights. *ESAIM: Proc. Surv.* **65** (2019) 445–475.
- [21] O. Faugeras, E. Soret and E. Tanré, Asymptotic behavior of a network of neurons with random linear interactions. arXiv preprint arXiv:2006.03083 (2020).
- [22] O.D. Faugeras, J.D. Touboul and B. Cessac, A constructive mean-field analysis of multi population neural networks with random synaptic weights and stochastic inputs. *Front. Computat. Neurosci.* **3** (2009) 323.
- [23] A. De Masi, A. Galves, E. Löcherbach and E. Presutti, Hydrodynamic limit for interacting neurons. *J. Statist. Phys.* **158** (2015) 866–902.
- [24] N. Fournier and E. Löcherbach, On a toy model of interacting neurons. *Ann. Inst. Henri Poincaré Probab. Statist.* **52** (2016) 1844–1876.
- [25] G. La Camera, The mean field approach for populations of spiking neurons, in *Computational Modelling of the Brain: Modelling Approaches to Cells, Circuits and Networks*. Springer (2021) 125–157.
- [26] M. Tamborrino and P. Lansky, Shot noise, weak convergence and diffusion approximations. *Physica D: Nonlinear Phenomena* **418** (2021) 132845.
- [27] G. Ascione, G. D’Onofrio, L. Kostal and E. Pirozzi, An optimal Gauss–Markov approximation for a process with stochastic drift and applications. *Stoch. Processes Appl.* **130** (2020) 6481–6514.
- [28] P.H Baxendale and P.E. Greenwood, Sustained oscillations for density dependent Markov processes. *J. Math. Biol.* **63** (2011) 433–457.
- [29] S. Ditlevsen and P. Greenwood, The Morris–Lecar neuron model embeds a leaky integrate-and-fire model. *J. Math. Biol.* **67** (2013) 239–259.
- [30] Q. Cormier, E. Tanré and R. Veltz, Long time behavior of a mean-field model of interacting neurons. *Stoch. Processes Appl.* **130** (2020) 2553–2595.

- [31] E. Löcherbach, Fluctuations for mean field limits of interacting systems of spiking neurons. *Ann. Inst. Henri Poincaré Probab. Statist.* **60** (2024) 790–823.
- [32] G. Ascione and G. D'Onofrio, Deterministic control of SDEs with stochastic drift and multiplicative noise: a variational approach. *Appl. Math. Optim.* **88** (2023) 11.
- [33] E. Löcherbach, Spiking neurons: interacting Hawkes processes, mean field limits and oscillations. *ESAIM: Proc. Surv.* **60** (2017) 90–103.
- [34] V. Panaretos and Y. Zemel, *An Invitation to Statistics in Wasserstein Space*. SpringerBriefs in Probability and Mathematical Statistics. Cambridge University Press (2020).
- [35] R.F. Arens and J. Eells Jr., On embedding uniform and topological spaces. *Pacific J. Math.* **6** (1956) 397–403.
- [36] Z.-M. Ma and M. Röckner, Introduction to the Theory of (Non-symmetric) Dirichlet Forms. Springer Science & Business Media (2012).
- [37] G. Da Prato and J. Zabczyk, Stochastic Equations in Infinite Dimensions, vol. 152. Cambridge University Press (2014).
- [38] A.-S. Sznitman, Topics in propagation of chaos. *Ecole Probab. Saint-Flour XIX* **1464** (1991) 165–251.
- [39] L.-P. Chaintron and A. Diez, Propagation of chaos: a review of models, methods and applications. I. models and methods. *Kinetic Related Models* **15** (2022) 895.
- [40] L.-P. Chaintron and A. Diez, Propagation of chaos: a review of models, methods and applications. II. applications. *Kinetic Related Models* **15** (2022) 1017.
- [41] P. Billingsley, Convergence of probability measures. Wiley Series in Probability and Statistics: Probability and Statistics, 2nd edn. John Wiley & Sons, Inc., New York (1999). A Wiley-Interscience Publication.
- [42] A. Dembo and O. Zeitouni, Large Deviations Techniques and Applications. Jones and Bartlett Publishers, Boston, MA (1993).
- [43] P. Lánský and V. Lanska, Diffusion approximation of the neuronal model with synaptic reversal potentials. *Biol. Cybernet.* **56** (1987) 19–26.
- [44] K. Pakdaman, B. Perthame and D. Salort, Dynamics of a structured neuron population. *Nonlinearity* **23** (2009) 55.
- [45] J.A. Cañizo and H. Yoldaş, Asymptotic behaviour of neuron population models structured by elapsed-time. *Nonlinearity* **32** (2019) 464.
- [46] H. Brézis, *Opérateurs maximaux monotones et semi-groupes de contractions dans les espaces de Hilbert*. North-Holland Mathematics Studies. North-Holland, Amsterdam–London; American Elsevier, New York (1973).
- [47] L. Manca, Kolmogorov operators in spaces of continuous functions and equations for measures. Vol. 10 of *Tesi. Scuola Normale Superiore di Pisa (Nuova Series)*. Edizioni della Normale, Pisa (2008).

Please help to maintain this journal in open access!



This journal is currently published in open access under the Subscribe to Open model (S2O). We are thankful to our subscribers and supporters for making it possible to publish this journal in open access in the current year, free of charge for authors and readers.

Check with your library that it subscribes to the journal, or consider making a personal donation to the S2O programme by contacting subscribers@edpsciences.org.

More information, including a list of supporters and financial transparency reports, is available at <https://edpsciences.org/en/subscribe-to-open-s2o>.

APPENDIX A. WELL-POSEDNESS OF ODES IN BANACH SPACES

Let us recall the following Theorem by Brezis [46], Theorem 1.4 about the well-posedness of ODEs in Banach spaces.

Theorem A.1. [46] *Let $(E, \|\cdot\|_E)$ be a Banach space, C a closed convex subset of E , and let $A(t, \cdot) : C \rightarrow E$, $t \in [0, T]$, be a family of operators satisfying the following properties:*

(i) there exists a constant $L \geq 0$ such that for every $c_1, c_2 \in C$ and $t \in [0, T]$

$$\|A(t, c_1) - A(t, c_2)\|_E \leq L \|c_1 - c_2\|_E;$$

(ii) for every $c \in C$ the map $t \mapsto A(t, c)$ is continuous in $[0, T]$;
 (iii) for every $R > 0$ there exists $\theta > 0$ such that

$$c \in C, \|c\|_E \leq R \Rightarrow c + \theta A(t, c) \in C.$$

Then for every $\bar{c} \in C$ there exists a unique curve $c : [0, T] \rightarrow C$ of class C^1 satisfying $c_t \in C$ for all $t \in [0, T]$ and

$$\frac{d}{dt} c_t = A(t, c_t) \text{ in } [0, T], c_0 = \bar{c}.$$

Moreover, if c^1, c^2 are the solutions starting from the initial data $\bar{c}^1, \bar{c}^2 \in C$, respectively, we have

$$\|c_t^1 - c_t^2\| \leq e^{Lt} \|\bar{c}^1 - \bar{c}^2\|_E \text{ for every } t \in [0, T].$$

In what follows, we consider the following generalization of the previous result proved in [13], Corollary 2.3.

Corollary A.2 ([13], Cor. 2.3). *Let hypotheses (ii) and (iii) of Theorem A.1 hold for a family of operators $A(t, \cdot) : C \rightarrow E$, $t \in [0, T]$. Assume in addition, that*

(i') for every $R > 0$ there exists a constant $L_R \geq 0$ such that for every $c_1, c_2 \in C \cap B_R$ and $t \in [0, T]$,

$$\|A(t, c_1) - A(t, c_2)\|_E \leq L_R \|c_1 - c_2\|_E;$$

(ii') there exists $M > 0$ such that for every $c \in C$, there holds

$$\|A(t, c)\|_E \leq M(1 + \|c\|_E).$$

Then for every $\bar{c} \in C$ there exists a unique curve $c : [0, T] \rightarrow C$ of class C^1 satisfying $c_t \in C$ for all $t \in [0, T]$, and

$$\frac{d}{dt} c_t = A(t, c_t) \text{ in } [0, T], c_0 = \bar{c}.$$

Moreover, if c^1, c^2 are the solutions starting from the initial data $\bar{c}^1, \bar{c}^2 \in C \cap B_R$, respectively, there exists a constant $L = L(M, R, T) > 0$ such that

$$\|c_t^1 - c_t^2\| \leq e^{Lt} \|\bar{c}^1 - \bar{c}^2\|_E \text{ for every } t \in [0, T].$$

APPENDIX B. A KOLMOGOROV EQUATION FOR MEASURES

Here, suppose that X is a separable Banach space with norm $\|\cdot\|$, and let $\mathcal{B}(X)$ be its Borel σ -algebra. We denote by $C_b(X)$, the Banach space of all uniformly continuous and bounded functions $f : X \rightarrow \mathbb{R}$, endowed with the supremum norm $\|\cdot\|_0$. Furthermore, we consider a strongly continuous semigroup of linear operators $\{P_t\}_{t \geq 0} \subset \mathcal{B}(C_b(X))$ with the Markovian property, that is, there exists a family $\{\pi_t(x, \cdot) : t \geq 0, x \in X\}$ of probability measures on X such that

- the map $\mathbb{R}_+ \times X \rightarrow [0, 1]$, $(t, x) \mapsto \pi_t(x, \Gamma)$ is measurable, for any Borel set $\mathcal{B}(X)$;

- $\pi_{t+s}(x, \Gamma) = \int_X \pi_s(y, \Gamma) \pi_t(x, dy)$ for all $t, s \geq 0$, $x \in X$, and $\Gamma \in \mathcal{B}(X)$;
- for any $x \in X$, $\pi_0(x, \cdot) = \delta_x(\cdot)$, the probability measure concentrated in x ;
- $P_t \phi(x) = \int_X \phi(y) \pi_t(x, dy)$, for any $t \geq 0$, $\phi \in C_b(X)$, $x \in X$;
- for any $\phi \in C_b(X)$, $x \in X$, the function $\mathbb{R}_+ \rightarrow \mathbb{R}$, $t \mapsto P_t \phi(x)$ is continuous.

Then we define the linear operator $(\mathcal{L}, D(\mathcal{L}))$ where

$$\begin{aligned} D(\mathcal{L}) &:= \left\{ u \in X : \lim_{t \downarrow 0} \frac{P_t u - u}{t} \text{ exists} \right\}, \\ \mathcal{L}u &:= \lim_{t \downarrow 0} \frac{P_t u - u}{t} \quad u \in D(\mathcal{L}). \end{aligned} \tag{B.1}$$

Definition B.1. Let $\bar{\mu} \in \mathcal{M}_b(X)$. We say that a family of measures $\{\mu_t\}_{t \geq 0}$ is a solution of the measure equation

$$\begin{cases} \frac{d}{dt} \int_X \phi \mu_t(dx) = \int_X \mathcal{L} \phi(x) \mu_t(dx), & t \geq 0, \phi \in D(\mathcal{L}), \\ \mu_0 = \bar{\mu}, & \mu \in \mathcal{M}_b(X), \end{cases} \tag{B.2}$$

if the following conditions are fulfilled:

- the total variation of the measure μ_t satisfies

$$\int_0^T \|\mu_t\|_{TV} dt < +\infty, \quad T > 0;$$

- for any $\phi \in D(\mathcal{L})$, the real valued function $t \mapsto \int_X \phi(x) \mu_t(dx)$ is absolutely continuous, and for any $t \geq 0$ it holds

$$\int_X \phi(x) \mu_t(dx) - \int_X \phi(x) \bar{\mu}(dx) = \int_0^t \int_X \mathcal{L} \phi(x) \mu_s(dx) ds.$$

Let us now recall an important result about the well-posedness of (B.2) when X is a Hilbert space, [47], Theorem 1.2.

Theorem B.2. Let $\{P_t\}_{t \geq 0}$ be a strongly continuous Markov semigroup and let $(\mathcal{L}, D(\mathcal{L}))$ be its infinitesimal generator, defined as in (B.1). Then the formula

$$\langle \phi, P_t^* F \rangle_{\mathcal{B}(C_b(X), (C_b(X))^*)} := \langle F, P_t \phi \rangle_{\mathcal{B}(C_b(X), (C_b(X))^*)}$$

defines a semigroup $(P_t^*)_{t \geq 0}$ of linear and continuous operators on $(C_b(X))^*$ that maps $\mathcal{M}_b(X)$ into $\mathcal{M}_b(X)$. Moreover, for any $\bar{\mu} \in \mathcal{M}_b(X)$, $\phi \in C_b(X)$ the map $t \mapsto \int_X \phi(x) P_t^* \bar{\mu}(dx)$ is continuous, and if $\phi \in D(\mathcal{L})$ then it is also differentiable with continuous differential

$$\frac{d}{dt} \int_X \phi(x) P_t^* \bar{\mu}(dx) = \int_X \mathcal{L} \phi(x) P_t^* \bar{\mu}(dx). \tag{B.3}$$

Finally, for any $\mu \in \mathcal{M}_b(X)$ there exists a unique solution of the measure equation (B.2) given by $\{P_t^* \mu\}_{t \geq 0}$.

Proof. The proof of this theorem can be performed by using the same argument as the one in [47], Theorem 1.2. The only difference is that we consider a separable Banach space rather than a Hilbert space. \square

APPENDIX C. ALDOUS TIGHTNESS CRITERION

Let E be a Polish space, and for each $m \in \mathbb{N}$, we denote by $\mathcal{D}_E^m = \mathcal{D}_E[0, m]$ the Skorokhod space. That is, the space of all càdlàg trajectories defined on $[0, m]$ with values in E . In the same way, we denote by \mathcal{D}_E^∞ the Skorokhod space $\mathcal{D}_E[0, +\infty)$. Let X_n be random variables of \mathcal{D}_E^∞ .

Theorem C.1 ([41], Thm. 16.10). *Suppose that*

1.

$$\lim_{a \rightarrow +\infty} \limsup_n \mathbb{P}(\|X_n\|_{\mathcal{D}_E^m} \geq a) = 0,$$

holds for all $m \in \mathbb{N}$.

2. *For each $\varepsilon > 0, \eta > 0, m$ there exists $\delta_0 > 0$ and $N_0 \in \mathbb{N}$ such that, if $\delta \leq \delta_0$, and $N \geq N_0$, and if τ is a discrete X_n -stopping time satisfying $\tau \leq m$, then*

$$\mathbb{P}(|X_n(\tau + \delta) - X_n(\tau)| \geq \varepsilon) \leq \eta.$$

Then X_n is tight.