

Ecohydrological monitoring of two Alpine ecosystems representing possible broader future conditions in the Alps

Original

Ecohydrological monitoring of two Alpine ecosystems representing possible broader future conditions in the Alps / Hamza, T., Gisolo, D., Gentile, A., Canone, D., Ferraris, S.. - (2025). (EGU General Assembly 2025) [10.5194/egusphere-egu25-11799].

Availability:

This version is available at: 11583/3008991 since: 2026-03-20T13:23:20Z

Publisher:

Copernicus GmbH

Published

DOI:10.5194/egusphere-egu25-11799

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)



Ecohydrological monitoring of two Alpine ecosystems representing possible broader future conditions in the Alps

Tanzeel Hamza et al. ▶

High-altitude non-glacial grasslands and peri-urban new-growth forests are still two poorly studied ecosystems that represent present and future conditions in the Alps. Hence, studying the functioning of these environments is crucial, especially if land surface models' capability of representing the ecosystems' processes is assessed.

The data are collected at two eddy covariance sites located in the Northwestern Alps, respectively on a high-altitude grassland (2550 m a.s.l.) and in a forest (650 m a.s.l. with a 25 m high mast). The data are characterised by time series spanning 365 days per year, since 2018 for the grassland site and since 2021 for the forest site.

The ecosystem and soil information (energy fluxes and actual evapotranspiration, ET_a, soil moisture) obtained from measurements is combined with simulation results obtained with the land surface CLM model (The Community Land Model, NCAR, US). A rather good agreement is found between observations and simulations.

A particular focus on dry conditions in 2022 on the forest site is also presented. The results show that soil moisture, pressure head, and net radiation are more important than vapor pressure deficit, wind speed, and air temperature as ET_a drivers. During the drought, despite the low soil moisture, both cases of water- and energy-limited conditions occurred. A weak effect of the drought on ET_a is observed, likely due to the deep root system. The cosmic ray neutron sensor (CRNS) measurements revealed a good agreement with capacitive probes profile (CPS) ones.

This work was supported by the NODES project, funded under MUR – M4C2 1.5 of the PNRR with resources from the European Union - NextGenerationEU (Grant agreement no. ECS00000036), as well as the MUR PRIN project SUNSET (202295PFKP_003).

How to cite: Hamza, T., Gisolo, D., Gentile, A., Canone, D., and Ferraris, S.: Ecohydrological monitoring of two Alpine ecosystems representing possible broader future conditions in the Alps, EGU General Assembly 2025, Vienna, Austria, 27 Apr–2 May 2025, EGU25-11799, <https://doi.org/10.5194/egusphere-egu25-11799>, 2025.