

Relational Bundle Geometric Formulation of Non-Relativistic Quantum Mechanics

Original

Relational Bundle Geometric Formulation of Non-Relativistic Quantum Mechanics / Francois, J.T., Ravera, L.. - In: FORTSCHRITTE DER PHYSIK. - ISSN 0015-8208. - ELETTRONICO. - 73:12(2025), pp. 1-24. [10.1002/prop.70040]

Availability:

This version is available at: 11583/3012407 since: 2026-06-24T14:47:54Z

Publisher:

John Wiley and Sons

Published

DOI:10.1002/prop.70040

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Relational Bundle Geometric Formulation of Non-Relativistic Quantum Mechanics

J. T. François* and L. Ravera*

A bundle geometric formulation of non-relativistic many-particles Quantum Mechanics is presented. A wave function is seen to be a $\mathbb{C}$ -valued *cocyclic* tensorial 0-form on configuration space-time seen as a principal bundle, while the Schrödinger equation flows from its covariant derivative, with the action functional supplying a (flat) *cocyclic connection* 1-form on the configuration bundle. In line with the historical motivations of Dirac and Feynman, ours is thus a *Lagrangian* geometric formulation of QM, in which the Dirac–Feynman path integral arises in a geometrically natural way.

Applying the *dressing field method*, we obtain a relational reformulation of this geometric non-relativistic QM: a relational wave function is realised as a basic cocyclic 0-form on the configuration bundle. In this relational QM, any particle position can be used as a dressing field, i.e., as a “physical reference frame.”

The dressing field method naturally accounts for the freedom in choosing the dressing field, which is readily understood as a covariance of the relational formulation under changes of physical reference frame.

1. Introduction

Contrary to the other fundamental pillars of modern physics, General Relativity (GR) and Gauge Field Theory (GFT)—or broadly, general-relativistic Gauge Field Theory (gRGFT)—which are fundamentally understood in geometric terms, Quantum Mechanics (QM) is usually still primarily conceived algebraically. Indeed many of its key puzzling features, from an interpretive standpoint, stems from its algebraic linearity. Yet, the geometric structure of QM has been a topic of interest for many. Notably, the field of geometric quantization^[1] pioneered by Souriau, Kostant, and Kirillov has been fruitful, mainly in mathematics and group representation theory. Also, isolated works throughout the years have made interesting contributions, e.g., refs. [2–4].¹

As part of our program of a relational (re)formulation of fundamental physics, started in refs. [6, 7] (see also refs. [8, 9]), in this work, we propose an original contribution to this tradition. We give a bundle geometric formulation of non-relativistic QM, which relies on a *generalisation* of standard bundle geometric notions, where group representations are replaced by 1-cocycles for the action of the group: *cocyclic bundle geometry*^[10]—originally called *twisted bundle geometry*. A key notion of which is that of *cocyclic connection*, extending the usual Ehresmann (and/or Cartan) connections.² We further stress that ours is naturally a *Lagrangian* approach to QM, which soundly resonates with the historical motivations of Dirac^[13] and Feynman,^[14,15] leading them to the path integral (PI) formulation. As a matter of fact, the Dirac–Feynman PI fits well with our geometric approach.

Our bundle geometric version of QM makes it possible to apply the *dressing field method* (DFM) of symmetry reduction,^[7,16,17]

J. T. François
Department of Mathematics & Statistics
Masaryk University – MUNI
Kotlářská 267/2, Veveří, Brno, Czech Republic
E-mail: jordan.francois@uni-graz.at

J. T. François
Department of Philosophy – University of Graz
Heinrichstrasse 26/5, Graz 8010, Austria

J. T. François
Department of Physics
Mons University – UMONS
20 Place du Parc, Mons 7000, Belgium

L. Ravera
DISAT
Politecnico di Torino – PoliT
Corso Duca degli Abruzzi 24, Torino 10129, Italy
E-mail: lucrezia.ravera@polito.it

L. Ravera
Istituto Nazionale di Fisica Nucleare
Section of Torino – INFN
Via P. Giuria 1, Torino 10125, Italy

L. Ravera
Grupo de Investigación en Física Teórica – GIFT
Universidad Católica De La Santísima Concepción
Concepción, Chile

 The ORCID identification number(s) for the author(s) of this article can be found under <https://doi.org/10.1002/prop.70040>

© 2025 The Author(s). Fortschritte der Physik published by Wiley-VCH GmbH. This is an open access article under the terms of the [Creative Commons Attribution](#) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

DOI: 10.1002/prop.70040

¹ Recently was published the result^[5] of a years-long concerted effort to couch QM in differential geometric terms. Our approach, to be described momentarily, is entirely independent.

² The notion of cocyclic (or twisted) connection features prominently in twistor geometry,^[11] as well as in its real counterpart, conformal tractor geometry^[12] – and in projective geometry. Tractor and twistors are cocyclic gauge fields regarding Weyl rescaling symmetry, and the conformal tractor and twistor connections are cocyclic Cartan connections.

³ Dirac sought a Lagrangian formulation of QM so that it could be made compatible with Special Relativity from first principles. Feynman was motivated to find a Lagrangian approach to QM by the particular approach he and Wheeler took at the time to tackle QED, which rested on a classical version of electrodynamics that could not be cast in an Hamiltonian form.

thereby obtaining a geometric *relational* formulation of QM. It is distinct from relational QM à la Rovelli,^[18] even though the interpretation of our results is broadly in line with it. In spirit, it is also broadly in line with the heuristic idea behind the recent burgeoning literature on “quantum reference frames,”^[19–21] though we approach the subject in a mathematically more sophisticated way, akin to geometric quantization.

The paper is organised as follows: For the benefit of the reader, in Section 2, we provide the essential background material. In Section 2.1, we remind the basics of bundle geometry. This is necessary for two reasons: First to introduce *cocyclic* bundle geometry, in terms of which we shall formulate QM, and second because bundle geometry is the proper framework to express the DFM, which we do in Section 2.2.

With this preparatory work done, in Section 3, we give our bundle geometric formulation of non-relativistic QM. The configuration space $\mathcal{C}_t$ of a set of N particles at a time t is seen as the fiber of a principal bundle $\mathcal{C}$, which we call the configuration space-time bundle, over the Newtonian time line $\mathcal{T}$, its base space. The gauge group of $\mathcal{C}$ is shown to exactly capture the so-called *extended Galilean transformations*.^[22] The action functional is shown to induce a *flat cocyclic connection* 1-form. It induces a *cocyclic covariant derivative* on $\mathbb{C}$ -valued *cocyclic tensorial forms* on $\mathcal{C}$. Quantum mechanical wave functions ψ are the subspace of form degree 0 of its kernel. The cocyclic covariant derivative of ψ splits—in bundle coordinates—as two equations: One gives the quantum mechanical prescription for the conjugate momentum operator $\hat{\pi} = -i\hbar \frac{\partial}{\partial x}$, while the other gives the Schrödinger equation. We then show how the Dirac–Feynman PI formulation may arise naturally from this geometric framework.

In Section 4, we describe a bundle geometric relational reformulation of both classical and quantum mechanics, achieved via the DFM. Relational wave functions are realized as *dressed*, or *basic*, 0-forms on $\mathcal{C}$. Correspondingly, a *relational Schrödinger equation* gives the quantum dynamics. The DFM further naturally accounts for the freedom in choosing the dressing field: in this case, any particle position can be used as a dressing field, i.e., as a *physical reference frame*. The dressed, relational formulation of QM via the DFM thus automatically implements a “covariance under change of physical reference frame.”

In the conclusion 5, we discuss the salient points our results and we indicate the next phase of our program. The main text being kept general, Appendix A illustrates our formalism for free particles.

2. Bundle Geometry and the Dressing Field Method

2.1. Cocyclic Bundle Geometry in a Nutshell

The differential geometry of principal bundles is the underpinning of our formulation of non-relativistic classical and quantum mechanics, and of the Dressing Field Method.

The central object is a principal bundle P over a base manifold M with structure group H and projection $\pi : P \rightarrow M$, $p \mapsto \pi(p) = x$. A fiber over $x \in M$, $\pi^{-1}(x) = P|_x \subset P$, is an orbit of the right action of the structure group, $P \times H \rightarrow P$, $(p, h) \mapsto ph =: R_h p$, which is free and transitive on fibers, and s.t. $\pi \circ R_h = \pi$. This means the base is isomorphic as a manifold to the space of fibers $P/H \simeq M$.

The linearisation of the action of H induces vectors tangent to the fibers: $\forall X \in \mathfrak{h}$ corresponds a *fundamental* vertical vector X_p^ν at p . At $p \in P$, the span of these vectors is a subvector space $V_p P$ of the tangent space $T_p P$. The canonical vertical subbundle is $VP = \cup_p V_p P \subset TP$. We denote by $\Gamma(VP)$ the space of sections of VP , i.e., vertical vector fields $X^\nu : P \rightarrow VP$.

The natural maximal group of transformation of P is its group of automorphisms

$$\text{Aut}(P) := \{ \Xi \in \text{Diff}(P) \mid \Xi(ph) = \Xi(p)h \}. \quad (1)$$

It is the subgroup of $\text{Diff}(P)$ that preserves the fibration structure by sending fibers to fibers, and thus induces diffeomorphisms of the base space M , i.e., projects onto $\text{Diff}(M)$. Its maximal normal subgroup is the group of vertical automorphisms

$$\text{Aut}_v(P) := \{ \Xi \in \text{Aut}(P) \mid \pi \circ \Xi = \pi \}. \quad (2)$$

It is the subgroup of those automorphisms that induce the identity transformation on M . It is isomorphic to the *gauge group* of P ,

$$\mathcal{H} := \{ \gamma : P \rightarrow H \mid R_h^* \gamma = h^{-1} \gamma h \}, \quad (3)$$

the isomorphism being given by $\Xi(p) = p\gamma(p)$. The action of $\text{Aut}_v(P) \simeq \mathcal{H}$ on differential forms of P defines their *active gauge transformations*. We have thus the characteristic short exact sequence (SES) of groups of P ,

$$\text{id}_P \rightarrow \text{Aut}_v(P) \simeq \mathcal{H} \xrightarrow{\bar{\cdot}} \text{Aut}(P) \xrightarrow{\bar{\cdot}} \text{Diff}(M) \rightarrow \text{id}_M, \quad (4)$$

whose linear version, the SES of Lie algebras, characterizes the Atiyah Lie algebroid canonically associated to P .

Spaces of Remarkable Differential Forms: As a smooth manifold P has a graded ring of differential forms $\Omega^*(P) = \bigoplus_{k \geq 0} \Omega^k(P)$, with graded Lie algebra of derivations $\text{Der}^*(\Omega^*(P)) = \bigoplus_{k \geq 0} \text{Der}^k(\Omega^*(P))$ s.t. for $D_1 \in \text{Der}^p$ and $D_2 \in \text{Der}^q$ we have $[D_1, D_2] := D_1 \circ D_2 - (-1)^{pq} D_2 \circ D_1 \in \text{Der}^{p+q}$. In particular, we have the exterior de Rham derivative $d \in \text{Der}^1$, the inner derivation $\iota_X \in \text{Der}^{-1}$ for any $X \in \Gamma(TP)$, and the Lie derivative $\mathfrak{L}_X := [d, \iota_X]$. The de Rham complex of P is $(\Omega^*(P), d)$.

The action of the structure group on P induces a right action on forms $\Omega^*(P) \times H \rightarrow \Omega^*(P)$, $(\alpha, h) \mapsto R_h^* \alpha$, which defines their *equivariance*. The action by pullback of the group of automorphisms is also a right action: $\Omega^*(P) \times \text{Aut}(P) \rightarrow \Omega^*(P)$, $(\alpha, \Xi) \mapsto \Xi^* \alpha$. In particular, gauge transformations are defined by the right action $\Omega^*(P) \times \text{Aut}_v(P) \simeq \mathcal{H} \rightarrow \Omega^*(P)$, $(\alpha, \Xi \sim \gamma) \mapsto \Xi^* \alpha =: \alpha^\gamma$.

There are a number of remarkable spaces of forms worth describing. The first, which can be defined canonically, are *horizontal* forms, i.e., those that vanish when evaluated on vertical vector fields

$$\Omega_{\text{hor}}^*(P) := \{ \alpha \in \Omega^*(P) \mid \iota_{X^\nu} \alpha = 0 \text{ for } X^\nu \in \Gamma(VP) \}. \quad (5)$$

Then, there is the space of *equivariant* forms, whose equivariance is “simple.” In standard bundle geometry, equivariant forms are valued in representations (ρ, V) of H , which control their equiv-

ariance:

$$\Omega_{\text{eq}}^*(P, \rho) := \{\alpha \in \Omega^*(P, V) \mid R_h^* \alpha = \rho(h)^{-1} \alpha\}. \quad (6)$$

Their linear equivariance is then given by the Lie derivative along fundamental vector fields: $\mathfrak{L}_{X^v} \alpha = -\rho_*(X) \alpha$.

This is generalised in *cocyclic* bundle geometry (formerly introduced as *twisted* bundle geometry^[10]), where one defines *cocyclic* equivariant forms whose equivariance is controlled by a 1-cocycle for the action of the structure group: Given a Lie group G we define

$$C : P \times H \rightarrow G, \quad (7)$$

$$(p, h) \mapsto C(p, h), \quad \text{s.t.} \quad C(p, hh') = C(p, h) \cdot C(ph, h').$$

We see that a 1-cocycle generalises representations. From this defining property follows that

$$C(p, e_H) = e_G = C(ph, e_H), \quad (8)$$

$$C(p, h)^{-1} = C(ph, h^{-1}), \quad \text{and} \quad C(p, h^{-1}) = C(ph^{-1}, h)^{-1}.$$

The linearisation of such a group cocycle gives a 1-cocycle for the action of $\mathfrak{h}$ on P :

$$a : P \times \mathfrak{h} \rightarrow \mathfrak{g},$$

$$(p, X) \mapsto a(X, p), \quad \text{s.t.} \quad X^v \cdot a(Y, p) - Y^v \cdot a(X, p) \quad (9)$$

$$+ [a(X, p), a(Y, p)]_{\mathfrak{g}} = a([X, Y], p).$$

This can indeed be deduced from (7) via $a(X, p) := \left. \frac{d}{d\tau} C(p, h_\tau) \right|_{\tau=0}$ and $X := \left. \frac{d}{d\tau} h_\tau \right|_{\tau=0} \in \mathfrak{h}$. One may observe that, in the context of the field space of a gauge theory, $P \rightarrow \Phi$, this generalises the Wess–Zumino consistency condition for (classical and/or quantum) gauge anomalies (see refs. [7, 10]).

Now, for V a G -space, we define *cocyclic equivariant* forms as

$$\Omega_{\text{eq}}^*(P, C) := \{\alpha \in \Omega^*(P, V) \mid R_h^* \alpha = C(\cdot, h)^{-1} \alpha\}. \quad (10)$$

One may check that the 1-cocycle defining property ensures compatibility with the right action of H , $R_{hh'}^* = R_h^* R_{h'}^*$. The linear equivariance is then given by $\mathfrak{L}_{X^v} \alpha = -a(X, \cdot) \alpha$. The 1-cocycle property (9) ensures compatibility with $[\mathfrak{L}_{X^v}, \mathfrak{L}_{Y^v}] = \mathfrak{L}_{[X^v, Y^v]} = \mathfrak{L}_{[X, Y]^v}$ —which uses the fact that the map $\mathfrak{h} \rightarrow \Gamma(VP)$, $X \mapsto X^v$, is a Lie algebra morphism.

Forms that are both equivariant and horizontal are said *tensorial*. As above, *cocyclic tensorial* forms generalise standard tensorial forms:

$$\Omega_{\text{tens}}^*(P, \rho) := \Omega_{\text{hor}}^*(P) \cap \Omega_{\text{eq}}^*(P, \rho)$$

$$\subset \Omega_{\text{tens}}^*(P, C) := \Omega_{\text{hor}}^*(P) \cap \Omega_{\text{eq}}^*(P, C). \quad (11)$$

Observe that $\Omega_{\text{eq}}^0(P, \rho) = \Omega_{\text{tens}}^0(P, \rho)$ and $\Omega_{\text{eq}}^0(P, C) = \Omega_{\text{tens}}^0(P, C)$ since 0-forms are horizontal.

It is an elementary result of bundle geometry that there is a bijection $\Omega_{\text{tens}}^k(P, \rho) \simeq \Omega^k(M) \otimes E$ between the space of standard

tensorial forms and the space of forms on M valued in associated vector bundles $E = P \times_{\rho} V := P \times V / \sim$, where the equivalence relation is $[p, v] = (p, v) \sim (ph, \rho(h)^{-1}v)$: the bijection is given by $\tilde{\alpha}_{|x} = [p, \alpha_{|p}] = [ph, R_h^* \alpha_{|ph}] = [ph, \rho(h)^{-1} \alpha_{|p}]$. In degree 0, this is the well-known isomorphism $\Omega_{\text{tens}}^0(P, \rho) \simeq \Omega^0(M) \otimes E \simeq \Gamma(E)$ between representation-valued equivariant maps on P and sections of associated bundles $\Gamma(E) = \{s : M \rightarrow E\}$. This in particular gives another characterisation of the gauge group $\mathcal{H}$, whose elements are H -valued equivariant 0-forms, as the space of sections of the associated bundle $\text{Conj}(P) = P \times_{\text{Conj}} H = P \times H / \sim$, with equivalence relation given by $[p, h'] = (p, h') \sim (ph, h^{-1}h'h)$: so we have the isomorphisms $\text{Aut}_*(P) \simeq \mathcal{H} \simeq \Gamma(\text{Conj}(P))$.

The result extends to the bijection $\Omega_{\text{tens}}^k(P, C) \simeq \Omega^k(M) \otimes E^C$ between the space of cocyclic tensorial forms and the space of forms on M valued in *cocyclic associated bundles* $E^C = P \times_C V := P \times V / \sim$, where the equivalence relation is $[p, v] = (p, v) \sim (ph, C(p, h)^{-1}v)$: the bijection being $\tilde{\alpha}_{|x} = [p, \alpha_{|p}] = [ph, R_h^* \alpha_{|ph}] = [ph, C(p, h)^{-1} \alpha_{|p}]$. In degree 0, this gives the isomorphism $\Omega_{\text{tens}}^0(P, C) \simeq \Gamma(E^C)$, with $\Gamma(E^C) = \{\tilde{s} : M \rightarrow E^C\}$.

Finally, one may define the important space of *basic* forms

$$\Omega_{\text{basic}}^*(P) := \{\alpha \in \Omega^*(P) \mid \exists \beta \in \Omega^*(M) \text{ s.t. } \alpha = \pi^* \beta\}. \quad (12)$$

These are horizontal: $\alpha(X^v) = \pi^*(\beta)(X^v) = \beta(\pi_* X^v) = 0$. They are *invariant*, i.e., have trivial equivariance: $R_h^* \alpha = R_h^* \pi^* \beta = (\pi \circ R_h)^* \beta = \pi^* \beta =: \alpha$. This makes for an alternative definition of basic forms, as follows: $\Omega_{\text{basic}}^*(P) := \{\alpha \in \Omega^*(P) \mid \iota_{X^v} \alpha = 0 \text{ and } R_h^* \alpha = \alpha\} = \Omega_{\text{hor}}^*(P) \cap \Omega_{\text{inv}}^*(P)$.

Connections and Covariant Derivatives: The exterior derivative is not a derivation of the spaces of tensorial forms. To define first-order differential operators preserving those spaces, the *covariant derivatives*, we need non-canonical structures on P : the appropriate notions of connection.

In standard bundle geometry, the solution is well-known and given by the celebrated Ehresmann (or principal) connection 1-form $\omega \in \Omega^1(P, \mathfrak{h})$ defined as

$$R_h^* \omega = \text{Ad}_{h^{-1}} \omega, \quad \text{i.e., } \omega \in \Omega_{\text{eq}}^1(P, \text{Ad}), \quad \text{and}$$

$$\omega_p(X_p^v) = X \in \mathfrak{h}, \quad \forall X_p^v \in V_p P, \quad (13)$$

whose associated curvature can be found via the Cartan structure equation $\Omega = d\omega + \frac{1}{2}[\omega, \omega] \in \Omega_{\text{tens}}^2(P, \text{Ad})$. The connection ω allows to define the standard covariant derivative $D = d + \rho_*(\omega) : \Omega_{\text{tens}}^*(P, \rho) \rightarrow \Omega_{\text{tens}}^{*+1}(P, \rho)$, which is s.t. $D \circ D = \rho_*(\Omega)$. Furthermore, we have the Bianchi identity: $D\Omega = 0$.

By $\Omega_{\text{tens}}^k(P, \rho) \simeq \Omega^k(M) \otimes E$, a connection ω on P induces a connection on the space $\Omega^*(M) \otimes E$ by

$$D : \Omega_{\text{tens}}^k(P, \rho) \rightarrow \Omega_{\text{tens}}^{k+1}(P, \rho), \quad \Rightarrow \quad \nabla : \Omega^k(M) \otimes E \rightarrow \Omega^{k+1}(M) \otimes E,$$

$$\alpha \mapsto D\alpha, \quad \tilde{\alpha} \mapsto \nabla \tilde{\alpha} = [_, D\alpha].$$

In particular, for $k = 0$ we have $\varphi \in \Omega_{\text{tens}}^0(P, \rho) \sim s \in \Gamma(E)$, and the above restricts to a connection on the space of sections of associated bundles:

$$\nabla : \Gamma(E) \rightarrow \Omega^1(M) \otimes E, \quad (14)$$

$$s \mapsto \nabla s = [_, D\varphi].$$

Finally, we observe that the space of connections C is an affine space modelled on the vector space $\Omega_{\text{tens}}^1(P, \mathfrak{h})$, meaning that for $\omega \in C$ and $\alpha \in \Omega_{\text{tens}}^1(P, \mathfrak{h})$, $\omega' := \omega + \alpha \in C$.⁴

This generalises to cocyclic bundle geometry. One first defines the notion of *cocyclic connection* 1-form $\varpi \in \Omega^1(P, \mathfrak{g})$ by

$$R_h^* \varpi|_{ph} = \text{Ad}_{C(p,h)^{-1}} \varpi|_p + C(p, h)^{-1} dC(\cdot, h)|_p, \quad (15)$$

$$\varpi_p(X_p^\nu) = \frac{d}{d\tau} C(p, h_\tau) \Big|_{\tau=0} := a(X, p) \in \mathfrak{g},$$

where $X := \frac{d}{d\tau} h_\tau \Big|_{\tau=0} \in \mathfrak{h}$. The *cocyclic curvature* is a cocyclic tensorial 2-form defined by the Cartan structure equation $\bar{\Omega} := d\varpi + \frac{1}{2}[\varpi, \varpi] \in \Omega_{\text{tens}}^2(P, C)$. A cocyclic connection allows to define a *cocyclic covariant derivative* $\bar{D} = d + \varpi : \Omega_{\text{tens}}^*(P, C) \rightarrow \Omega_{\text{tens}}^{*+1}(P, C)$, which is s.t. $\bar{D} \circ \bar{D} = \bar{\Omega}$. The cocyclic curvature satisfies a Bianchi identity, $\bar{D}\bar{\Omega} = 0$.

By the isomorphism $\Omega_{\text{tens}}^k(P, C) \simeq \Omega^k(M) \otimes E^C$, a cocyclic connection ϖ on P induces a cocyclic connection on the space $\Omega^*(M) \otimes E^C$ by

$$\bar{D} : \Omega_{\text{tens}}^k(P, C) \rightarrow \Omega_{\text{tens}}^{k+1}(P, C), \Rightarrow \bar{V} : \Omega^k(M) \otimes E^C \rightarrow \Omega^{k+1}(M) \otimes E^C,$$

$$\alpha \mapsto \bar{D}\alpha, \quad \tilde{\alpha} \mapsto \bar{V}\tilde{\alpha} = [_, \bar{D}\alpha].$$

For $k = 0$, we have $\varphi \in \Omega_{\text{tens}}^0(P, C) \sim \tilde{\varphi} \in \Gamma(E^C)$ and the above restricts to a connection on the space of sections of cocyclic associated bundles:

$$\bar{V} : \Gamma(E^C) \rightarrow \Omega^1(M) \otimes E^C, \quad (16)$$

$$\tilde{\varphi} \mapsto \bar{V}\tilde{\varphi} = [_, \bar{D}\varphi].$$

The space $\bar{C}$ of cocyclic connections is an affine space modeled on cocyclic tensorial 1-forms $\Omega_{\text{tens}}^1(P, C)$, so that for $\varpi \in \bar{C}$ and $\alpha \in \Omega_{\text{tens}}^1(P, C)$, $\varpi' = \varpi + \alpha$.

We observe that the exterior derivative d is a covariant derivative for basic forms, $d : \Omega_{\text{basic}}^*(P) \rightarrow \Omega_{\text{basic}}^{*+1}(P)$. This means that $(\Omega_{\text{basic}}^*(P), d)$ is a subcomplex of the de Rham complex of P , called the *basic subcomplex*, defining the *equivariant cohomology* of P .^[23] It is of special importance, notably in Physics, since it is isomorphic to the de Rham complex of the base M : $(\Omega_{\text{basic}}^*(P), d) \simeq (\Omega^*(M), d)$. As we shall see shortly, the dressing field method (DFM) is a way to associate to a form on P a corresponding basic form.

Gauge Transformations: As observed above, the group of automorphisms $\text{Aut}(P)$ acts on $\Omega^*(P)$ by pullback: it preserves the various spaces defined above, in particular tensorial forms $\Omega_{\text{tens}}^*(P)$ and the space of connections C . This stems from the functoriality of the definition of those spaces, automorphisms being special cases of morphisms in the bundle category. It follows that *gauge transformations*, defined by the action of vertical automorphisms $\text{Aut}_v(P)$, preserve these spaces too. Since $\text{Aut}_v(P) \simeq \mathcal{H}$, gauge transformations may be expressed via elements of $\mathcal{H}$: We define the gauge transformation of $\alpha \in \Omega^*(P)$ by $\alpha'(\mathfrak{X}) := \Xi^* \alpha(\mathfrak{X})$, for

$\gamma \in \mathcal{H} \sim \Xi \in \text{Aut}_v(P)$ and $\mathfrak{X} \in \Gamma(TP)$. The explicit result is found by $\Xi^* \alpha(\mathfrak{X}) = \alpha(\Xi_* \mathfrak{X})$ and via the well-known result

$$\Xi_* \mathfrak{X}|_p = R_{\gamma(p)^*} \mathfrak{X}|_p + \{\gamma^{-1} d\gamma|_p(\mathfrak{X}|_p)\}_{|\Xi(p)}^\vee$$

$$= R_{\gamma(p)^*} \left(\mathfrak{X}|_p + \{d\gamma \gamma^{-1}(\mathfrak{X}|_p)\}_{|\Xi(p)}^\vee \right). \quad (17)$$

From this follows that standard and cocyclic tensorial forms have homogeneous gauge transformations, controlled by their equivariance: they are “gauge tensorial,” hence their name,

$$\alpha' = \rho(\gamma)^{-1} \alpha, \quad \text{for } \alpha \in \Omega_{\text{tens}}^*(P, \rho),$$

$$\alpha' = C(\gamma)^{-1} \alpha, \quad \text{for } \alpha \in \Omega_{\text{tens}}^*(P, C), \quad (18)$$

where we introduce the shorthand notation $C(\gamma(p)) := C(p, \gamma(p))$. The $\mathcal{H}$ -transformations of standard and cocyclic connections are

$$\omega' := \Xi^* \omega = \gamma^{-1} \omega \gamma + \gamma^{-1} d\gamma, \quad \text{for } \omega \in C, \quad (19)$$

$$\varpi' := \Xi^* \varpi = C(\gamma)^{-1} \varpi C(\gamma) + C(\gamma)^{-1} dC(\gamma), \quad \text{for } \varpi \in \bar{C}.$$

The first result is well-known, and easily read from (17). For the proof of the second one see Section 4.2 of [10]. The result (18) gives, in particular, the $\mathcal{H}$ -transformations of the curvatures Ω and $\bar{\Omega}$ of the Ehresmann and cocyclic connections, as well as that of the (cocyclic) covariant derivative $\bar{D}\alpha$ of a (cocyclic) form α . It also gives the action of the gauge group $\mathcal{H}$ on its own elements: for $\eta, \gamma \in \mathcal{H}$ we have $\eta' = \gamma^{-1} \eta \gamma$.

The action of $\text{Aut}_v(P) \simeq \mathcal{H}$, giving (18)–(19), are called *active* gauge transformations. These are to be distinguished from *passive* gauge transformations, the name given to the *gluings* of local representatives on the base M , stemming from the local structure of P .

Local Structure: A principal bundle P is locally trivial, i.e. for any open sets $U \subset M$ we have $P|_U \simeq U \times H$: the isomorphism being realised by a *trivialisation* $\phi = (\pi, t) : P|_U \rightarrow U \times H$, $p \mapsto \phi(p) = (\pi(p) = x, t(p))$, which provides a “bundle coordinate chart” on $P|_U$. A trivialisation is compatible with the structure group action, $\phi \circ R_h = R_h \circ \phi$, meaning that the fiber coordinate map t is s.t. $R_h^* t = t h$, or $t(ph) = t(p)h$. Often t itself is called a trivialisation. A local section of P is a map $\sigma : U \rightarrow P|_U$, $x \mapsto \sigma(x)$ —which allows to define a trivialisation of $P|_U$. Any two local sections σ and σ' over U —or over overlapping sets $U \cap U'$ —are related via $\sigma' = \sigma g$, where g is a transition function of P , that is a map $g : U \rightarrow H$, $x \mapsto g(x)$.

Given a local section σ , one may define the *local representative* of a form $\beta \in \Omega^*(P)$ as $b := \sigma^* \beta \in \Omega^*(U)$. In particular, the local representative of a standard or cocyclic tensorial form is $a := \sigma^* \alpha \in \Omega^*(U, V)$, that of Ehresmann and cocyclic connections are respectively $A := \sigma^* \omega \in \Omega^*(U, \mathfrak{h})$ and $\bar{A} := \sigma^* \varpi \in \Omega^*(U, \mathfrak{g})$. One may also define the pullback of a 1-cocycle $C := \sigma^* C : U \times H \rightarrow G$, $(x, h) \mapsto C(x, h) = C(\sigma(x), h)$. It allows to define the map $C(g) := C(\sigma, g) : U \rightarrow G$, which are *cocyclic transition function*—i.e., none other than the transition functions of cocyclic associated bundles E^C . Under change of local section $\sigma' = \sigma g$, i.e., of bundle

⁴ Or, as is clear from the above defining properties, for $\omega, \omega' \in C$, $\omega + \omega' \notin C$ and $\omega' - \omega \in \Omega_{\text{tens}}^1(P, \mathfrak{h})$.

coordinates, the local representatives $a' := \sigma'^* a$, $A' := \sigma'^* \omega$ and $\bar{A}' := \sigma'^* \varpi$ satisfy *gluing relations*

$$\begin{aligned} a' &= \rho(g)^{-1} a, & A' &= g^{-1} A g + g^{-1} d g, \\ a' &= C(g)^{-1} a, & \bar{A}' &= C(g)^{-1} \bar{A} C(g) + C(g)^{-1} d C(g). \end{aligned} \quad (20)$$

These are also known as *passive gauge transformations*. These must be conceptually distinguished from *local active gauge transformations*: Let us define the local gauge group $\mathcal{H}_{\text{loc}} := \{\gamma := \sigma^* \gamma : U \rightarrow H \mid \gamma \in H, \text{ and } \eta' = \gamma^{-1} \eta \gamma\}$, and consider the map $C(\gamma) := C(\sigma, \gamma) : U \rightarrow G$. Now, through a given section σ , the local representatives of the active gauge transformations (18)–(19) are

$$\begin{aligned} a' &= \rho(\gamma)^{-1} a, & A' &= \gamma^{-1} A \gamma + \gamma^{-1} d \gamma, \\ a' &= C(\gamma)^{-1} a, & \bar{A}' &= C(\gamma)^{-1} \bar{A} C(\gamma) + C(\gamma)^{-1} d C(\gamma). \end{aligned} \quad (21)$$

These are the local active gauge transformations, formally indistinguishable from the passive gauge transformations (20), but conceptually as distinct as coordinate changes and diffeomorphisms in GR.

2.2. The Dressing Field Method: Basics

The dressing field method (DFM) has been conceived as a tool of gauge symmetry reduction in Gauge Field Theory^[16,24,25] (see also refs. [26, 27]) whose foundational implications were first explored in refs. [28, 29] (see also ref. [30]). It is extended to general-relativistic theories, whose covariance group is the group of diffeomorphisms, in ref. [17]. The most general and synthetic mathematical and conceptual presentation, for general-relativistic Gauge Field Theory, is to be found in ref. [7].

Mathematically, it is a result about realisation of subbundles and of basic, or partially basic, forms on a bundle. In that respect it is close to the well-known bundle reduction theorem, a standard result of bundle geometry.^[31–33] A key difference is that, in the DFM the dressing map, or *dressing field*, is assumed to have a functional dependence on one or several forms of P . In physical terms, the dressing field is a functional of the degrees of freedom (d.o.f.) of the theory under consideration. The DFM thus has a straightforward *relational* interpretation:⁵ the basic forms produced, or dressed fields, are naturally seen as relational variables as articulated precisely in refs. [6, 7, 17]—and close to the views expressed, e.g., in refs. [35–39].

We shall now give a synthetic review of the fundamentals of the DFM, as it is essential to our subsequent relational reformulation of non-relativistic classical and quantum mechanics.

Dressing Field and Basic Dressed Fields: Suppose there exists a subgroup $K \subseteq H$ of the structure group, to which corresponds a subgroup $\mathcal{K} \subset \mathcal{H}$ of the gauge group, which is isomorphic to the subgroup $\text{Aut}_v(P)^K \subset \text{Aut}_v(P)$ of the group of vertical automorphisms. Consider further a Lie group G s.t. $K \subseteq G \subseteq H$. A K -

dressing field is defined as

$$\begin{aligned} u : P \rightarrow G, \quad \text{s.t.} \quad u^k &:= \Xi^* u = \kappa^{-1} u, \quad \Xi \in \text{Aut}_v(P)^K \sim \kappa \in \mathcal{K}, \\ \text{or s.t.} \quad R_k^* u &:= k^{-1} u, \quad k \in K. \end{aligned} \quad (22)$$

We denote the space of G -valued K -dressing fields on P by $\text{Dr}[K, G]$: K is called the equivariance group of the dressing field and G its target group. In the DFM, one considers more generally dressing fields having a functional dependence on objects of P (typically forms):

$$\begin{aligned} u : \Omega^*(P) &\rightarrow \text{Dr}[K, G], \\ \alpha &\mapsto \{u[\alpha] : P \rightarrow G, \quad \text{s.t.} \quad u[\alpha]^k := \Xi^* u[\alpha] = \kappa^{-1} u[\alpha], \\ &\quad \text{or s.t.} \quad R_k^* u[\alpha] := k^{-1} u[\alpha], \quad k \in K\}. \end{aligned} \quad (23)$$

Alternatively, the property can be defined as—or required compatible with—an equivariance w.r.t. transformation of the functional argument: $u[\alpha]^k := u[\alpha^k] = u[\Xi^* \alpha] = \kappa^{-1} u[\alpha]$. In physical applications, dynamical d.o.f. are represented by (or embedded into) objects on P , so one may understand dressing fields (23) as being determined by physical (sub)systems. We may also observe that for $K = G = H$ a dressing field (22) is equivalent to a trivialisation $t := u^{-1}$, i.e., provides a bundle coordinate chart. Therefore, dressing fields (23) provide bundle coordinatisations via physical degrees of freedom.

Given such a K -dressing field, one may define the dressing map

$$\begin{aligned} f_u : P \rightarrow P' \subset P, \quad \text{s.t.} \quad f_u \circ R_k &= f_u, \quad \text{and} \quad f_u \circ \Xi = f_u \\ &\quad \text{for} \quad \Xi \in \text{Aut}_v(P)^K \sim \mathcal{K}, \\ p &\mapsto f_u(p) := pu(p), \end{aligned} \quad (24)$$

where $P' = P/K$ is a quotient space since f_u is constant along the action of K . It allows to define, for any form $\beta \in \Omega^*(P)$, its *dressed counterpart*

$$\beta^u := f_u^* \beta \in \Omega_{K\text{-basic}}^*(P). \quad (25)$$

It is K -basic since it has both trivial K -equivariance and is K -horizontal, so that it is $\mathcal{K} \simeq \text{Aut}_v(P)^K$ -invariant, $(\beta^u)^k := \Xi^* (\beta^u) = \Xi^* f_u^* \beta = f_u^* \beta =: \beta^u$. A dressed form β^u therefore naturally “lives” on the quotient space P/K . In particular, the dressings of an Ehresmann connection ω and standard tensorial form α have explicit expressions

$$\omega^u := u^{-1} \omega u + u^{-1} du \quad \text{and} \quad \alpha^u := \rho(u)^{-1} \alpha. \quad (26)$$

The dressing of cocyclic connections and cocyclic tensorial forms is

$$\varpi^u := C(u)^{-1} \varpi C(u) + C(u)^{-1} d C(u) \quad \text{and} \quad \alpha^u := C(u)^{-1} \alpha, \quad (27)$$

where, from the 1-cocycle $C : P \times H \rightarrow G$ (where we distinguish the target group G of C from that $G \subseteq H$ of the dressing field),

⁵ Which sets it apart from gauge-fixing, a notion, in Gauge Field Theory, with which it is often conflated. See ref. [34]

one defines the K -cocyclic dressing field

$$C(\mathbf{u}) := C(\cdot, \mathbf{u}) : P \rightarrow G, \quad \text{s.t.} \quad R_k^* C(\mathbf{u}) = C(\cdot, k)^{-1} C(\mathbf{u}). \quad (28)$$

$$p \mapsto C(\mathbf{u}(p)) := C(p, \mathbf{u}(p)),$$

whose equivariance is secured by the 1-cocycle property (7) and (8):

$$C(\mathbf{u}(pk)) = C(pk, \mathbf{u}(pk)) = C(pk, k^{-1}\mathbf{u}(p))$$

$$= C(pk, k^{-1}) C(pk k^{-1}, \mathbf{u}(p)) = C(p, k)^{-1} C(p, \mathbf{u}(p)). \quad (29)$$

Similarly, we find the $C(\mathbf{u})^\kappa = C(\kappa)^{-1} C(\mathbf{u})$, from which, together with (18)–(19), the $\mathcal{K} \simeq \text{Aut}_v(P)^K$ -invariance of (27) is easily checked.

From the above, comparing (18)–(19) and (26)–(27), we derive the “rule of thumb” of the DFM: To find the dressing of a form $\beta \in \Omega^*(P)$, on account of the formal similarity of the action of the dressing map f_u and a vertical automorphism $\text{Aut}_v(P)^K$, first compute its $\mathcal{K}$ -gauge transformation $\beta^\kappa := \Xi^* \beta$ in terms of $\kappa \in \mathcal{K} \sim \Xi \in \text{Aut}_v(P)^K$, and then simply substitute $\kappa \rightarrow \mathbf{u}$ in this expression to get the dressed, K -basic, form β^u .

Let us stress the following important fact: It should be clear from its definition that a dressing field is not a gauge group element, $\mathbf{u} \notin \mathcal{K}$, and that a dressing map f_u is not a vertical automorphism. So that despite a superficial formal resemblance with (18)–(19), dressed fields (26)–(27) are not gauge transformations: dressed fields β^u are not points in the $\mathcal{K}$ -gauge orbits $\mathcal{O}_\kappa[\beta]$ of the form β —and in particular must not be conflated with a gauge-fixing of β . For example, a dressed connection is no more a connection, $\omega^u \notin C$.

We observe that a dressing field induces a (partial, i.e., LieK-valued) flat Ehresmann connection $\omega_0 := -d\mathbf{u}\mathbf{u}^{-1}$.

As we stressed above, since dressings fields (23) depend on the dynamical d.o.f. at play, dressed forms (25)–(27) have a natural interpretation as *relational variables*: They encode the $\mathcal{K}$ -gauge-invariant, physical, relational d.o.f. among the dynamical d.o.f. describing the system.

Residual Transformations of the 1st Kind: Being K -basic, i.e., $\mathcal{K}$ -invariant, the dressed forms β^u may be expected to display residual transformations under what remains of the structure and gauge groups. For these to be meaningful, it must be the case that K is a normal subgroup of the structure group, $K \triangleleft H$, so that their quotient is itself a group $H/K = J$. Then, $P' = P/K$ is a J -principal subbundle of P , on which dressed form β^u “live,” with isomorphic vertical automorphism and gauge groups $\text{Aut}(P') = \text{Aut}_v(P)^J \simeq J$. The structure group of P can then be written (assuming the trivial homomorphism $\phi : J \rightarrow H$, $\phi(j) = j \in H$) as $H = J \ltimes K$, correspondingly its vertical automorphism groups is $\text{Aut}_v(P) = \text{Aut}_v(P)^J \ltimes \text{Aut}_v(P)^K$, and its gauge group is $\mathcal{H} = J \ltimes \mathcal{K}$.

Since the J -equivariance and J -horizontality of β are already known by assumption, those of β^u depend only on the J -equivariance of $\mathbf{u}$: $R_j^* \mathbf{u}$. Alternatively, one may say that the J -transformation of β being known, to get that of β^u one needs only to find the J -transformation of $\mathbf{u}$: $\mathbf{u}^j$. Finding the residual J -equivariance and/or J -transformation of $\mathbf{u}$ will depend on the specifics of the situation under consideration.

As an illustrative example, consider the case where $R_j^* \mathbf{u} = j^{-1} \mathbf{u} j$ for $j \in J$, so that $\mathbf{u}^j = \eta^{-1} \mathbf{u} \eta$ for $\eta \in J$. The residual J -transformations of the dressed forms (26) are then easily found to be: $(\omega^u)^j = \eta^{-1} \omega^u \eta + \eta^{-1} d\eta$ and $(\alpha^u)^j = \rho(\eta)^{-1} \alpha^u$. They behave as standard connections and tensorial forms on $P' \subset P$. See refs. [12, 29] for other cases.

Residual Transformations of the 2nd Kind: The dressed forms may exhibit residual transformations resulting from a possible ambiguity in the choice of dressing field. Two dressings $\mathbf{u}, \mathbf{u}' \in \text{Dr}[K, G]$ may a priori be related by $\mathbf{u}' = \mathbf{u}^\xi$, where $\xi \in \mathcal{G} := \{ \xi : P \rightarrow G \mid R_k^* \xi = \xi \}$. We may write the $\mathcal{G}$ -action on a dressing by $\mathbf{u}^\xi = \mathbf{u}^\xi$. By definition, $\mathcal{G}$ has no action on $\Omega^*(P)$, so $\beta^\xi = \beta$. The $\mathcal{G}$ -action on dressed forms is thus easily found via $(\beta^u)^\xi := (\beta^\xi)^{u^\xi} = \beta^{u^\xi}$. For example, the residual $\mathcal{G}$ -transformations of (26) are found to be: $(\omega^u)^\xi := (\omega^\xi)^{u^\xi} = \omega^{u^\xi} = \xi^{-1} \omega^u \xi + \xi^{-1} d\xi$ and $(\alpha^u)^\xi := (\alpha^\xi)^{u^\xi} = \alpha^{u^\xi} = \rho(\xi^{-1}) \alpha^u$.

For cocyclic dressing fields we get $C(\mathbf{u})^\xi := C(\mathbf{u}^\xi)$, which is $C(p, \mathbf{u}(p)\xi(p)) = C(p, \mathbf{u}(p)) C(p\mathbf{u}(p), \xi(p)) = C(p, \mathbf{u}(p)) C(f_u(p), \xi(p))$, where we may write $f_u(p) = p' \in P' \subset P$. This result we may denote by

$$C(\mathbf{u})^\xi = C(\mathbf{u}^\xi) = C(\mathbf{u}) C(\xi). \quad (30)$$

So, the residual $\mathcal{G}$ -transformations of cocyclic dressed forms (27) are easily found to be:

$$(\omega^u)^\xi = C(\xi)^{-1} \omega^u C(\xi) + C(\xi)^{-1} dC(\xi) \quad \text{and} \quad (\alpha^u)^\xi = C(\xi)^{-1} \alpha^u. \quad (31)$$

Transformations like (31) are the basis of the frame change covariance of the relational version of our geometric formulation of QM, both of which we now describe.

3. Bundle Geometric Non-Relativistic Mechanics and Quantum Mechanics

As said in the introduction, until now, the main applications of the DFM have been in classical general-relativistic Gauge Field Theory,^[7] the natural relational interpretation of the former fitting the conceptual foundation of the latter.^[6] In the following, the DFM will be applied for the first time in a quantum context, allowing a manifestly relational formulation of non-relativistic many-particle quantum mechanics. This is made possible by first providing a bundle geometric description of both non-relativistic Mechanics and then Quantum Mechanics.

3.1. Bundle Geometry of Configuration Space-Time $\mathcal{E}$

Our geometric formulation of non-relativistic QM is based on the bundle geometry of the configuration space-time of N classical structureless, point-like, particles, which generalises the description of Galilean space-time.

The configuration space-time is a principal bundle $\mathcal{E} \xrightarrow{\pi} \mathcal{T}$ over the 1-dimensional non-compact manifold $\mathcal{T}$, representing the Newtonian universal time line. The fiber over an instant $t \in \mathcal{T}$ is $\mathcal{E}_t := \pi^{-1}(t) = \mathbb{E}^{3N}$, i.e., the configuration space of N classical

structureless point particles. A point $p = (p_1, \dots, p_N) \in \mathcal{C}_t \subset \mathcal{C}$, $\pi(p) = t \in \mathcal{T}$, is the instantaneous configuration of the N particles. The structure group of $\mathcal{C}$ is $H = \mathbb{R}^{3N} := \underbrace{\mathbb{R}^3 \times \dots \times \mathbb{R}^3}_{N \text{ times}}$.

Its right action is

$$\begin{aligned} \mathcal{C} \times H &\rightarrow \mathcal{C}, \\ (p, X) &\mapsto R_X p = p + X, \quad \text{s.t.} \quad \pi \circ R_X = \pi. \end{aligned} \quad (32)$$

We observe that this action reflects the fact that $\mathbb{E}^{3N}$ is an affine space over $\mathbb{R}^{3N}$, because the Euclidean plane $\mathbb{E}^3$ is affine over $\mathbb{R}^3$. Since H is (additive) Abelian, we have $R_{X'} R_X p = p + X + X' = R_{X+X'} p$, i.e., $R_{X+X'} = R_{X'} R_X$.

As usual, the linearisation of the right action of H defines the canonical vertical subbundle $V\mathcal{C} \subset T\mathcal{C}$, and a fundamental vertical vector is

$$\chi_p^\nu := \left. \frac{d}{d\tau} R_{X_\tau} p \right|_{\tau=0} = p + \chi \in V_p \mathcal{C}, \quad \text{with} \quad \left. \frac{d}{d\tau} X_\tau \right|_{\tau=0} := \chi \in \mathfrak{h}, \quad (33)$$

so that a fundamental vector field is $\chi^\nu \in \Gamma(V\mathcal{C})$. Because H is Abelian, these are right invariant: $R_{X*} \chi_p^\nu = \chi_{p+X}^\nu$. Equation (33) defines the “verticality map” $|\cdot|^\nu : \mathfrak{h} \rightarrow \Gamma(V\mathcal{C})$, applying to any $\mathfrak{h}$ -valued (0-)form on $\mathcal{C}$. As we shall see shortly, contrary to the case of Gauge Field Theory, vertical “motion” along the fibers of $\mathcal{C}$, with “velocity” in $\Gamma(V\mathcal{C})$, is not always unphysical, or “pure gauge”: it also contains genuine physical changes within the set of particles. Of course, the true *physical* dynamics of the latter unfolds along a curve in $\mathcal{C}$ that projects onto a curve in $\mathcal{T}$. Over an open set $\mathcal{U} \subset \mathcal{T}$, i.e., a time interval, a local trivialisation is a map

$$\begin{aligned} \phi : \mathcal{C}|_{\mathcal{U}} &\rightarrow \mathcal{U} \times H, \\ p &\mapsto \phi(p) := (\pi(p), h(p)) = (t, x) = (t, (x_1, \dots, x_N)). \end{aligned} \quad (34)$$

The fiber coordinate $h(p) = x$ may also be written $x = \{x_i\}_{i \in \{1, \dots, N\}} = \{x_i^a e_a\}$ where $\{e_a\}$ is a basis of $\mathbb{R}^3$, meaning that $x = \{x_i\}$ is an intrinsic vector of $\mathbb{R}^{3N}$, independent of a choice of basis. The trivialisation is s.t. $\phi \circ R_X = R_X \circ \phi$, i.e., $\phi(p + X) = \phi(p) + X$, meaning that the fiber coordinate satisfies $R_X^* h = h + X$, i.e. $h(p + X) = h(p) + X$. A local (trivialising) section is $\sigma : \mathcal{U} \subset \mathcal{T} \rightarrow \mathcal{C}|_{\mathcal{U}} \simeq \mathcal{U} \times H$, $t \mapsto \sigma(t) = (t, x(t))$, i.e., it gives the graph of the trajectories of the particles described by $p = \sigma(t)$. Two local sections are related by $\sigma' = R_g \sigma = \sigma + g$, where $g : \mathcal{U} \rightarrow H$, $t \mapsto g(t)$ is a transition function of $\mathcal{C}$. These are involved in *gluings* of local representatives on $\mathcal{T}$ of forms on $\mathcal{C}$, i.e., in *passive gauge transformations*. Again, beware that here the latter may relate two physically distinct particle configurations. Since $\mathcal{T}$ is contractible, $\mathcal{C}$ is *globally trivial*, $\mathcal{C} \simeq \mathcal{T} \times H$.

Under the action $R_X : \mathcal{C}|_t \rightarrow \mathcal{C}|_t$, $p = (p_1, \dots, p_N) \mapsto p + X = (p_1 + X_1, \dots, p_N + X_N)$, of the structure group $H = \mathbb{R}^{3N}$, each particle position is shifted by a different vector $X_i \in \mathbb{R}^3$. It is clear that any configuration of particles is left invariant by the




Figure 1. The 2-fibration structure of the configuration space-time bundle $\mathcal{C}$ of N classical particles.

subgroup

$$H_\Delta = \mathbb{R}^{3N}_\Delta := \text{diag} \mathbb{R}^{3N} \simeq \mathbb{R}^3, \quad \text{whose elements are } X = \underbrace{(X, \dots, X)}_{N \text{ times}}. \quad (35)$$

Now, since any subgroup of an Abelian group is *normal*, we have $H_\Delta \triangleleft H$, so that H/H_Δ is an Abelian group. The structure group thus decomposes as

$$H = H/H_\Delta \ltimes H_\Delta =: H_{\text{int}} \ltimes H_{\text{ext}}, \quad (36)$$

where the nomenclature is chosen to reflect the fact that H_{int} implements physical changes in the configuration of particles as seen from any particle within it, while H_{ext} only reflects a global change of the whole configuration as seen from a viewpoint external to the configuration.⁶

The action of H_{int} maps a configuration to a physically distinct configuration, where the relative positions of the particles are different. The decomposition (36) implies that the configuration space-time bundle is a 2-fibration:

$$\begin{aligned} \mathcal{C} &\xrightarrow{H_{\text{ext}}} \mathcal{C}_{\text{rel}} \xrightarrow{H_{\text{int}}} \mathcal{T}, \\ \text{or} \quad \mathcal{C} &\xrightarrow{\pi_1} \mathcal{C}_{\text{rel}} := \mathcal{C}/H_{\text{ext}} \xrightarrow{\pi_2} \mathcal{T}, \end{aligned} \quad (37)$$

$$\text{with} \quad \pi_2 \circ \pi_1 = \pi,$$

where the quotient subbundle $\mathcal{C}_{\text{rel}} := \mathcal{C}/H_{\text{ext}} \xrightarrow{H_{\text{int}}} \mathcal{T}$ is the *relational configuration space-time bundle*—here realised as a quotient—encoding the internal relative dynamics of the N particles system. Figure 1 sketches the 2-fibration structure of $\mathcal{C}$.

⁶ So, $H_{\text{ext}} = H_\Delta$ can be seen as inducing unphysical transformation of the configuration. Yet, it is not so for the corresponding gauge subgroup, H_{ext} . See below.

The group of automorphism of $\mathcal{C}$ is $\text{Aut}(\mathcal{C}) := \{\Xi \in \text{Diff}(\mathcal{C}) \mid \Xi \circ R_X = R_X \circ \Xi\}$, and projects onto $\text{Diff}(\mathcal{T})$, i.e., induces diffeomorphisms of the time line. Its maximal normal subgroup is the group of vertical automorphisms $\text{Aut}_v(\mathcal{C}) := \{\Xi \in \text{Aut}(\mathcal{C}) \mid \pi \circ \Xi = \pi\}$, inducing the identity transformation on $\mathcal{T}$.

The latter is isomorphic to the gauge group $\mathcal{H} := \{X : \mathcal{C} \rightarrow H \mid R_X^* X = \text{Conj}(-X)X = X\}$. The isomorphism being $\Xi(p) = R_{X(p)}p = p + X(p)$. It is noteworthy that the elements of the gauge group are *basic*, being horizontal as usual (as 0-forms on $\mathcal{C}$) and *invariant* due to H being Abelian. So, we have that $\mathcal{H} \simeq \Omega^0_{\text{basic}}(\mathcal{C}, H) \simeq \Omega^0(\mathcal{T}, H)$, i.e., any $X \in \mathcal{H}$ arises from some $\tilde{X} : \mathcal{T} \rightarrow H$ s.t. $X = \pi^* \tilde{X}$. Elements of the gauge group $\mathcal{H}$ thus depend only on $t \in \mathcal{T}$. The restriction of $\mathcal{H}$ to a trivial patch $\mathcal{C}_{|U} \subset \mathcal{C}$ thus carries the same information as the local gauge group $\mathcal{H}_{\text{loc}} := \{\tilde{X} : \sigma^* X : U \rightarrow H \mid \tilde{X}^{\tilde{X}} = \tilde{X}\} \simeq \Omega^0(U, H) =: \Omega^0(\mathcal{T}, H)_{|U}$. The definition of $\mathcal{H}_{\text{loc}}$ is actually independent of the choice of local section σ , its elements having trivial gluings. The use of the same notation $\tilde{X}$ for elements of $\mathcal{H}_{\text{loc}}$ and $\Omega^0(\mathcal{T}, H)$ is meant to highlight that they are the same objects.

We observe that $\mathcal{H}$ includes *Galilean boosts* via elements of the form $X(t) = v_{\text{ext}} t$, where $v_{\text{ext}} \in H_{\text{ext}}$ is constant and interpretable as the relative velocity between the configuration p of N particles and an external observer $\mathcal{O}_{\text{ext}}$. The gauge group also encodes constant relative accelerations $a_{\text{ext}} \in H_{\text{ext}}$ between p and $\mathcal{O}_{\text{ext}}$ via $X(t) = \frac{1}{2} a_{\text{ext}} t^2$, whereby inertial forces manifest. More generally, the gauge group $\mathcal{H} \simeq \Omega^0_{\text{basic}}(\mathcal{C}, H) \simeq \Omega^0(\mathcal{T}, H)$ encodes precisely what is known as “*extended Galilean transformations*”: Indeed, the action $\Xi : \mathcal{C} \rightarrow \mathcal{C}, p \mapsto \Xi(p) = p + X(p)$, of a vertical automorphism $\Xi \in \text{Aut}_v(\mathcal{C}) \simeq X \in \mathcal{H}$ is, in bundle coordinates, $(t, x) \mapsto (t', x') = (t, x + \tilde{X}(t))$. The latter is precisely the usual definition of extended Galilean transformations as described, e.g., in ref. [22]. The characteristic SES of the bundle $\mathcal{C}$ is then

$$\text{id}_{\mathcal{C}} \rightarrow \text{Aut}_v(\mathcal{C}) \simeq \mathcal{H} \xrightarrow{\Delta} \text{Aut}(\mathcal{C}) \rightarrow \text{Diff}(\mathcal{T}) \rightarrow \text{id}_{\mathcal{T}}. \quad (38)$$

Given that the structure group H decomposes as (36), the group of vertical automorphisms and the gauge group decompose correspondingly: $\text{Aut}_v(\mathcal{C}) = \text{Aut}_v(\mathcal{C})^{H_{\text{int}}} \ltimes \text{Aut}_v(\mathcal{C})^{H_{\text{ext}}} \simeq \mathcal{H} = \mathcal{H}_{\text{int}} \ltimes \mathcal{H}_{\text{ext}}$, with $\text{Aut}_v(\mathcal{C})^{H_{\text{int}}} = \text{Aut}_v(\mathcal{C}_{\text{rel}})$. So, we have the SES characterizing the relational configuration space-time bundle,

$$\text{id}_{\mathcal{C}_{\text{rel}}} \rightarrow \text{Aut}_v(\mathcal{C}_{\text{rel}}) \simeq \mathcal{H}_{\text{int}} \xrightarrow{\Delta} \text{Aut}(\mathcal{C}_{\text{rel}}) \rightarrow \text{Diff}(\mathcal{T}) \rightarrow \text{id}_{\mathcal{T}}. \quad (39)$$

The action of $\text{Aut}_v(\mathcal{C}) \simeq \mathcal{H}$ on $\beta \in \Omega^*(\mathcal{C})$ defines the *active gauge transformations* $\beta^X := \Xi^* \beta$, geometrically found via $\Xi^* \beta(\mathfrak{X}) = \beta(\Xi_* \mathfrak{X})$, where $\mathfrak{X} \in \Gamma(T\mathcal{C})$ is a generic vector field on $\mathcal{C}$ and, as special case of (17),

$$\Xi_* \mathfrak{X}_{|p} = R_{X(p)}^* \mathfrak{X}_{|p} + \{dX_{|p}(\mathfrak{X}_{|p})\}_{|\Xi(p)}^v = R_{X(p)}^* \mathfrak{X}_{|p} + \{dX_{|p}(\mathfrak{X}_{|p})\}_{|p}^v, \quad (40)$$

where the second equality arises because of the right invariance of fundamental vector fields. For example, for an Ehresmann connection $\omega \in \Omega^1_{\text{eq}}(\mathcal{C}, \text{Ad})$, s.t. $R_X^* \omega = \omega$ and $\omega(\chi^v) = \chi \in \mathfrak{h}$, we

have

$$\omega^X := \Xi^* \omega = \omega + dX, \quad (41)$$

which is an Abelian version of the first line of (19).⁷

Cocyclic geometry on $\mathcal{C}$ is defined via a 1-cocycle

$$\begin{aligned} C : \mathcal{C} \times H &\rightarrow G, \\ (p, X) &\mapsto C(p, X), \quad \text{s.t. } C(p, X + X') = C(p, X) \cdot C(p + X, X'), \end{aligned} \quad (42)$$

with corresponding linear 1-cocycle $a(\chi, p) := \frac{d}{d\tau} C(p, X_\tau) \Big|_{\tau=0}$,

$$\begin{aligned} a : \mathcal{C} \times \mathfrak{h} &\rightarrow \mathfrak{g}, \\ (p, \chi) &\mapsto a(\chi, p), \quad \text{s.t. } \chi^v \cdot a(\chi', p) - \chi'^v \cdot a(\chi, p) \\ &\quad + [a(\chi, p), a(\chi', p)]_{\mathfrak{g}} = 0. \end{aligned} \quad (43)$$

So that cocyclic tensorial forms $\alpha \in \Omega^*_{\text{tens}}(\mathcal{C}, C)$ and cocyclic connections $\varpi \in \Omega^*_{\text{eq}}(\mathcal{C}, C)$, s.t.

$$\begin{aligned} R_X^* \varpi_{|p+X} &= C(p, X)^{-1} \varpi_{|p} C(p, X) + C(p, X)^{-1} dC(\cdot, X)_{|p} \quad \text{and} \\ \varpi_{|p}(\chi_{|p}^v) &= a(\chi, p), \end{aligned} \quad (44)$$

have gauge transformations given by

$$\alpha^X = C(X)^{-1} \alpha \quad \text{and} \quad \varpi^X = C(X)^{-1} \varpi C(X) + C(X)^{-1} dC(X), \quad (45)$$

with the map $C(X) : \mathcal{C} \rightarrow G$ defined by $C(X(p)) = C(p, X(p))$, being special cases of (18)–(19).

3.2. Classical Mechanics on $\mathcal{C}$: The Action as a Cocyclic Connection

The dynamics of the configuration of N classical particles is given by the Lagrangian, which we take to be in general a *cocyclic tensorial* 1-form $L \in \Omega^1_{\text{tens}}(\mathcal{C}, c)$ where the 1-cocycle c takes values in $(\Omega^1(\mathcal{C}), +)$ seen as an additive Abelian group: It thus satisfies the defining property

$$c(p, X + X') = c(p, X) + c(p + X, X'), \quad (46)$$

the Abelian version of (7). A Lagrangian thus satisfies

$$\iota_{\chi^v} L = 0 \quad \text{and} \quad R_X^* L_{|p+X} = L_{|p} + c(p, X). \quad (47)$$

⁷ We may observe that, as is standard in bundle geometry, the kernel of a connection defines an horizontal subbundle $\ker \omega := H\mathcal{C}$ s.t. $T_p \mathcal{C} = V_p \mathcal{C} \oplus H_p \mathcal{C}$ at any $p \in \mathcal{C}$: Given a curve $c(\tau) \in \mathcal{C}$ s.t. $c(0) = t$ and $c(1) = t'$, it allows to define the horizontal lift $c^h(\tau) \in \mathcal{C}$ s.t. $\frac{d}{d\tau} c^h(\tau) \Big|_{\tau=0} \in \Gamma(H\mathcal{C})$ is a horizontal vector field. In turn, this allows to define the parallel transport map $c^h_* : \mathcal{C}_t \rightarrow \mathcal{C}_{t'}$, $p \mapsto c^h_*(p) = p'$, which prescribes a standard of identification across distinct fibers, i.e., determines (for each choice of connection ω) what it means for a configuration of particles p to remains “the same” across time.

The infinitesimal equivariance is then given by the Lie derivative,

$$\mathfrak{L}_{X^\vee} L = \frac{d}{d\epsilon} R_{X_\epsilon}^* L \Big|_{\epsilon=0} = \frac{d}{d\epsilon} c(-, X_\epsilon) \Big|_{\epsilon=0} =: a(\chi, -), \quad (48)$$

where $\frac{d}{d\epsilon} X_\epsilon \Big|_{\epsilon=0} = \chi \in \mathfrak{h}$, and $a(\chi; -)$ is the $\mathfrak{h}$ -1-cocycle (linear in χ) associated to L . The $\mathcal{H} \sim \text{Aut}_v(\mathcal{E})$ -transformation of L is then easily deduced to be

$$L^X := \Xi^* L = L + c(-, X) =: L + c(X). \quad (49)$$

The $\text{Lie}\mathcal{H} \sim \mathfrak{aut}_v(\mathcal{E})$ -transformation of L is $\mathfrak{L}_{X^\vee} L = a(\chi, -) =: a(\chi)$, with $\chi \in \text{Lie}\mathcal{H}$. The cocycle $a(\chi)$ will be shown to encode the Euler-Lagrange equations.

In a trivialisation, ϕ , i.e., in a bundle coordinate chart, we may therefore write (using the same notation above) $L|_{(t,x)} = (\phi^{-1})^* L|_p = \mathcal{L}(t, x) dt$. Given a local section $\sigma : \mathcal{U} \subset \mathcal{T} \rightarrow \mathcal{E}$, $t \mapsto \sigma(t) \simeq (t, x(t))$, where $\simeq$ is meant to indicate the expression in a bundle chart, the local representative of L on $\mathcal{T}$ is the familiar expression:

$$\sigma^* L|_t = L|_{\sigma(t)} \simeq L|_{(t, x(t))} = \mathcal{L}(t, x(t)) dt \in \Omega^1(\mathcal{T}, c). \quad (50)$$

Given a curve $\gamma : I \subset \mathbb{R} \rightarrow \mathcal{E}$, $\tau \mapsto \gamma(\tau) \simeq (t(\tau), x(\tau))$, with tangent vector field $\dot{\gamma} = \frac{dx}{d\tau} \frac{\partial}{\partial x} + \frac{dt}{d\tau} \frac{\partial}{\partial t} = \dot{x} \frac{\partial}{\partial x} + \dot{t} \frac{\partial}{\partial t} \in \Gamma(T\mathcal{E})$, the restriction of L “on” γ , actually its pullback to I , is:

$$\begin{aligned} \ell_{\gamma|I} &:= \gamma^* L|_I = L|_{\gamma} \simeq L|_{(t(\tau), x(\tau))} = \mathcal{L}(t(\tau), x(\tau)) \gamma^* dt|_\tau \\ &= \mathcal{L}(t(\tau), x(\tau)) i d\tau \in \Omega^1(I, c). \end{aligned} \quad (51)$$

Observe that (51) is the so-called “parametrized” formulation of non-relativistic mechanics (as exposed in, e.g., ref. [37]), while the standard (50) is the “deparametrized” or “unparametrized” version. We obtain the latter from the former via the choice $\dot{t} = 1$, which identifies the parameter τ with the time variable t , getting again the familiar expression $\ell_{\gamma|I} = \mathcal{L}(t, x(t)) dt$. Remark also that $\dot{x} = \frac{dx}{dt} \dot{t} =: x'$, where x' is the physical (measurable) velocity of x —i.e., the set of velocities $\{x'_i\}_{i=1, \dots, N}$ of the N particles under consideration. So, the condition $\dot{t} = 1$ thus identifies $\dot{x}$ with the velocity x' .⁸ This dual picture is unsurprising: the image of a local section σ is a curve in $\mathcal{E}$ parametrized by $t \in \mathcal{T}$.

Although there is no action of H and $\mathcal{H}$ on I , one may define indirectly their “action” on $\ell \in \Omega^1(I)$ —in the similar way that one may define their action on local representatives $\sigma^* L \in \Omega^1(\mathcal{T})$. The equivariance of ℓ_γ may be defined via the action of H on γ : since $R_X \gamma =: \gamma'$,

$$\begin{aligned} R_X^* \ell_\gamma &:= \gamma'^* L = (R_X \circ \gamma)^* L = \gamma^* R_X^* L = \gamma^* (L + c(-, X)) \\ &= \ell_\gamma + c_\gamma(-, X), \end{aligned} \quad (52)$$

where $c_\gamma(-, X) := c(-, X) \circ \gamma : I \rightarrow \Omega^1(\mathcal{E})$. Hence, the notation $\ell_\gamma \in \Omega^1(\mathcal{E}, c)$ and the legitimacy of considering ℓ_γ as a cocyclic 1-form. The same may be done for local representatives, using

$\sigma' = R_X \sigma$. The infinitesimal equivariance is obtained by linearising the above, which may be written as

$$\delta_X \ell_\gamma = a_\gamma(\chi, -) := a(\chi, -) \circ \gamma. \quad (53)$$

The action of the gauge group $\mathcal{H} \simeq \text{Aut}_v(\mathcal{E})$ is likewise obtained: considering $\gamma^X := \Xi \circ \gamma$, one defines

$$\ell_\gamma^X := (\gamma^X)^* L = (\Xi \circ \gamma)^* L = \gamma^* \Xi^* L = \gamma^* (L + c(-, X)) = \ell_\gamma + c_\gamma(X), \quad (54)$$

where $c_\gamma(X) := \gamma^* c(-, X) : I \rightarrow \Omega^1(\mathcal{E})$. This is analogous to the way one obtains the local version the active gauge transformation of L , and the way one defines passive gauge transformations of local representatives, i.e., gluings. The infinitesimal gauge transformation, i.e., the action of $\text{Lie}\mathcal{H} \sim \mathfrak{aut}_v(\mathcal{E})$, is obtained by linearising the above,

$$\delta_X \ell_\gamma = a_\gamma(\chi) := \gamma^* a(\chi, -), \quad (55)$$

which is also just $\gamma^* (\mathfrak{L}_{X^\vee} L)$, as expected from consistency of the definitions. Observe that $\delta_X \gamma = \chi(\gamma)$.

Let us observe that the tangent vector field $\dot{\gamma} \in \Gamma(T\mathcal{E})$ of a curve γ is right-invariant: This is clear from the fact that $\dot{\gamma} = \frac{d}{d\tau} \gamma(\tau) \Big|_{\tau=0} \simeq \dot{x} \frac{\partial}{\partial x} + \dot{t} \frac{\partial}{\partial t}$ and $R_{X^*} \dot{\gamma} := \frac{d}{d\tau} R_X \circ \gamma(\tau) \Big|_{\tau=0} = \frac{d}{d\tau} \gamma(\tau) + X \Big|_{\tau=0} = \dot{\gamma}$. Therefore, its pushforward by a vertical automorphism is, by (40), $\Xi_* \dot{\gamma} = \dot{\gamma} + \{dX(\dot{\gamma})\}^\vee$. We have $dX(\dot{\gamma}) = \frac{\partial X}{\partial t} \dot{t} + \frac{\partial X}{\partial x} \dot{x} =: \frac{dX}{d\tau} =: \dot{X}$, and since $\frac{\partial X}{\partial x} = 0$ as $X \in \Omega_{\text{inv}}^0(\mathcal{E})$, we get $\dot{X} = \frac{\partial X}{\partial t} \dot{t}$. So,

$$\Xi_* \dot{\gamma} = \dot{\gamma} + \{\dot{X}\}^\vee \simeq (\dot{x} + \dot{X}) \frac{\partial}{\partial x} + \dot{t} \frac{\partial}{\partial t}. \quad (56)$$

The choice $\dot{t} = 1$ gives $\dot{X} = \frac{\partial X}{\partial t} = X'$, with X' the physical velocity shift induced by the extended Galilean transformation $X \in \mathcal{H}$. In particular, the above specialises to vertical tangent vectors of vertical curves, s.t. $\dot{t} = 0$.

The action functional is defined by

$$S[\gamma] := \int_\gamma L = \int_I \gamma^* L =: \int_I \ell_\gamma \simeq \int_I \mathcal{L}(t(\tau), x(\tau)) \dot{t} d\tau. \quad (57)$$

Here again, the standard “unparametrised” version of the action is obtained by setting $\dot{t} = 1$.⁹ The equivariance of $S[\gamma]$ is derived from that of ℓ_γ : $R_X^* S[\gamma] = \int_I R_X^* \ell_\gamma = \int_I \ell_\gamma + c_\gamma(-, X) = S[\gamma] + \int_I c_\gamma(-, X)$. Which, for consistency, is cross-checked by

$$\begin{aligned} R_X^* S[\gamma] &:= S[\gamma'] = \int_{\gamma'} L = \int_I \gamma'^* L = \int_I (R_X \circ \gamma)^* L = \int_I \gamma^* R_X^* L \\ &= \int_I R_X^* L = \int_I L + c(-, X) =: S[\gamma] + \int_I c(-, X). \end{aligned} \quad (58)$$

⁸ We stress that $\dot{t} = 1$ is not a “gauge choice” (as many would be inclined to understand it), but actually reflects a dressing operation. We shall fully elaborate on this remark in a forthcoming paper.^[40]

⁹ Or by defining it as $S[\sigma] := \int_{\mathcal{U} \subset \mathcal{T}} \sigma^* L$, for a local section $\sigma : \mathcal{U} \subset \mathcal{T} \rightarrow \mathcal{E}|_{\mathcal{U}}$, $t \mapsto \sigma(t) \simeq (t, x(t))$ representing a trajectory/history.

Its infinitesimal equivariance is then defined by linearisation of the above,

$$\begin{aligned}\delta_\chi S|_\gamma &:= \frac{d}{d\epsilon} R_{X_\epsilon}^* S[\gamma] \Big|_{\epsilon=0} = \int_\gamma \frac{d}{d\epsilon} R_{X_\epsilon}^* L \Big|_{\epsilon=0} = \int_\gamma \mathfrak{L}_{X^\nu} L = \int_\gamma a(\chi, -), \\ &= \int_I a_\gamma(\chi, -),\end{aligned}\quad (59)$$

which is indeed the expression $\delta_\chi S|_\gamma = \int_I \delta_\chi \ell_\gamma$ intuitively expected from the definition of $S[\gamma]$. The $\mathcal{H} \sim \text{Aut}_v(\mathcal{C})$ -transformation of the action is likewise defined, via $\gamma^X := \Xi \circ \gamma$, as

$$\begin{aligned}S[\gamma]^X &:= S[\gamma^X] = \int_{\gamma^X} \Xi^* L = \int_{\gamma^X} L = \int_I (\gamma^X)^* L = \int_I \ell_\gamma^X \\ &= \int_I \ell_\gamma + c_\gamma(X) = S[\gamma] + \int_\gamma c(X),\end{aligned}\quad (60)$$

with $c(X) := c(-, X)$, which is consistent with the intuitive formula $S[\gamma]^X = \int_I \ell_\gamma^X$. By linearisation, the $\text{Lie}\mathcal{H} \sim \mathfrak{aut}_v(\mathcal{C})$ -transformation of S is easily found:

$$\delta_\chi S|_\gamma := \frac{d}{d\epsilon} \int_\gamma \Xi_\epsilon^* L \Big|_{\epsilon=0} = \int_\gamma \mathfrak{L}_{X^\nu} L = \int_\gamma a(\chi) = \int_I a_\gamma(\chi) =: \int_I \delta_\chi \ell_\gamma. \quad (61)$$

Equivalently, this is seen to be the Gateaux derivative along $\mathcal{H} \sim \text{Aut}_v(\mathcal{C})$: for $\gamma_\epsilon := \Xi_\epsilon \circ \gamma = \gamma + X_\epsilon(\gamma)$, we have

$$\delta_\chi S|_\gamma := \frac{d}{d\epsilon} [S[\gamma_\epsilon] - S[\gamma]] \Big|_{\epsilon=0} = \frac{d}{d\epsilon} \int_\gamma c(-, X_\epsilon) \Big|_{\epsilon=0} = \int_\gamma a(\chi). \quad (62)$$

This allows to define the classical dynamics as the critical points (curves) of S .

Considering the set $\mathcal{P}_{0,1} := \{\gamma : I \rightarrow \mathcal{C} \mid \gamma(\partial I) = \{p_0, p_1\}\}$ of paths with beginning and end points $p_0 \simeq (t_0, x_0)$ and $p_1 \simeq (t_1, x_1)$, respectively. Consider also the gauge subgroup $\mathcal{H}_{0,1} := \{X \in \mathcal{H} \mid X(t \leq t_0) = 0 = X(t \geq t_1) \text{ and } X(t_0 < t < t_1) \neq 0\} \subset \mathcal{H}$. We have that $\mathcal{H}_{0,1}$ acts freely and transitively on $\mathcal{P}_{0,1}$, i.e., for any choice of initial reference curve $\gamma \in \mathcal{P}_{0,1}$, any other can be written as γ^X for some $X \in \mathcal{H}_{0,1}$: so $\mathcal{P}_{0,1} \simeq \mathcal{H}_{0,1}$. The classical physical trajectory γ_c , i.e., the classical history, between the configurations p_0 and p_1 , is the critical point of $S[\gamma]$ for $\gamma \in \mathcal{P}_{0,1}$. By (61)–(62), we see that this is given by the vanishing of the linearised 1-cocycle. Said otherwise, the 1-cocycle $a(\chi)$ encodes the Euler–Lagrange equations, i.e., the dynamics, up to vanishing boundary terms:

$$\delta_\chi S|_{\gamma_c} = \int_I a(\chi) = \int_I a_{\gamma_c}(\chi) = \int_I E(\chi, \gamma_c) \equiv 0, \quad \forall \chi \in \text{Lie}\mathcal{H}_{0,1}, \quad (63)$$

where $E(\chi, \gamma) := \chi(\gamma) \cdot EL[\gamma] d\tau = \delta_\chi \gamma \cdot EL[\gamma] d\tau$ is the *equations of motion 1-form*, with $EL[\gamma]$ the Euler–Lagrange equations for the curve γ . The critical curve is s.t. $EL[\gamma_c] \equiv 0$. Then,

for any $\gamma \in \mathcal{P}_{0,1}$ there is a unique $X \in \mathcal{H}_{0,1}$ s.t. $\gamma = \gamma_c^X$, and by (60) one has

$$S[\gamma] = S[\gamma_c] + \int_{\gamma_c} c(X). \quad (64)$$

One may define the function $S_\gamma = S[\gamma; -, p_0] \in \Omega_{\text{tens}}^0(\mathcal{C}; c_\gamma)$ where $\gamma(\partial I) = \{p_0, p\}$ and $c_\gamma(-, X) := \int_\gamma c(-, X)$ is a 1-cocycle inherited from (58): i.e., we have $R_X^* S_\gamma = S_\gamma + c_\gamma(-, X)$. So its $\mathcal{H}$ -transformation is immediately read to be $S_\gamma^X := \Xi^* S_\gamma = S_\gamma + c_\gamma(X)$, where $c_\gamma(X) := c_\gamma(-, X)$. For $\gamma = \gamma_c$, the cocyclic 0-form S_{γ_c} is the *Hamilton Principal Function* (HPF). It satisfies the Hamilton equation: $\frac{\partial S_{\gamma_c}}{\partial t} = -H\left(x, \frac{\partial S_{\gamma_c}}{\partial x}; t\right)$, where the canonical conjugate momenta is by definition $\pi := \frac{\partial S_{\gamma_c}}{\partial x}$ and $H(x, \pi, t)$ is the Hamiltonian. Defining the 1-cocycle $C_\gamma(-, X) := \exp\{-\frac{i}{\hbar} c_\gamma(-, X)\}$, we find that the action induces the *flat cocyclic connection* $\varpi_{0,\gamma} := -\frac{i}{\hbar} dS_\gamma$ on $\mathcal{C}$, s.t.

$$R_X^* \varpi_{0,\gamma} = \varpi_{0,\gamma} + C_\gamma(-, X)^{-1} dC_\gamma(-, X).$$

$$\begin{aligned}\iota_{X^\nu} \varpi_{0,\gamma} &= -\frac{i}{\hbar} \iota_{X^\nu} dS_\gamma = -\frac{i}{\hbar} L_{X^\nu} S_\gamma = -\frac{i}{\hbar} \frac{d}{d\epsilon} R_{X_\epsilon}^* S_\gamma \Big|_{\epsilon=0} \\ &= -\frac{i}{\hbar} \frac{d}{d\epsilon} c_\gamma(-, X_\epsilon) \Big|_{\epsilon=0} = -\frac{i}{\hbar} \int_\gamma a(\chi) \\ &= \frac{d}{d\epsilon} C_\gamma(-, X_\epsilon) \Big|_{\epsilon=0},\end{aligned}\quad (65)$$

which is a special case of (15). From which follows that the $\mathcal{H}$ -transformation of $\varpi_{0,\gamma}$ is

$$\varpi_{0,\gamma}^X := \Xi^* \varpi_{0,\gamma} = \varpi_{0,\gamma} + C_\gamma(X)^{-1} dC_\gamma(X), \quad (66)$$

which is a special case of (19). It is also $(dS_\gamma)^X := \Xi^* dS_\gamma = dS_\gamma + dc_\gamma(X)$, consistent with the $\mathcal{H}$ -transformation of S_γ and $[d, \Xi^*] = 0$. Notice that $\iota_{X^\nu} \varpi_{0,\gamma} \neq 0$ since $\chi \in \text{Lie}\mathcal{H}$, contrary to what is done in (63) to determine the critical point γ_c . Actually, since $E(\chi; \gamma_c) \equiv 0$ by definition of γ_c , the verticality property of ϖ_{0,γ_c} gives the conserved Noether charge: In a bundle chart, $\chi^\nu \simeq \chi \frac{\partial}{\partial x}$ and $dS_{\gamma_c} = \frac{\partial S_{\gamma_c}}{\partial x} dx + \frac{\partial S_{\gamma_c}}{\partial t} dt$, so $\iota_{X^\nu} \varpi_{0,\gamma_c} = -\frac{i}{\hbar} \chi \frac{\partial S_{\gamma_c}}{\partial x} =: -\frac{i}{\hbar} \chi \pi$.¹⁰ Finally, we may observe that (64) implies a relation analogous to (66): For $X \in \mathcal{H}_{0,1}$ we have

$$\varpi_{0,\gamma} = \varpi_{0,\gamma_c} + C_{\gamma_c}(X)^{-1} dC_{\gamma_c}(X). \quad (67)$$

We are now but a couple of steps from a formulation of Quantum Mechanics.

¹⁰ This is also none other than the evaluation of χ^ν on the canonical 1-form (also known as the presymplectic potential) $-\frac{i}{\hbar} \theta := -\frac{i}{\hbar} \pi dx$, familiar in that form from Geometric Quantization.

3.3. Quantum Mechanics on $\mathcal{E}$: Schrödinger Equation as a Cocyclic Covariant Derivative

One may quite naturally consider the simplest possible cocyclic tensorial objects on $\mathcal{E}$ to which (65) is associated:

$$\begin{aligned} \psi_\gamma : \mathcal{E} &\rightarrow \mathbb{C} \quad \text{s.t.} \quad \psi_\gamma \in \Omega_{\text{tens}}^0(\mathcal{E}, C_\gamma) \\ p &\mapsto \psi_\gamma(p) \\ p + X &\mapsto \psi_\gamma(p + X) = C_\gamma(p, X)^{-1} \psi_\gamma(p), \end{aligned} \quad (68)$$

i.e., $R_X^* \psi_\gamma = C_\gamma(-, X)^{-1} \psi_\gamma$ with the 1-cocycle $C_\gamma : \mathcal{E} \times H \rightarrow U(1)$, $(p, X) \mapsto C_\gamma(p, X) := \exp\{-\frac{i}{\hbar} c_\gamma(p, X)\}$ defined via $S[\gamma]$. Their gauge transformation is thus $\psi_\gamma^X = C_\gamma(X)^{-1} \psi_\gamma =: \psi_{\gamma^X}$, for $X \in \mathcal{H}$. These can also be seen as sections $\tilde{\psi}_\gamma$ of the cocyclic associated bundle $E^{C_\gamma} := \mathcal{E} \times_{C_\gamma} \mathbb{C}$ over $\mathcal{T}$, so that $\Omega_{\text{tens}}^0(\mathcal{E}, C_\gamma) \simeq \Gamma(E^{C_\gamma})$. Let us name the latter the space of “pre-wave functions.” One then has the cocyclic covariant derivative preserving cocyclic tensoriality,

$$\begin{aligned} \bar{D}_{0,\gamma} : \Omega_{\text{tens}}^0(\mathcal{E}, C_\gamma) &\rightarrow \Omega_{\text{tens}}^1(\mathcal{E}, C_\gamma), \\ \psi_\gamma &\mapsto \bar{D}_{0,\gamma} \psi_\gamma := d\psi_\gamma + \varpi_{0,\gamma} \psi_\gamma. \end{aligned} \quad (69)$$

Notice that from the horizontality property of $\bar{D}_{0,\gamma} \psi_\gamma$ follows, in bundle coordinates, that $-i\hbar \frac{\partial}{\partial x} = \frac{\partial S_\gamma}{\partial x}$. Then, in view of (67), it must be the case that

$$\psi_\gamma = C_{\gamma_c}(X)^{-1} \psi_{\gamma_c}, \text{ and } \bar{D}_{0,\gamma} \psi_\gamma = C_{\gamma_c}(X)^{-1} \bar{D}_{0,\gamma_c} \psi_{\gamma_c} \text{ for } X \in \mathcal{H}_{0,1}. \quad (70)$$

Now, from (69) one defines the linear space of “wave functions” as $\mathcal{K} := \ker \bar{D}_{0,\gamma} \subset \Omega_{\text{tens}}^0(\mathcal{E}, C_\gamma)$, i.e., the covariantly constant cocyclic pre-wave functions. The name is justified by the observation that QM is encoded in the statement

$$\psi_{\gamma_c} \in \mathcal{K}, \quad \text{i.e.,} \quad \bar{D}_{0,\gamma_c} \psi_{\gamma_c} = 0. \quad (71)$$

Indeed, remember from the previous section that $\varpi_{0,\gamma_c} = -\frac{i}{\hbar} dS_{\gamma_c}$ where $S_{\gamma_c} = S[\gamma_c; -, p_0]$ is the HPF, satisfying the Hamilton–Jacobi equation. So, in bundle coordinate, we have

$$\begin{aligned} \bar{D}_{0,\gamma_c} \psi_{\gamma_c} &= d\psi_{\gamma_c} + \varpi_{0,\gamma_c} \psi_{\gamma_c} = 0 \\ \Rightarrow \quad dt \frac{\partial}{\partial t} \psi_{\gamma_c} + dx \frac{\partial}{\partial x} \psi_{\gamma_c} - \frac{i}{\hbar} \left(dt \frac{\partial S_{\gamma_c}}{\partial t} + dx \frac{\partial S_{\gamma_c}}{\partial x} \right) \psi_{\gamma_c} &= 0 \quad (72) \\ \hookrightarrow \quad dt \left(\frac{\partial}{\partial t} \psi_{\gamma_c} + \frac{i}{\hbar} H(x, \pi, t) \psi_{\gamma_c} \right) + dx \left(\frac{\partial}{\partial x} - \frac{i}{\hbar} \pi \right) \psi_{\gamma_c} &= 0. \end{aligned}$$

The horizontality property of $\bar{D}_{0,\gamma_c} \psi_{\gamma_c}$ means the identical vanishing of the coefficient of dx : which is an *eigenvalue problem* implying the quantum mechanical prescription for the momentum operator $\hat{\pi} := -i\hbar \frac{\partial}{\partial x}$. It is then immediate that, upon defining (still in bundle coordinates) the position operator $\hat{x}\psi = x\psi$, one gets the canonical commutation relation: $[\hat{x}, \hat{\pi}] = i\hbar \delta(x - x)$. These facts are “kinematical”, so to speak, arising from the geometric nature (the tensoriality) of the objects considered. Cocyclic

covariant constancy is a further dynamical constraint: It specifically implies the vanishing of the coefficient of dt in (72), which is just the eigenvalue problem for the Schrödinger equation. In summary, we have

$$\bar{D}_{0,\gamma_c} \psi_{\gamma_c} = 0 \quad \Rightarrow \quad \hat{\pi} := -i\hbar \frac{\partial}{\partial x} \quad \text{and} \quad i\hbar \frac{\partial}{\partial t} \psi_{\gamma_c} = H(\hat{x}, \hat{\pi}, t) \psi_{\gamma_c}. \quad (73)$$

The cocyclic covariant constancy condition (71) naturally encodes both the essential kinematical and dynamical axioms of Quantum Mechanics: the momentum operator and the Schrödinger equation. It extends to any $\psi_\gamma \in \mathcal{K}$ by (70). We remark that, per Section 2.1, QM is equivalently encoded in the statement $\tilde{\psi}_\gamma \in \tilde{\mathcal{K}}$, where $\tilde{\mathcal{K}} := \ker \tilde{\nabla}_0$ with $\tilde{\nabla}_0 : \Gamma(E^{C_\gamma}) \rightarrow \Omega^1(\mathcal{T}) \otimes E^{C_\gamma}$ the cocyclic covariant derivative induced on associated bundles $E^{C_\gamma} \rightarrow \mathcal{T}$.

One defines the standard (Hermitian) inner product, seen as a fiber-wise integration on $\mathcal{E}$,

$$\begin{aligned} \langle -, - \rangle : \mathcal{K} \times \mathcal{K} &\rightarrow C^\infty(\mathcal{T}, \mathbb{C}), \\ (\psi_\gamma, \psi'_\gamma) &\mapsto \langle \psi_\gamma, \psi'_\gamma \rangle := \int_{\mathcal{E}} \psi_\gamma^* \psi'_\gamma \text{vol}(\mathcal{E}), \end{aligned} \quad (74)$$

with ψ_γ^* the $\mathbb{C}$ -conjugate of ψ_γ , and $\text{vol}(\mathcal{E})$ the H_Δ -invariant volume form on fibers $\mathcal{E}_t = \mathbb{E}^{3N}$, $R_X^* \text{vol}(\mathcal{E}) = \text{vol}(\mathcal{E})$. Then, $\mathcal{H} := (\mathcal{K}; \langle -, - \rangle)$ is a Hilbert space.

We observe that one may obtain (71)–(73) as the field equation of a “meta-action” $\mathfrak{S}(\psi_\gamma, \varpi_{0,\gamma}) := \langle \psi_\gamma, \iota_{\mathfrak{X}} \bar{D}_{0,\gamma} \psi_\gamma \rangle$ —yet, we are aware, a not very apt name since this functional is dimensionless—where $\mathfrak{X} \in \Gamma(T\mathcal{E})$ is any non-vertical vector field ($\pi_* \mathfrak{X} \neq 0$), via the associated variational principle: $\frac{\delta}{\delta \psi} \mathfrak{S} = 0 \Rightarrow \bar{D}_{0,\gamma} \psi_\gamma = 0$. In typical quantum field-theoretical parlance, $\mathfrak{S}(\psi_\gamma, \varpi_{0,\gamma}) = \langle \psi_\gamma, \iota_{\mathfrak{X}} \bar{D}_{0,\gamma} \psi_\gamma \rangle$ may be understood as the expectation value of the cocyclic covariant derivative operator $\bar{D}_{0,\gamma}$. QM arises from requiring stationarity of this expectation value.

Path Integral: The a priori dependence of the above considerations on curves $\gamma \in \mathcal{P}_{0,1}$ between initial and final points $\gamma(\partial I) = \{p_0, p\}$ may seem unusual, but it naturally leads to the path integral description of QM. One may formally define the quantity depending only on the boundary by functional integration

$$K(p, p_0) := \int_{\mathcal{P}_{0,1}} D\gamma \quad \psi_\gamma = \int_{\mathcal{H}_{0,1}} DX \quad \psi_{\gamma^X} = \int_{\mathcal{H}_{0,1}} DX \quad C_{\gamma^X}(X)^{-1} \psi_{\gamma^X}, \quad (75)$$

where $D\gamma$ and DX are formal measures on the space of curves $\mathcal{P}_{0,1}$ and on $\mathcal{H}_{0,1}$ respectively. The first equality arises from the isomorphism $\mathcal{P}_{0,1} \simeq \mathcal{H}_{0,1}$ mentioned above when discussing classical dynamics, and specified by any reference curve $\gamma_r \in \mathcal{P}_{0,1}$ so that for any other $\gamma \in \mathcal{P}_{0,1}$, $\exists! X \in \mathcal{H}_{0,1}$ s.t. $\gamma = \gamma_r^X$. Stated otherwise, $\mathcal{H}_{0,1}$ acts freely and transitively on $\mathcal{P}_{0,1}$. One may then define the pre-wave function depending only on the final point by fiber integration over all initial conditions:

$$\psi(p) := \int_{\mathcal{E}_{|t_0}} K(p, p_0) \psi(p_0) \text{vol}(\mathcal{E}_{|t_0}), \quad (76)$$

where $\text{vol}(\mathcal{C}|_{t_0})$ is the volume form of the fiber $\mathcal{C}|_{t_0} \subset \mathcal{C}$ over $\pi(p_0) = t_0 \in \mathcal{T}$. In bundle coordinates, $\psi(p) \simeq \psi(t, x)$. The quantity (75) is usually seen as a propagator, describing the “transition amplitude” between the initial and final states. We may take the classical history as reference curve $\gamma_r = \gamma_c$, so that

$$K(p, p_0) := \int_{\mathcal{P}_{0,1}} \mathcal{D}\gamma \, \psi_\gamma = \int_{\mathcal{H}_{0,1}} \mathcal{D}X \, C_{\gamma_c}(X)^{-1} \psi_{\gamma_c}. \quad (77)$$

Restricting furthermore the attention to wave functions $\psi_\gamma \in \mathcal{X}$, solutions of $\bar{D}_{0,\gamma} \psi_\gamma = 0$ of the form

$$\psi_\gamma = \psi_0 \exp \int_\gamma \varpi_{0,\gamma} = \psi_0 \exp \frac{i}{\hbar} S_\gamma = \psi_0 \exp \frac{i}{\hbar} S[\gamma], \quad (78)$$

then the quantity (77) becomes the standard Dirac–Feynman propagator, or path integral (PI),

$$\begin{aligned} K(p, p_0) &= \int_{\mathcal{P}_{0,1}} \mathcal{D}\gamma \, \psi_0 \exp \frac{i}{\hbar} S[\gamma] \\ &= \int_{\mathcal{H}_{0,1}} \mathcal{D}X \, C_{\gamma_c}(X)^{-1} \psi_0 \exp \frac{i}{\hbar} S[\gamma_c] \\ &= \psi_0 \exp \frac{i}{\hbar} S[\gamma_c] \int_{\mathcal{H}_{0,1}} \mathcal{D}X \, \exp \frac{i}{\hbar} c_{\gamma_c}(X). \end{aligned} \quad (79)$$

Then, $\psi(p)$ in (76) is a wave function. Notice how geometric considerations automatically provide, in view of (73), a splitting of the PI in a contribution from the classical history, $\psi_{\gamma_c} = \psi_0 \exp \frac{i}{\hbar} S[\gamma_c] \in \mathcal{X}$, and a contribution from the functional integral of the 1-cocycle $c_{\gamma_c}(X) := \int_{\gamma_c} c(-, X) = \int_1 c(-, X) \circ \gamma_c$, inherited from the Lagrangian. In the semi-classical (e.g., WKB) analysis, the former contribution is known to contain the essential dynamical physical information, while the latter is a normalization factor.

The flat cocyclic connection $\varpi_{0,\gamma}$ is at the centre of our geometric formulation of both classical and quantum mechanics. As an intriguing exercise, one may entertain the possibility of allowing a non-flat cocyclic connection ϖ_γ and see what kind of generalisation of classical and quantum mechanics this leads to.

4. Relational Bundle Geometric Classical and Quantum Mechanics

In this section, we apply the Dressing Field Method to the configuration space-time bundle $\mathcal{C}$ to obtain relational reformulations of Classical and Quantum Mechanics. First, we consider how to realise the relational configuration space-time subbundle $\mathcal{C}_{\text{rel}} := \mathcal{C}/H_{\text{ext}} \xrightarrow{H_{\text{int}}} \mathcal{T}$, encoding the physical kinematics. In particular, we show how the notion of residual transformations of the 2nd kind in the DFM, parametrizing the choice of dressing, here parametrizes the multiple realisations of $\mathcal{C}_{\text{rel}}$ within $\mathcal{C}$. Then, we show how to obtain the dressed Lagrangian/action and wave function encoding the physical classical and quantum dynamics. In the context of relational QM, the group of residual transformations of the 2nd kind implements a notion of “physical reference frame covariance.”

4.1. Relational Configuration Space-Time Subbundle via DFM

Given that the configuration space-time bundle $\mathcal{C}$ is a 2-fibration

$$\mathcal{C} \xrightarrow{H_{\text{ext}}} \mathcal{C}_{\text{rel}} := \mathcal{C}/H_{\text{ext}} \xrightarrow{H_{\text{int}}} \mathcal{T}, \quad (80)$$

illustrated in Figure 1, with gauge group $H = H_{\text{int}} \ltimes H_{\text{ext}}$, we may use the DFM to realise the relational configuration space-time $\mathcal{C}_{\text{rel}}$, thereby reducing the gauge group H_{ext} . Specializing 2.2 to this context, a dressing field fit for that purpose is defined as

$$\begin{aligned} u : \mathcal{C} &\rightarrow H_{\text{ext}} = H_\Delta \triangleleft H \quad \text{s.t.} \quad R_X^* u = u - X, \quad \text{i.e.,} \\ u(p + X) &= u(p) - X. \end{aligned} \quad (81)$$

This is typically the rule for an Abelian dressing field. It is a H_{ext} -tensorial 0-form, thus it gauge transforms as $u^X := \Xi^* u = u - X$ for $X \in \mathcal{H}_{\text{ext}}$. It allows to define the map $f_u : \mathcal{C} \rightarrow \mathcal{C}$,

$$\begin{aligned} p &\mapsto f_u(p) := p + u(p), \quad \text{s.t.} \quad f_u \circ R_X = f_u \quad \text{and} \\ f_u \circ \Xi &= f_u \quad \text{for} \quad \Xi \in \text{Aut}_v(\mathcal{C})^{H_{\text{ext}}} \sim X \in \mathcal{H}_{\text{ext}}, \end{aligned} \quad (82)$$

whose image is a subbundle of $\mathcal{C}$ isomorphic to $\mathcal{C}_{\text{rel}}$: we thus denote it $\mathcal{C}_{\text{rel}}^u$.

The question is: Can we produce such a dressing field? As a matter of fact we can: Take $p \in \mathcal{C}$ as describing an instantaneous configuration of N particles, and, given bundle coordinates s.t. $p \simeq (t, x) \in \mathcal{U} \times H$ with $\mathcal{U} \subset \mathcal{T}$, let us define

$$u_i(p) = u_i(p_1, \dots, p_i, \dots, p_N) := -x_i = \overbrace{(-x_i, \dots, -x_i)}^{N \text{ times}} \in H_{\text{ext}} = \mathbb{R}^{3N}, \quad (83)$$

which clearly satisfies (81). So, the dressing field u_i outputs the vertical bundle coordinate (the spatial coordinate) of the i th particle of the configuration.¹¹ We have the associated map

$$\begin{aligned} f_{u_i}(p) &= p + u_i(p) = (p_1 - x_i, \dots, p_i - x_i, \dots, p_N - x_i) \\ &\simeq (t; x_1 - x_i, \dots, 0, \dots, x_N - x_i). \end{aligned} \quad (84)$$

That is, $\text{Im} f_{u_i} =: \mathcal{C}_{\text{rel}}^{u_i}$ is the realisation of the relational configuration space-time bundle $\mathcal{C}_{\text{rel}}$ from the viewpoint of the i th particle: it is a description of the N -configuration *from within*, by a participating “observer.” Notice that (84) is independent of the choice of bundle coordinates (i.e., of trivialisation).

Residual Transformations of the 1st Kind: Since H_{ext} is reduced via dressing, and thus acts trivially on $\mathcal{C}_{\text{rel}}^{u_i}$, one expects that there is still a residual action of the non-reduced subgroup $H_{\text{int}} \subset H$. It is indeed so, but with an interesting refinement that brings a better understanding of the isomorphism between $\mathcal{C}_{\text{rel}}^{u_i}$ and $\mathcal{C}_{\text{rel}}$.

The group H_{int} acts via $R_Y p = p + Y$. However, it is intuitive that from the viewpoint of any particle in the N -configuration, it

¹¹ It induces the Lie H_{ext} -valued Ehresmann connection $\omega_0 = -du_i$, giving the position of the i th particle as the standard of identification of fibers of $\mathcal{C}$ (i.e., of space) at distinct moments of time $\mathcal{T}$ —recall, without a connection, there is no canonical identification between fibers.

1521.3978, 2025, 12. Downloaded from <https://onlinelibrary.wiley.com/doi/10.1002/prop.70040> by Cochrane Italia, Wiley Online Library on [24/06/2026]. See the Terms and Conditions (<https://onlinelibrary.wiley.com/terms-and-conditions>) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

$$f_{u_i}(p) \simeq \left(t; x_1 - x_i, \dots, \overbrace{x_i - x_i}^0, \dots, \overbrace{x_j - x_i}^{-Z_{ij}(p)}, \dots, x_N - x_i \right),$$

$$f_{u_j}(p) \simeq \left(t; x_1 - x_j, \dots, \underbrace{x_i - x_j}_{Z_{ij}(p)}, \dots, \underbrace{x_j - x_j}_0, \dots, x_N - x_j \right)$$

$$= f_{u_i}(p) + Z_{ij}(p), \quad (93)$$

which showcases the relational symmetry of the viewpoint permutation. In the generic notation of (91)–(92),

$$f_{u^Z}(p) = f_u(p) + Z(p). \quad (94)$$

In other words, it is clear that the group $\mathcal{G}$ of transformations of the 2nd kind precisely encodes the notion of *physical reference frame covariance* of the relational description.

As seen in Section 3.2, a kinematical history is a curve $\gamma : I \rightarrow \mathcal{C}$. Its “dressing” is the projection on $\mathcal{C}_{\text{rel}}^u$,

$$\gamma^u := f_u \circ \gamma : I \rightarrow \mathcal{C}_{\text{rel}}^u,$$

$$\tau \mapsto \gamma^u(\tau) = \gamma(\tau) + \mathbf{u}(\gamma(\tau)) \quad (95)$$

$$\simeq (t(\tau); \bar{x}_1(\tau), \dots, \bar{x}_{N-1}(\tau)) = (t(\tau), \bar{x}(\tau)).$$

It is a *relational kinematical history* of a N particles configuration, as described from the internal viewpoint of any one of them, e.g., the i th if $\mathbf{u} = \mathbf{u}_i$. By definition it is $\mathcal{H}_{\text{ext}}$ -invariant: we define $(\gamma^u)^X := f_u \circ \Xi \circ \gamma = f_u \circ \gamma = \gamma^u$, for $X \in \mathcal{H}_{\text{ext}}$ generating $\Xi \in \text{Aut}_v(\mathcal{C})^{\mathcal{H}_{\text{ext}}}$. On account of (86), it supports the residual transformation of the 1st kind

$$(\gamma^u)^Y = \gamma^u + \bar{Y}(\gamma) =: (\gamma^u)^{\bar{Y}}, \quad \text{for } Y \in \mathcal{H}_{\text{int}} \simeq \bar{Y} \in \mathcal{H}_{\text{int}}^u, \quad (96)$$

$$\simeq (t(\tau); \bar{x}_1(\tau), \dots, \bar{x}_{N-1}(\tau)) + (\bar{Y}_1(\tau), \dots, \bar{Y}_{N-1}(\tau)),$$

which describes a *physically distinct* relational kinematical history in $\mathcal{C}_{\text{rel}}^u$. The infinitesimal variation of a physical relational kinematics is then: $\delta_Y \gamma^u := \frac{d}{d\varepsilon} (\gamma^u)^{Y_\varepsilon} - \gamma^u \Big|_{\varepsilon=0} = \bar{Y}(\gamma)$, with $\frac{d}{d\varepsilon} Y_\varepsilon \Big|_{\varepsilon=0} = \mathbf{Y} \in \text{Lie} \mathcal{H}_{\text{int}}$.

On account of (93)–(94), γ^u support the action of $\mathcal{G}$

$$(\gamma^u)^Z = \gamma^u + Z(\gamma), \quad \text{for } Z \in \mathcal{G}, \quad (97)$$

where $(\gamma^u)^Z$ and γ^u describe the *same* relational kinematical history: specialising the notation, this is indeed by (93)

$$\gamma^u(\tau) = \gamma^{u_i}(\tau) \simeq (t(\tau); \bar{x}_1^i(\tau), \dots, \bar{x}_{N-1}^i(\tau)) = (t(\tau), \bar{x}^i(\tau)), \quad \text{and}$$

$$(\gamma^u)^Z(\tau) = \gamma^{u_i}(\tau) = \gamma^{u_i}(\tau) + Z_{ij}(\gamma(\tau)) \simeq (t(\tau); \bar{x}_1^j(\tau), \dots, \bar{x}_{N-1}^j(\tau))$$

$$= (t(\tau), \bar{x}^j(\tau)). \quad (98)$$

Equation (97) demonstrates the physical reference frame covariance of the relational description of a kinematical history. We can now move on to the relational description of classical and quantum mechanics.

4.2. Relational Classical Mechanics, and Physical Frame Covariance

As a special case of the general formulae (25)–(27) of the DFM, the dressed Lagrangian 1-form is

$$L^u := f_u^* L = L + c(\mathbf{u}), \quad (99)$$

where we used the notation $c(\mathbf{u}) := c(_, \mathbf{u})$. It is $\mathcal{H}_{\text{ext}}$ -basic by construction, hence $\mathcal{H}_{\text{ext}}$ -invariant. This is checked explicitly using the $\mathcal{H}_{\text{ext}}$ -transformation of L (49) and the fact that, by the Abelian version (46) of (8), we get,

$$c(\mathbf{u})^X(p) := c(_, \mathbf{u})^X(p) = (\Xi^* c(_, \mathbf{u}))(p) = c(\Xi(p), \Xi^* \mathbf{u}(p))$$

$$= c(p + X(p), -X(p) + \mathbf{u}(p))$$

$$= c(p + X(p), -X(p)) + c(p + X(p) - X(p), \mathbf{u}(p))$$

$$= -c(p, X(p)) + c(p, \mathbf{u}(p)),$$

$$c(\mathbf{u})^X = c(\mathbf{u}) - c(X), \quad \text{for } X \in \mathcal{H}_{\text{ext}}, \quad (100)$$

i.e., an Abelian version of (the gauge transformation of) a cocyclic dressing field (29). Remark that we thus have

$$\mathfrak{L}_{\chi^v} c(\mathbf{u}) = -a(\chi) \quad \text{for } \chi \in \text{Lie} \mathcal{H}_{\text{ext}}, \quad (101)$$

which is the “classical anomaly” (48). We may also observe that the notion of cocyclic dressing field $c(\mathbf{u})$, as it appears here in Mechanics, encompasses the notion of *Wess-Zumino counterterms* introduced to “restore” gauge-invariance and/or implement anomaly cancellation—see, e.g., Section 12.3 in ref. [42], Chap. 15 in [43], or the end of Chap.4 in [44]. See ref. [7] for further discussion of this point in the context of general-relativistic Gauge Field Theory.

Notice that (99) can be obtained from (49) substituting the gauge parameter by the dressing field, $X \rightarrow \mathbf{u}$. This illustrates the DFM *rule of thumb* described in Section 2.2: To obtain the dressing of an object, compute first its gauge transformation, then substitute $X \rightarrow \mathbf{u}$ in the result. We shall often see instances of it in what follows.

On account of (86), L^u exhibits residual $\mathcal{H}_{\text{int}} \simeq \mathcal{H}_{\text{int}}^u$ -transformations (of the 1st kind), which by (99) may be found by: $(L^u)^Y = L^Y + c(\mathbf{u})^Y$. The $\mathcal{H}_{\text{int}}$ -transformation of L is known as a special case of (49), so we need only to find $c(\mathbf{u})^Y$. For this let us first notice the very simple lemma

$$c(p, \mathbf{u} + \mathbf{Y}) = c(p, \mathbf{u}) + c(p + \mathbf{u}, \mathbf{Y}) = c(p, \mathbf{Y}) + c(p + \mathbf{Y}, \mathbf{u}),$$

$$\Rightarrow c(p + \mathbf{Y}, \mathbf{u}) - c(p + \mathbf{u}, \mathbf{Y}) = c(p, \mathbf{u}) - c(p, \mathbf{Y}), \quad (102)$$

where we used the definition of an Abelian 1-cocycle. So, for $Y \in \mathcal{H}_{\text{int}}$ generating $\Xi \in \text{Aut}_v(\mathcal{E})^{\mathcal{H}_{\text{int}}}$ and $\mathbf{u} = \mathbf{u}_i$:

$$\begin{aligned} c(\mathbf{u}_i)^Y(p) &:= c(_, \mathbf{u}_i)^Y(p) = (\Xi^* c(_, \mathbf{u}_i))(p) = c(\Xi(p), \Xi^* \mathbf{u}_i(p)) \\ &= c(p + Y(p), \mathbf{u}_i(p) - Y_i(p)) \\ &= c(p + Y(p) + \mathbf{u}_i(p), -Y_i(p)) + c(p + Y(p), \mathbf{u}_i(p)) \\ &= c(p + \mathbf{u}_i(p), Y(p) - Y_i(p)) - c(p + \mathbf{u}_i(p), Y(p)) \\ &\quad + c(p + Y(p), \mathbf{u}_i(p)) \\ &= c(p + \mathbf{u}_i(p), \tilde{Y}^i(p)) + c(p, \mathbf{u}_i(p)) - c(p, Y(p)) \\ &= c(p, \mathbf{u}_i(p) + \tilde{Y}^i(p)) - c(p, Y), \end{aligned}$$

$$\begin{aligned} \hookrightarrow c(\mathbf{u})^Y &= c(\mathbf{u}) + c(f_u(_, \tilde{Y})) - c(Y) \\ &= c(\mathbf{u} + \tilde{Y}) - c(Y), \end{aligned} \quad (103)$$

using again the defining property of c . We thus compute the residual $\mathcal{H}_{\text{int}} \simeq \mathcal{H}_{\text{int}}^u$ -transformation of L^u to be

$$\begin{aligned} (L^u)^Y &= L^Y + c(\mathbf{u})^Y = L + c(Y) + c(\mathbf{u}) + c(f_u(_, \tilde{Y})) - c(Y) \\ &= L + c(\mathbf{u}) + c(f_u(_, \tilde{Y})) \\ &= L^u + c(f_u(_, \tilde{Y})), \end{aligned} \quad (104)$$

which is also $(L^u)^Y = L + c(p, \mathbf{u} + \tilde{Y})$. Equation (104) is precisely what geometric reasoning immediately gives: Considering L^u as a form on the subbundle $\mathcal{E}_{\text{rel}}^u = \text{Im} f_u$, its gauge group $\mathcal{H}_{\text{int}}^u$ acts in exactly the same formal way as $\mathcal{H}_{\text{int}}$ acts on L , i.e., for $\tilde{Y} \in \mathcal{H}_{\text{int}}^u$ generating $\Xi \in \text{Aut}_v(\mathcal{E}_{\text{rel}}^u)$ we have

$$(L^u)^{\tilde{Y}} := \Xi^* L^u = L^u + c(f_u(_, \tilde{Y})). \quad (105)$$

We also see that indeed $(L^u)^Y = (L^u)^{\tilde{Y}}$, as expected from $\mathcal{H}_{\text{int}} \simeq \mathcal{H}_{\text{int}}^u$, but ensured by the peculiar $\mathcal{H}_{\text{int}}^u$ -transformation (103) of the cocyclic dressing field $c(\mathbf{u})$. For $\mathbf{Y} \in \text{Lie} \mathcal{H}_{\text{int}}$ we read from (104) the infinitesimal transformation

$$\mathfrak{L}_{Y^v} L^u = \mathfrak{L}_{\tilde{Y}^v} L^u = a(\tilde{Y}, f_u(_)). \quad (106)$$

This cross-checked via (99) and from the transformation of the cocyclic dressing read from (103)

$$\mathfrak{L}_{Y^v} c(\mathbf{u}) = a(\tilde{Y}, f_u(_)) - a(\mathbf{Y}), \quad (107)$$

by which we get,

$$\mathfrak{L}_{Y^v} L^u = \mathfrak{L}_{Y^v} L + \mathfrak{L}_{Y^v} c(\mathbf{u}) = a(\mathbf{Y}) + a(\tilde{Y}, f_u(_)) - a(\mathbf{Y}) = a(\tilde{Y}, f_u(_)). \quad (108)$$

This computation is instructive by itself for the following reason: As we observed in Section 3.2, the infinitesimal 1-cocycle $a(\mathbf{Y}) = a(\mathbf{Y}, _)$ essentially encodes the Euler-Lagrange equations, and

therefore the dressed infinitesimal cocycle $a(\tilde{Y}, f_u(_))$ —which may be understood as a “residual classical anomaly,” in analogy with field theory—will similarly be shown to encode the *relational Euler–Lagrange equations*. So, Equation (108) establishes the link between the bare and relational equations of motions.

On account of (91), L^u supports $\mathcal{G}$ -transformations (of the 2nd kind), which by (99) may be found by writing: $(L^u)^Z = L^Z + c(\mathbf{u})^Z = L + c(\mathbf{u})^Z$. So we just need to determine $c(\mathbf{u})^Z$, which is easily done:

$$\begin{aligned} c(\mathbf{u})^Z(p) &:= c(p, \mathbf{u}(p))^Z = c(p, \mathbf{u}^Z(p)) = c(p, \mathbf{u}(p) + Z(p)) \\ &= c(p, \mathbf{u}(p)) + c(p + \mathbf{u}(p), Z(p)), \\ &\hookrightarrow c(\mathbf{u})^Z = c(\mathbf{u} + Z) = c(\mathbf{u}) + c(f_u(_, Z)). \end{aligned} \quad (109)$$

This is the Abelian version of (30). We thus find the $\mathcal{G}$ -transformation of L^u to be

$$(L^u)^Z = L^u + c(f_u(_, Z)), \quad (110)$$

which is also $(L^u)^Z = L^{u^Z} = L + c(\mathbf{u} + Z)$. Equation (110) is but a special case of the general formulae (31) of the DFM.

Despite their formal similarity, Equations (104) and (110) carry very different meanings: Equation (104) specifies how the description of the Lagrangian form changes from the point of view of the same particle participating in a distinct physical configuration, a change which is continuous and parametrized by the Lie group $\mathcal{H}_{\text{int}} \simeq \mathcal{H}_{\text{int}}^u$. So that both $(L^u)^Y$ and (L^u) “live” on the same subbundle $\mathcal{E}_{\text{rel}}^u$. Although (110) specifies how the description of the Lagrangian changes when shifting to a distinct particle in the same physical configuration, a change which is discrete and parametrized by the finite group $\mathcal{G}$ (89): Specialising the notation, (110) is $L^{u_i} = L^{u_i} + c(f_{u_i}(_, Z_{ij}))$, and indicates how to relate the relational descriptions of the Lagrangian on the bundles $\mathcal{E}_{\text{rel}}^{u_i}$ and $\mathcal{E}_{\text{rel}}^{u_j}$ in Figure 2.

Since both $(L^u)^Z = L^{u_j}$ and $(L^u) = L^{u_i}$ are the same object seen from two distinct perspectives, it is of interest to see explicitly how their residual $\mathcal{H}_{\text{int}}$ -transformations are related. By (104) we know that

$$\begin{aligned} (L^{u_j})^Y &= L^{u_j} + c(f_{u_j}(_, \tilde{Y}^j)), \quad \text{with } \tilde{Y}^j := Y - Y_j, \\ (L^{u_i})^Y &= L^{u_i} + c(f_{u_i}(_, \tilde{Y}^i)), \quad \text{with } \tilde{Y}^i := Y - Y_i. \end{aligned} \quad (111)$$

By (110), we have $(L^{u_j})^Y = (L^{u_i})^Y + c(f_{u_i}(_, Z_{ij}))^Y$. To compute the $c(f_{u_i}(_, Z_{ij}))^Y$ the $\mathcal{H}_{\text{int}}$ -transformation of $Z = Z_{ij}$ is needed. Since by definition $p_j = p_i - Z_{ij}(p)$, under the action of $\mathcal{H}_{\text{int}}$ we have $p_j + Y_j = p_i + Y_i - Z_{ij}^Y(p)$, so that $p_i - Z_{ij}(p) + Y_j = p_i + Y_i - Z_{ij}^Y(p)$, and we obtain finally

$$Z_{ij}^Y = Z_{ij} + (Y_i - Y_j) = Z_{ij} + (\tilde{Y}^j - \tilde{Y}^i). \quad (112)$$

So, we may compute

$$\begin{aligned}
 c(f_{u_i}(-), Z_{ij})^Y &= c(f_{u_i}(-)^Y, Z_{ij}^Y) \\
 &= c(f_{u_i}(-) + \tilde{Y}^i, Z_{ij} + (\tilde{Y}^j - \tilde{Y}^i)) \\
 &= c(f_{u_i}(-), \tilde{Y}^i + (\tilde{Y}^j - \tilde{Y}^i) + Z_{ij}) - c(f_{u_i}(-), \tilde{Y}^i) \\
 &= c(f_{u_i}(-), \tilde{Y}^j + Z_{ij}) - c(f_{u_i}(-), \tilde{Y}^i) \\
 &= c(f_{u_i}(-), Z_{ij}) + c(f_{u_i}(-) + Z_{ij}, \tilde{Y}^j) - c(f_{u_i}(-), \tilde{Y}^i) \\
 &= c(f_{u_i}(-), Z_{ij}) + c(f_{u_j}(-), \tilde{Y}^j) - c(f_{u_i}(-), \tilde{Y}^i).
 \end{aligned} \tag{113}$$

Remark the similarity with (103). We thus find

$$\begin{aligned}
 (L^{u_j})^Y &= (L^{u_i})^Y + c(f_{u_i}(-), Z_{ij})^Y \\
 &= L^{u_i} + \cancel{c(f_{u_i}(-), \tilde{Y}^i)} + c(f_{u_i}(-), Z_{ij}) \\
 &\quad + c(f_{u_i}(-) + Z_{ij}, \tilde{Y}^j) - \cancel{c(f_{u_i}(-), \tilde{Y}^i)} \\
 &= L^{u_j} + c(f_{u_i}(-) + Z_{ij}, \tilde{Y}^j) = L^{u_j} + c(f_{u_j}(-), \tilde{Y}^j),
 \end{aligned} \tag{114}$$

as expected. Again, the subtle cocyclic structure enforces consistency. The linear version of the above is of special interest. From (106) we have that

$$\mathfrak{L}_{Y^v} L^{u_j} = a(\tilde{Y}^j, f_{u_j}(-)), \quad \text{and} \quad \mathfrak{L}_{Y^v} L^{u_i} = a(\tilde{Y}^i, f_{u_i}(-)). \tag{115}$$

By (110) we must have $\mathfrak{L}_{Y^v} L^{u_j} = \mathfrak{L}_{Y^v} L^{u_i} + \mathfrak{L}_{Y^v} c(f_{u_i}(-), Z_{ij})$, and by (113) we get

$$\begin{aligned}
 \mathfrak{L}_{Y^v} c(f_{u_i}(-), Z_{ij}) &= a(\tilde{Y}^j, f_{u_i}(-) + Z_{ij}) - a(\tilde{Y}^i, f_{u_i}(-)) \\
 &= a(\tilde{Y}^j, f_{u_j}(-)) - a(\tilde{Y}^i, f_{u_i}(-)).
 \end{aligned} \tag{116}$$

Remark the similarity with (107). So, analogously to (108), we have

$$\begin{aligned}
 \mathfrak{L}_{Y^v} L^{u_j} &= \mathfrak{L}_{Y^v} L^{u_i} + \mathfrak{L}_{Y^v} c(f_{u_i}(-), Z_{ij}) = a(\tilde{Y}^i, f_{u_i}(-)) \\
 &\quad + a(\tilde{Y}^j, f_{u_i}(-) + Z_{ij}) - a(\tilde{Y}^i, f_{u_i}(-)) \\
 &= a(\tilde{Y}^j, f_{u_j}(-)),
 \end{aligned} \tag{117}$$

as expected. As we show below, at the end of this section, this computation is key to establish the correspondence between formulations of the *relational variational principle* as seen from distinct perspectives within the same kinematical history γ .

The dressed Lagrangian restricted along a kinematical history γ is,

$$\begin{aligned}
 \mathcal{L}_\gamma^u &:= \gamma^* L^u = (f_u \circ \gamma)^* L =: (\gamma^u)^* L := \mathcal{L}_{\gamma^u} \\
 &= (\mathcal{L} \circ \gamma^u) (\gamma^u)^* dt \\
 &= (\mathcal{L} \circ \gamma^u) \dot{t} d\tau,
 \end{aligned} \tag{118}$$

i.e., $\mathcal{L}_\gamma^u(\tau) = \mathcal{L}_{\gamma^u}(\tau) = \mathcal{L}(t(\tau), \tilde{x}(\tau)) \dot{t} d\tau \in \Omega^1(I, \mathbb{C})$.

This expression is conceptually clear: it is the relational parametrized Lagrangian from the viewpoint of one of the N particles in γ , which thus describes the kinematics as γ^u . From (99), we get the computationally useful expression

$$\mathcal{L}_{\gamma^u} = \mathcal{L}_\gamma^u := \gamma^* L^u = \gamma^* (L + c(u)) =: \mathcal{L}_\gamma + c_\gamma(u), \tag{119}$$

giving the relational Lagrangian in terms of the bare Lagrangian and its 1-cocycle. It could have been obtained from the $\mathcal{H}_{\text{ext}}$ -transformation of $\mathcal{L}$ (54) via the DFM rule of thumb. The residual $\mathcal{H}_{\text{int}} \simeq \mathcal{H}_{\text{int}}^u$ -transformation is, by (104),

$$\begin{aligned}
 (\mathcal{L}_\gamma^u)^Y &= (\mathcal{L}_\gamma^u)^{\tilde{Y}} := \gamma^* (L^u + c(f_u(-), \tilde{Y})) = \mathcal{L}_\gamma^u + c(f_u(\gamma), \tilde{Y}(\gamma)), \\
 &\text{or equivalently } (\mathcal{L}_{\gamma^u})^Y = (\mathcal{L}_{\gamma^u})^{\tilde{Y}} = \mathcal{L}_{\gamma^u} + c_{\gamma^u}(\tilde{Y}).
 \end{aligned} \tag{120}$$

It is the relational parametrized Lagrangian from the internal viewpoint of the same particle in a *physically distinct* kinematical history $(\gamma^u)^{\tilde{Y}}$ given by (96). This may be checked by $(\mathcal{L}_{\gamma^u})^Y := \gamma^* \Xi^* f_u^* L = (\Xi \circ \gamma)^* f_u^* L = (\gamma^Y)^* f_u^* L = (f_u \circ \gamma^Y)^* L$. Then, observe that $f_{u_i} \circ \gamma^Y = \gamma^Y + u_i(\gamma^Y) = \gamma + Y(\gamma) + u_i(\gamma + Y(\gamma)) = \gamma + Y(\gamma) + u_i(\gamma) - Y_i(\gamma) = f_{u_i}(\gamma) + \tilde{Y}^i(\gamma) = \gamma^{u_i} + \tilde{Y}^i(\gamma) =: (\gamma^{u_i})^{\tilde{Y}}$. So that $(\mathcal{L}_{\gamma^u})^Y = \mathcal{L}_{(\gamma^{u_i})^{\tilde{Y}}}$. The infinitesimal transformation is then

$$\delta_Y \mathcal{L}_\gamma^u := \frac{d}{d\epsilon} (\mathcal{L}_\gamma^u)^{Y_\epsilon} - \mathcal{L}_\gamma^u \Big|_{\epsilon=0} = \frac{d}{d\epsilon} c_{\gamma^u}(\tilde{Y}_\epsilon) \Big|_{\epsilon=0} =: a_{\gamma^u}(\tilde{Y}), \tag{121}$$

which is the variation of the relational Lagrangian under variation of the physical relational kinematical history $\delta\gamma^u = \tilde{Y}(\gamma)$. This is consistent with (106), writing $\delta_Y \mathcal{L}_\gamma^u = \gamma^* (\mathfrak{L}_{Y^v} L^u) = \gamma^* a(\tilde{Y}, f_u(-))$.

The $\mathcal{G}$ -transformation of $\mathcal{L}_\gamma^u$ is, by (110),

$$\begin{aligned}
 (\mathcal{L}_\gamma^u)^Z &:= \gamma^* (L^u + c(f_u(-), Z)) \\
 &= \mathcal{L}_\gamma^u + c(f_u(\gamma), Z(\gamma)) \\
 &= \mathcal{L}_\gamma^u + c(\gamma^u, Z(\gamma)), \quad \text{or equivalently } (\mathcal{L}_{\gamma^u})^Z \\
 &= \mathcal{L}_{\gamma^u} + c_{\gamma^u}(Z),
 \end{aligned} \tag{122}$$

which is the relational parametrized Lagrangian from the internal viewpoint of a distinct particle in the same physical kinematical history $(\gamma^u)^Z$ given by (97). Equation (122) showcases the physical reference frame covariance of the relational dynamics, controlled by the finite group $\mathcal{G}$. Taking a concrete case, (122) is $\mathcal{L}_\gamma^{u_j} = \mathcal{L}_\gamma^{u_i} + c(\gamma^{u_i}, Z_{ij}(\gamma))$.

Consequently, by (99) and (118)–(119), we find the dressed, relational action to be

$$\begin{aligned} S[\gamma]^u &:= \int_{\gamma} L^u = \int_I \ell_{\gamma}^u = \int_I (\mathcal{L} \circ \gamma^u) \dot{t} d\tau \\ &\simeq \int_I \mathcal{L}(t(\tau), \tilde{x}(\tau)) \dot{t} d\tau \\ &\stackrel{\text{b}}{=} \int_I \ell_{\gamma} + c_{\gamma}(u) =: S[\gamma] + c_{\gamma}(u), \end{aligned} \quad (123)$$

giving both the conceptually clear expression and the computationally useful form. The latter, as usual, could be directly read from the $\mathcal{H}_{\text{ext}}$ -transformation (60) of $S[\gamma]$ by the DFM rule of thumb. Quite naturally, it is the action associated to the relational kinematics γ^u , as it is clear from

$$S[\gamma]^u := \int_{\gamma} L^u = \int_{\gamma} f_{\gamma}^* L = \int_{f_{\gamma} \circ \gamma} L = \int_{\gamma^u} L =: S[\gamma^u]. \quad (124)$$

From (104) and (120), we find that the $\mathcal{H}_{\text{int}} \simeq \mathcal{H}_{\text{int}}^u$ -transformation of the dressed action is

$$\begin{aligned} (S[\gamma]^u)^Y &:= \int_{\gamma} (L^u)^Y = \int_{\gamma} L^u + c(f_u(-), \tilde{Y}) \\ &= S[\gamma]^u + c_{\gamma^u}(\tilde{Y}), \end{aligned} \quad (125)$$

$$\begin{aligned} \text{or equivalently } (S[\gamma^u])^Y &:= \int_I (\ell_{\gamma^u})^Y = \int_I \ell_{\gamma^u} + c_{\gamma^u}(\tilde{Y}) \\ &= S[\gamma^u] + c_{\gamma^u}(\tilde{Y}). \end{aligned}$$

It is the dressed action seen from the viewpoint of the same particle in a physically distinct kinematical history $(\gamma^u)^{\tilde{Y}}$, given by (96): indeed, one also writes $(S[\gamma^u])^Y := S[(\gamma^u)^{\tilde{Y}}] = S[(\gamma^u)^{\tilde{Y}}]$. The infinitesimal residual transformation is then, for $\mathbf{Y} \in \text{Lie}\mathcal{H}_{\text{int}}$,

$$\begin{aligned} \delta_{\mathbf{Y}} S_{|\gamma^u} &:= \frac{d}{d\epsilon} \int_{\gamma} \Xi_{\epsilon}^* L^u \Big|_{\epsilon=0} = \int_{\gamma} \mathfrak{L}_{\mathbf{Y}^v} L^u = \int_{\gamma} a(\tilde{\mathbf{Y}}, f_u(-)) \\ &= \int_I a_{\gamma^u}(\tilde{\mathbf{Y}}) =: \int_I \delta_{\mathbf{Y}} \ell_{\gamma^u}^u. \end{aligned} \quad (126)$$

Equivalently, this is the Gateaux derivative of $S[\gamma^u]$ along $\mathcal{H}_{\text{int}}^u \sim \text{Aut}_v(\mathcal{C}_{\text{rel}}^u)$: for $\gamma_{\epsilon}^u := (\gamma^u)^{Y_{\epsilon}} = \gamma^u + \tilde{\mathbf{Y}}_{\epsilon}(\gamma)$ and infinitesimal variation of the relational kinematical history $\delta_{\mathbf{Y}} \gamma^u = \tilde{\mathbf{Y}}(\gamma)$, we have

$$\begin{aligned} \delta_{\mathbf{Y}} S_{|\gamma^u} &:= \frac{d}{d\epsilon} S[\gamma_{\epsilon}^u] - S[\gamma^u] \Big|_{\epsilon=0} \\ &= \frac{d}{d\epsilon} \int_{\gamma} c(f_u(-), \mathbf{Y}_{\epsilon}) \Big|_{\epsilon=0} = \int_{\gamma} a(\tilde{\mathbf{Y}}, f_u(-)). \end{aligned} \quad (127)$$

This allows to define the critical points of S^u , i.e., the *classical relational dynamics*.

We proceed as in Section 3.2; Consider $\mathcal{P}_{0,1}^u := \{\gamma^u : I \rightarrow \mathcal{C}_{\text{rel}}^u \mid \gamma^u(\partial I) = \{q_0, q_1\}\}$ the set of paths with beginning and end points $q_0 = f_u(p_0) \simeq (t_0, \tilde{x}_0)$ and $q_1 = f_u(p_1) \simeq (t_1, \tilde{x}_1)$, i.e.,

$\mathcal{P}_{0,1}^u = \{\gamma^u \mid \gamma \in \mathcal{P}_{0,1}\}$. Consider also the gauge subgroup $\mathcal{H}_{\text{int}|0,1} := \{\mathbf{Y} \in \mathcal{H}_{\text{int}} \mid \mathbf{Y}(t \leq t_0) = 0 = \mathbf{Y}(t \geq t_1) \text{ and } \mathbf{Y}(t_0 < t < t_1) \neq 0\} \subset \mathcal{H}_{\text{int}}$, isomorphic to the similarly defined (*mutatis mutandis*) $\mathcal{H}_{\text{int}|0,1}^u \subset \mathcal{H}_{\text{int}}^u$. We have that $\mathcal{H}_{\text{int}|0,1} \simeq \mathcal{H}_{\text{int}|0,1}^u$ acts freely and transitively on $\mathcal{P}_{0,1}^u$, i.e., for any choice of initial reference curve $\gamma^u \in \mathcal{P}_{0,1}^u$, any other is written as $(\gamma^u)^{\mathbf{Y}} = (\gamma^u)^{\tilde{\mathbf{Y}}}$ for some $\mathbf{Y} \in \mathcal{H}_{\text{int}|0,1} \simeq \tilde{\mathbf{Y}} \in \mathcal{H}_{\text{int}|0,1}^u$: so $\mathcal{P}_{0,1}^u \simeq \mathcal{H}_{\text{int}|0,1} \simeq \mathcal{H}_{\text{int}|0,1}^u$.

The *classical relational history* γ_c^u between q_0 and q_1 is the critical point of $S[\gamma^u] = S[\gamma_c^u]$ for $\gamma^u \in \mathcal{P}_{0,1}^u$. By (126)–(127), it is given by the vanishing of the 1-cocycle $a(\mathbf{Y})$ on γ_c^u . It encodes the dressed Euler–Lagrange equations, i.e., the relational dynamics (up to vanishing boundary terms) as seen from within any particle in γ :

$$\begin{aligned} \delta_{\mathbf{Y}} S_{|\gamma_c^u} &= \int_{\gamma_c^u} a(\tilde{\mathbf{Y}}) = \int_I a_{\gamma_c^u}(\tilde{\mathbf{Y}}) = \int_I E(\tilde{\mathbf{Y}}, \gamma_c^u) \equiv 0, \\ \forall \mathbf{Y} \simeq \tilde{\mathbf{Y}} \in \text{Lie}\mathcal{H}_{\text{int}|0,1} &\simeq \text{Lie}\mathcal{H}_{\text{int}|0,1}^u, \end{aligned} \quad (128)$$

where $E(\tilde{\mathbf{Y}}, \gamma^u) := \tilde{\mathbf{Y}}(\gamma) \cdot EL[\gamma^u] d\tau = \delta_{\mathbf{Y}} \gamma^u \cdot EL[\gamma^u] d\tau$ is the equations of motion 1-form, with $EL[\gamma^u]$ the Euler–Lagrange equations for the curve γ^u . The critical curve is s.t. $EL[\gamma_c^u] \equiv 0$. Then, for any $\gamma^u \in \mathcal{P}_{0,1}^u$ there is a unique $\mathbf{Y} \in \mathcal{H}_{\text{int}|0,1}^u$ s.t. $\gamma^u = (\gamma_c^u)^{\mathbf{Y}}$, and by (125) one has

$$S[\gamma^u] = S[\gamma_c^u] + \int_{\gamma_c^u} c(\tilde{\mathbf{Y}}). \quad (129)$$

It is clear that $\gamma_c^u = f_u(\gamma_c)$, i.e., that the variational principle is well-behaved under dressing, meaning the *relational variational principle* (128) indeed produces the relational version of the dynamics obtained via the “bare” variational principle (63). This can be seen from a direct computation, similar to (108): using (107) we have indeed

$$\begin{aligned} \delta_{\mathbf{Y}} S_{|\gamma} &= \int_{\gamma} \mathfrak{L}_{\mathbf{Y}^v} L^u = \int_{\gamma} \mathfrak{L}_{\mathbf{Y}^v} L + \int_{\gamma} \mathfrak{L}_{\mathbf{Y}^v} c(\mathbf{u}) \\ &= \delta_{\mathbf{Y}} S_{|\gamma} + \int_{\gamma} \mathfrak{L}_{\mathbf{Y}^v} c(\mathbf{u}), \quad \forall \mathbf{Y} \in \text{Lie}\mathcal{H}_{\text{int}}, \\ &= \int_{\gamma} a(\mathbf{Y}) + \int_{\gamma} a(\tilde{\mathbf{Y}}, f_u(-)) - a(\mathbf{Y}) = \int_{\gamma} a(\tilde{\mathbf{Y}}, f_u(-)), \end{aligned} \quad (130)$$

which of course reproduces (128). But for $\mathbf{Y} \in \text{Lie}\mathcal{H}_{\text{int}|0,1}$, by (63)–(128), Equation (130) establishes the correspondence between the bare and the relational formulations of the variational principle. Notice the remarkable fact that from the above we see that *both* the bare and relational Euler–Lagrange equations can be obtained from the $\text{Lie}\mathcal{H}_{\text{int}|0,1}$ -transformation (107) of the cocyclic dressing field $c(\mathbf{u})$.

The $\mathcal{G}$ -transformation of $S[\gamma^u] = S[\gamma^u]$ is, by (110) and (122),

$$\begin{aligned} (S[\gamma^u])^Z &:= \int_{\gamma} (L^u)^Z = \int_{\gamma} L^u + c(f_u(-), Z) \\ &= S[\gamma^u] + c_{\gamma^u}(Z), \end{aligned} \quad (131)$$

$$\begin{aligned} \text{or equivalently } S[\gamma^u]^Z &:= \int_I (\ell_{\gamma^u})^Z = \int_I \ell_{\gamma^u} + c_{\gamma^u}(Z) \\ &= S[\gamma^u] + c_{\gamma^u}(Z). \end{aligned}$$

It the relational action from the internal viewpoint of a distinct particle in the same physical kinematical history $(\gamma^u)^Z$ given by (97): Specialising the notation in (131), it is $S[\gamma^u] = S[\gamma^{u_i}] + c(\gamma^{u_i}, Z_{ij}(\gamma))$. It shows the physical reference frame covariance of the relational dynamics.

We can go a step further, establishing the correspondence between the relational formulation of the variational principle as written, e.g., from the viewpoints of the i th and the j th particles. It can be obtained by linearising a computation all but similar to (114), using (113), relating $S[\gamma^u]^Y$ and $S[\gamma^{u_i}]^Y$. Or it is obtained directly from (117), or using (116),

$$\begin{aligned} \delta_Y S_{|\gamma}^{u_j} &= \int_{\gamma} \mathfrak{L}_{Y^v} L^{u_j} = \int_{\gamma} \mathfrak{L}_{Y^v} L^{u_i} + \int_{\gamma} \mathfrak{L}_{Y^v} c(f_{u_i}(-), Z_{ij}) \\ &= \delta_Y S_{|\gamma}^{u_i} + \int_{\gamma} \mathfrak{L}_{Y^v} c(f_{u_i}(-), Z_{ij}), \quad \forall Y \in \text{Lie}\mathcal{H}_{\text{int}}, \\ &= \int_{\gamma} \underline{a(\tilde{Y}^i, f_{u_i}(-))} + \int_{\gamma} a(\tilde{Y}^j, f_{u_j}(-)) \\ &\quad - \underline{a(\tilde{Y}^i, f_{u_i}(-))} = \int_{\gamma} a(\tilde{Y}^j, f_{u_j}(-)), \end{aligned} \quad (132)$$

which reproduces (126), as expected. Remark the similarity with (130), using (108)–(107). And, similarly, for $Y \in \text{Lie}\mathcal{H}_{\text{int}|0,1}$, by (128), Equation (132) establishes the equivalence between the relational formulations of the variational principle written by i th and the j th particles. We also notice the remarkable fact that from the above we see that *both* the relational Euler–Lagrange equations written from the viewpoints of the i th and the j th particles can be obtained from the $\text{Lie}\mathcal{H}_{\text{int}|0,1}$ -transformation (116) of the cocyclic quantity $c(f_{u_i}(-), Z_{ij})$.

The dressed action induces the function $S_{\gamma}^u = S_{\gamma^u} = S[\gamma^u; -, q_0]$ with $\gamma^u(\partial I) = \{q_0, q\}$: It is $\mathcal{H}_{\text{ext}}$ -invariant by construction and cocyclic $\mathcal{H}_{\text{int}} \simeq \mathcal{H}_{\text{int}}^u$ -tensorial, $S_{\gamma^u} \in \Omega_{\text{tens}}^0(\mathcal{E}_{\text{rel}}^u; c_{\gamma^u})$. It is then immediate that its $\mathcal{H}_{\text{int}} \simeq \mathcal{H}_{\text{int}}^u$ -transformation is given by $(S_{\gamma^u})^Y = (S_{\gamma^u})^{\tilde{Y}} = S_{\gamma^u} + c_{\gamma^u}(\tilde{Y})$. For $\gamma^u = \gamma_c^u$, the cocyclic 0-form $S_{\gamma_c^u}$ is the dressed Hamilton Principal Function, and satisfies the dressed Hamilton equation: $\frac{\partial S_{\gamma_c^u}}{\partial t} = -H(\tilde{x}, \frac{\partial S_{\gamma_c^u}}{\partial \tilde{x}}; t)$, where the dressed canonical conjugate momenta is by definition $\tilde{\pi} := \frac{\partial S_{\gamma_c^u}}{\partial \tilde{x}}$ and $H(\tilde{x}, \tilde{\pi}, t)$ is the dressed Hamiltonian.

Defining the $U(1)$ -valued $\mathcal{H}_{\text{int}}^u$ -1-cocycle $C_{\gamma^u}(-, \tilde{Y}) = C_{\gamma}(f_u(-), \tilde{Y}) := \exp\{-\frac{i}{\hbar} c_{\gamma^u}(-, \tilde{Y})\} = \exp\{-\frac{i}{\hbar} \int_{\gamma} c(f_u(-), \tilde{Y})\}$, the

dressed action induces the dressed flat cocyclic connection $\varpi_{0,\gamma}^u := -\frac{i}{\hbar} dS_{\gamma}^u$ on $\mathcal{E}_{\text{rel}}^u$, s.t.

$$\begin{aligned} R_{\tilde{Y}}^* \varpi_{0,\gamma}^u &= \varpi_{0,\gamma}^u + C_{\gamma^u}(-, \tilde{Y})^{-1} dC_{\gamma^u}(-, \tilde{Y}), \\ \iota_{\tilde{Y}^v} \varpi_{0,\gamma}^u &= -\frac{i}{\hbar} \iota_{\tilde{Y}^v} dS_{\gamma}^u = -\frac{i}{\hbar} L_{\tilde{Y}^v} S_{\gamma}^u = -\frac{i}{\hbar} \frac{d}{d\epsilon} R_{\tilde{Y}_\epsilon}^* S_{\gamma}^u \Big|_{\epsilon=0} \\ &= -\frac{i}{\hbar} \frac{d}{d\epsilon} c_{\gamma^u}(-, \tilde{Y}_\epsilon) \Big|_{\epsilon=0} = -\frac{i}{\hbar} \int_{\gamma} a(\tilde{Y}, f_u(-)) \\ &= \frac{d}{d\epsilon} C_{\gamma^u}(-, \tilde{Y}_\epsilon) \Big|_{\epsilon=0}, \end{aligned} \quad (133)$$

which is the dressing of (65): By (123), indeed,

$$\varpi_{0,\gamma}^u = \varpi_{0,\gamma} + C_{\gamma}(\mathbf{u})^{-1} dC_{\gamma}(\mathbf{u}), \quad (134)$$

a special case of (27). It follows that the $\mathcal{H}_{\text{int}} \simeq \mathcal{H}_{\text{int}}^u$ -transformation of $\varpi_{0,\gamma}^u$ is

$$(\varpi_{0,\gamma}^u)^{\tilde{Y}} := \Xi^* \varpi_{0,\gamma}^u = \varpi_{0,\gamma}^u + C_{\gamma^u}(\tilde{Y})^{-1} dC_{\gamma^u}(\tilde{Y}), \quad (135)$$

which is $(dS_{\gamma}^u)^{\tilde{Y}} = dS_{\gamma}^u + dC_{\gamma^u}(\tilde{Y})$, as expected from $\mathcal{H}_{\text{int}} \simeq \mathcal{H}_{\text{int}}^u$ -transformation of S_{γ}^u and $[d, \Xi^*] = 0$ for $\Xi \in \mathcal{H}_{\text{int}}^u$. Since $E(\tilde{Y}; \gamma_c^u) \equiv 0$ by definition of γ_c^u , the verticality property of $\varpi_{0,\gamma_c^u}^u = \varpi_{0,\gamma_c^u}$ gives the dressed conserved Noether charge: In a bundle chart, $\tilde{Y}^v \simeq \chi \frac{\partial}{\partial \tilde{x}}$ and $dS_{\gamma_c^u}^u = \frac{\partial S_{\gamma_c^u}^u}{\partial \tilde{x}} d\tilde{x} + \frac{\partial S_{\gamma_c^u}^u}{\partial t} dt$, so $\iota_{\tilde{Y}^v} \varpi_{0,\gamma_c^u}^u = -\frac{i}{\hbar} \chi \frac{\partial S_{\gamma_c^u}^u}{\partial \tilde{x}} =: -\frac{i}{\hbar} \tilde{Y} \tilde{\pi}$. Finally, notice that (129) implies a relation, analogous to (135): For $\tilde{Y} \in \mathcal{H}_{\text{int}|0,1}$ s.t. $\gamma^u = (\gamma_c^u)^{\tilde{Y}}$, we have

$$\varpi_{0,\gamma}^u = \varpi_{0,\gamma_c^u}^u + C_{\gamma_c^u}(\tilde{Y})^{-1} dC_{\gamma_c^u}(\tilde{Y}). \quad (136)$$

Under the action of $\mathcal{G}$, by (131), we have $(S_{\gamma}^u)^Z = S_{(\gamma^u)^Z} = S_{\gamma}^u + c_{\gamma^u}(Z)$. Specialising, $S_{\gamma}^u = S_{\gamma}^{u_i}$ and $(S_{\gamma}^u)^Z = S_{\gamma}^{u_j}$ are the same object seen from distinct viewpoint within γ . In particular, $(S_{\gamma_c^u}^u)^Z = S_{\gamma_c^u}^{u_j}$ is the dressed HPF described from the j th particle. This induces the cocyclic connection $(\varpi_{0,\gamma}^u)^Z := -\frac{i}{\hbar} d(S_{\gamma}^u)^Z$ on $\mathcal{E}_{\text{rel}}^u = \mathcal{E}_{\text{rel}}^{u_j}$, explicitly,

$$\begin{aligned} (\varpi_{0,\gamma}^u)^Z &= \varpi_{0,\gamma}^u + C_{\gamma^u}(Z)^{-1} dC_{\gamma^u}(Z), \quad \text{i.e.,} \\ (\varpi_{0,\gamma}^{u_j}) &= \varpi_{0,\gamma}^{u_i} + C_{\gamma^{u_i}}(Z_{ij})^{-1} dC_{\gamma^{u_i}}(Z_{ij}). \end{aligned} \quad (137)$$

This is another key instance showing how the discrete group $\mathcal{G}$ relates the relational descriptions on $\mathcal{E}_{\text{rel}}^{u_i}$ and $\mathcal{E}_{\text{rel}}^{u_j}$, and implement the physical reference frame covariance of this description.

After this thorough presentation of relational classical mechanics, we can now turn to the formulation of relational quantum mechanics.

4.3. Relational Quantum Mechanics, and Physical Frame Covariance

We define the dressed pre-wave function, using (25)–(27) or the DFM rule of thumb,

$$\psi_{\gamma}^u := \int_u^* \psi_{\gamma} = C_{\gamma}(\mathbf{u})^{-1} \psi_{\gamma}, \quad (138)$$

where $C_\gamma(\mathbf{u}) := C_\gamma(-, \mathbf{u}) = \exp\{-\frac{i}{\hbar} c_\gamma(-, \mathbf{u}(-))\}$. It is H_{ext} -basic on $\mathcal{E}$ by construction, and $\mathcal{H}_{\text{ext}}$ -invariant as is checked using the $\mathcal{H}_{\text{ext}}$ -transformation of ψ_γ , $\psi_\gamma^X = C_\gamma(X)^{-1} \psi_\gamma$, the fact that by (100) one has $C_\gamma(\mathbf{u})^X = C_\gamma(X)^{-1} C_\gamma(\mathbf{u})$. It is thus a 0-form on $\mathcal{E}_{\text{rel}}^u$, as it is clear from (92), and in bundle coordinates

$$\psi_\gamma^u(p) = \psi_\gamma \circ f_u(p) = \psi_\gamma(p + \mathbf{u}(p)) \simeq \psi_\gamma(t; \bar{x}_1, \dots, \bar{x}_{N-1}) = \psi_\gamma(t, \bar{x}), \quad (139)$$

which also shows that we may write $\psi_\gamma^u = \psi_{\gamma^u}$. Specialising this to, e.g., $\mathbf{u} = \mathbf{u}_i$, it is $\psi_{\gamma^u}(p) \simeq \psi_\gamma(t, \bar{x}^i) = \psi_\gamma(t, \bar{x}_1^i, \dots, \bar{x}_{N-1}^i) = \psi_\gamma(t; x_1 - x_i, \dots, 0, \dots, x_N - x_i)$, i.e., the pre-wave function as seen from the i th particle. Again, residual $\mathcal{H}_{\text{int}} \simeq \mathcal{H}_{\text{int}}^u$ -transformations are naturally expected and easily written, since, from (103), we have

$$C_\gamma(\mathbf{u})^Y = C_\gamma(Y)^{-1} C_\gamma(\mathbf{u}) C_\gamma(Y), \quad (140)$$

so that we find

$$\begin{aligned} (\psi_\gamma^u)^Y &= (\psi_{\gamma^u})^Y = (C_\gamma(\mathbf{u})^Y)^{-1} \psi_\gamma^Y, \\ &= C_{\gamma^u}(\bar{Y})^{-1} C_\gamma(\mathbf{u})^{-1} C_\gamma(Y) \cdot C_\gamma(Y)^{-1} \psi_\gamma, \\ &= C_{\gamma^u}(\bar{Y})^{-1} \psi_{\gamma^u} =: (\psi_{\gamma^u})^{\bar{Y}}. \end{aligned} \quad (141)$$

The dressed pre-wave function is thus a cocyclic tensorial 0-form on $\mathcal{E}_{\text{rel}}^u$, $\psi_\gamma^u = \psi_{\gamma^u} \in \Omega_{\text{tens}}^0(\mathcal{E}_{\text{rel}}^u, C_{\gamma^u})$. So it can also be seen as a section of the cocyclic-associated bundle $E^{C_{\gamma^u}} := \mathcal{E}_{\text{rel}}^u \times_{C_{\gamma^u}} \mathbb{C}$. So that $\Omega_{\text{tens}}^0(\mathcal{E}_{\text{rel}}^u, C_{\gamma^u}) \simeq \Gamma(E^{C_{\gamma^u}})$ is the space of “dressed pre-wave functions.”

The dressed cocyclic connection (134)–(135) induces the cocyclic covariant derivative preserving this space

$$\begin{aligned} \bar{D}_{0,\gamma^u} : \Omega_{\text{tens}}^0(\mathcal{E}_{\text{rel}}^u, C_{\gamma^u}) &\rightarrow \Omega_{\text{tens}}^1(\mathcal{E}_{\text{rel}}^u, C_{\gamma^u}), \\ \psi_{\gamma^u} &\mapsto \bar{D}_{0,\gamma^u} \psi_{\gamma^u} := d\psi_{\gamma^u} + \varpi_{0,\gamma^u} \psi_{\gamma^u}. \end{aligned} \quad (142)$$

It is the dressing of the bare cocyclic covariant derivative; by (134)–(138), we have $\bar{D}_{0,\gamma^u} \psi_{\gamma^u} = C_\gamma(\mathbf{u})^{-1} \bar{D}_{0,\gamma} \psi_\gamma$. From the horizontality of $\bar{D}_{0,\gamma^u} \psi_{\gamma^u}$ follows that in bundle coordinates $-i\hbar \frac{\partial}{\partial \bar{x}} = \frac{\partial S_{\gamma^u}}{\partial \bar{x}}$. By (136), we have that

$$\begin{aligned} \psi_{\gamma^u} &= C_{\gamma_c^u}(\bar{Y})^{-1} \psi_{\gamma_c^u}, \quad \text{and} \quad \bar{D}_{0,\gamma^u} \psi_{\gamma^u} = C_{\gamma_c^u}(\bar{Y})^{-1} \bar{D}_{0,\gamma_c^u} \psi_{\gamma_c^u} \\ \text{for } \bar{Y} &\in \mathcal{H}_{\text{int}|0,1}^u. \end{aligned} \quad (143)$$

The space of dressed wave function is $\mathcal{K}^u := \ker \bar{D}_{0,\gamma^u} \subset \Omega_{\text{tens}}^0(\mathcal{E}_{\text{rel}}^u, C_{\gamma^u})$, i.e., the covariantly constant dressed pre-wave functions. A relational version of QM is encoded in the statement

$$\psi_{\gamma_c^u} \in \mathcal{K}^u, \quad \text{i.e.,} \quad \bar{D}_{0,\gamma_c^u} \psi_{\gamma_c^u} = 0. \quad (144)$$

Since $S_{\gamma_c^u} = S[\gamma_c^u; -, q_0]$ is the dressed HPF, satisfying the relational Hamilton–Jacobi equation, in bundle coordinates we have

$$\bar{D}_{0,\gamma_c^u} \psi_{\gamma_c^u} = d\psi_{\gamma_c^u} + \varpi_{0,\gamma_c^u} \psi_{\gamma_c^u} = 0$$

$$\begin{aligned} \Rightarrow \quad dt \frac{\partial}{\partial t} \psi_{\gamma_c^u} + d\bar{x} \frac{\partial}{\partial \bar{x}} \psi_{\gamma_c^u} - \frac{i}{\hbar} \left(dt \frac{\partial S_{\gamma_c^u}}{\partial t} + d\bar{x} \frac{\partial S_{\gamma_c^u}}{\partial \bar{x}} \right) \psi_{\gamma_c^u} &= 0 \\ \hookrightarrow \quad dt \left(\frac{\partial}{\partial t} \psi_{\gamma_c^u} + \frac{i}{\hbar} H(\bar{x}, \bar{\pi}, t) \psi_{\gamma_c^u} \right) + d\bar{x} \left(\frac{\partial}{\partial \bar{x}} - \frac{i}{\hbar} \bar{\pi} \right) \psi_{\gamma_c^u} &= 0. \end{aligned} \quad (145)$$

This gives the relational version of the quantum mechanical prescription for the momentum operator $\hat{\pi} := -i\hbar \frac{\partial}{\partial \bar{x}}$, from which we get the relational canonical commutation relation: $[\hat{x}, \hat{\pi}] = i\hbar \delta(\bar{x} - \bar{x})$. So, altogether we have

$$\begin{aligned} \bar{D}_{0,\gamma_c^u} \psi_{\gamma_c^u} &\equiv 0 \quad \Rightarrow \quad \hat{\pi} := -i\hbar \frac{\partial}{\partial \bar{x}} \quad \text{and} \\ i\hbar \frac{\partial}{\partial t} \psi_{\gamma_c^u} &= H(\hat{x}, \hat{\pi}, t) \psi_{\gamma_c^u}. \end{aligned} \quad (146)$$

It extends to any $\psi_{\gamma^u} \in \mathcal{K}^u$ by (143). Relational QM is equivalently encoded in $\tilde{\psi}_{\gamma^u} \in \tilde{\mathcal{K}}^u$, where $\tilde{\mathcal{K}}^u := \ker \tilde{\nabla}_0^u$ with $\tilde{\nabla}_0^u : \Gamma(E^{C_{\gamma^u}}) \rightarrow \Omega^1(\mathcal{T}) \otimes E^{C_{\gamma^u}}$ the cocyclic covariant derivative induced on associated bundles $E^{C_{\gamma^u}} \rightarrow \mathcal{T}$.

Dressed Path Integral: The above allows to define the dressed, or relational, version of the PI, defined on $\mathcal{P}_{0,1}^u := \{\gamma^u : I \rightarrow \mathcal{E}_{\text{rel}}^u \mid \gamma^u(\partial I) = \{q_0, q_1\}\} = \{\gamma^u \mid \gamma \in \mathcal{P}_{0,1}\}$ the space of paths with beginning and end points $q_0 = f_u(p_0) \simeq (t_0, \bar{x}_0)$ and $q_1 = f_u(p_1) \simeq (t_1, \bar{x}_1)$. We define the dressed propagator,

$$\begin{aligned} K^u(p, p_0) &:= f_u^* K(p, p_0) = K(q, q_0) := \int_{\mathcal{P}_{0,1}^u} D\gamma^u \psi_{\gamma^u} \\ &= \int_{\mathcal{H}_{\text{int}|0,1}^u} D\bar{Y} \psi_{(\gamma^u)^{\bar{Y}}} = \int_{\mathcal{H}_{\text{int}|0,1}^u} D\bar{Y} C_{\gamma_c^u}(\bar{Y})^{-1} \psi_{\gamma_c^u}, \end{aligned} \quad (147)$$

where $D\gamma^u$ and $D\bar{Y}$ are formal measures on $\mathcal{P}_{0,1}^u$ and $\mathcal{H}_{\text{int}|0,1}^u$, respectively. The dressed pre-wave function depending only on the final point is

$$\psi^u(p) := f_u^* \psi(p) = \psi(q) = \int_{\mathcal{E}_{\text{rel}|t_0}^u} K(q, q_0) \psi(q_0) \text{vol}(\mathcal{E}_{\text{rel}|t_0}^u), \quad (148)$$

with $\text{vol}(\mathcal{E}_{\text{rel}|t_0}^u)$ the volume form of the fiber $\mathcal{E}_{\text{rel}|t_0}^u \subset \mathcal{E}_{\text{rel}}^u$ over $\pi(q_0) = t_0 \in \mathcal{T}$. In bundle coordinates, $\psi(q) \simeq \psi(t, \bar{x})$. Taking the relational classical history $\gamma_r^u = \gamma_c^u$ as reference curve, we write

$$K(q, q_0) := \int_{\mathcal{P}_{0,1}^u} D\gamma^u \psi_{\gamma^u} = \int_{\mathcal{H}_{\text{int}|0,1}^u} D\bar{Y} C_{\gamma_c^u}(\bar{Y})^{-1} \psi_{\gamma_c^u}. \quad (149)$$

Further restricting attention to dressed wave functions $\psi_{\gamma^u} \in \mathcal{K}^u$, which are solutions of $\bar{D}_{0,\gamma^u} \psi_{\gamma^u} = 0$ of the form

$$\psi_{\gamma^u} = \psi_0 \exp \int_{\gamma^u} \varpi_{0,\gamma^u} = \psi_0 \exp \frac{i}{\hbar} S_{\gamma^u} = \psi_0 \exp \frac{i}{\hbar} S[\gamma^u], \quad (150)$$

then the propagator (149) becomes the *relational path integral*

$$K(q, q_0) = \int_{\mathcal{P}_{0,1}^u} D\gamma^u \psi_0 \exp \frac{i}{\hbar} S[\gamma^u]$$

$$= \int_{\mathcal{H}_{\text{int}|0,1}^u} D\tilde{Y} C_{\gamma_c^u}(\tilde{Y})^{-1} \psi_0 \exp \frac{i}{\hbar} S[\gamma_c^u] \\ = \psi_0 \exp \frac{i}{\hbar} S[\gamma_c^u] \int_{\mathcal{H}_{\text{int}|0,1}^u} D\tilde{Y} \exp \frac{i}{\hbar} c_{\gamma_c^u}(\tilde{Y}). \quad (151)$$

Then, $\psi^u(p) = \psi(q)$ in (148) is a relational wave function, e.g., as seen from the i th particle if $\mathbf{u} = \mathbf{u}_i$. As in the bare case, we obtain the splitting of the relational PI in a contribution from the classical history, $\psi_{\gamma_c^u} = \psi_0 \exp \frac{i}{\hbar} S[\gamma_c^u]$, containing the essential dynamical physical information, and a normalisation factor given by functional integration of the 1-cocycle $c_{\gamma_c^u}(\tilde{Y}) := \int_{\gamma_c^u} c(-, \tilde{Y}) = \int_I c(-, \tilde{Y}) \circ \gamma_c^u$.

Reference Frame Covariance: The dressed pre-wave functions support $\mathcal{G}$ -transformations. The most direct way to write it is to notice that from (109) follows that the cocyclic dressing $C_\gamma(\mathbf{u})$ transforms as

$$C_\gamma(\mathbf{u})^Z = C_\gamma(\mathbf{u}) C_{\gamma^u}(\mathbf{Z}), \quad (152)$$

as a special case of (30), while ψ_γ is $\mathcal{G}$ -invariant by construction. A dressed pre-wave function (148) $\mathcal{G}$ -transforms as

$$\left(\psi_\gamma^u \right)^Z = \left(C_\gamma(\mathbf{u})^Z \right)^{-1} (\psi_\gamma)^Z = C_{\gamma^u}(\mathbf{Z})^{-1} C_\gamma(\mathbf{u})^{-1} \psi_\gamma \\ = C_{\gamma^u}(\mathbf{Z})^{-1} \psi_\gamma^u. \quad (153)$$

This is a case of (31), and can also be written as $(\psi_{\gamma^u})^Z = C_{\gamma^u}(\mathbf{Z})^{-1} \psi_{\gamma^u}$. This result is also obtained from defining $\left(\psi_\gamma^u \right)^Z = \psi_\gamma^{(u^Z)} := f_{u^Z}^* \psi_\gamma$ via (94), which is thus the same pre-wave function seen from a different viewpoint. The result (153) showcases the covariance of the relational description under change of physical reference frame. Specialising, e.g., to $\mathbf{u} = \mathbf{u}_i$ and $\mathbf{u}^Z = \mathbf{u}_j = \mathbf{u}_i + \mathbf{Z}_{ij}$, by (93) we have, in bundle coordinates,

$$\psi_\gamma^{u_i}(p) \simeq \psi_\gamma(t, \tilde{x}^j) = \psi_\gamma(t; x_1 - x_j, \dots, x_i - x_j, \dots, 0, \dots, x_N - x_j), \\ \psi_\gamma^{u_i}(p) \simeq \psi_\gamma(t, \tilde{x}^i) = \psi_\gamma(t; x_1 - x_i, \dots, 0, \dots, x_j - x_i, \dots, x_N - x_i), \\ \text{and } \psi_\gamma(t, \tilde{x}^j) = C((t, \tilde{x}^i), \mathbf{Z}_{ij})^{-1} \psi_\gamma(t, \tilde{x}^i). \quad (154)$$

Correspondingly, by (137) we have that

$$\left(\bar{D}_{0,\gamma^u} \psi_{\gamma^u} \right)^Z = C_{\gamma^u}(\mathbf{Z})^{-1} \bar{D}_{0,\gamma^u} \psi_{\gamma^u}, \\ \hookrightarrow \bar{D}_{0,\gamma^{u_i}} \psi_{\gamma^{u_i}} = C_{\gamma^{u_i}}(\mathbf{Z}_{ij})^{-1} \bar{D}_{0,\gamma^{u_i}} \psi_{\gamma^{u_i}}. \quad (155)$$

This shows that $\mathcal{K}^{u_i} := \ker \bar{D}_{0,\gamma^{u_i}} \simeq \mathcal{K}^{u_i} := \bar{D}_{0,\gamma^{u_i}}$, which in turn implies the physical reference frame covariance of the relational description of QM (146)—here displayed on $\mathcal{E}_{\text{rel}}^{u_i}$ and $\mathcal{E}_{\text{rel}}^{u_j}$ —as one should expect, a covariance explicitly controlled by the cocyclic structure. It is clear it similarly controls the frame covariance of the relational PI, via $K^{(u^Z)}(p, p_0)$ and $\psi^{(u^Z)}(p)$, so that, e.g., relational wave functions on $\mathcal{E}_{\text{rel}}^{u_i}$ and $\mathcal{E}_{\text{rel}}^{u_j}$ are, in bundle coordinates,

$$\psi^{u_i}(p) \simeq \psi(t, \tilde{x}^j) = \psi(t; x_1 - x_j, \dots, x_i - x_j, \dots, 0, \dots, x_N - x_j),$$

$$\psi^{u_i}(p) \simeq \psi(t, \tilde{x}^i) = \psi(t; x_1 - x_i, \dots, 0, \dots, x_j - x_i, \dots, x_N - x_i),$$

$$\text{and } \psi(t, \tilde{x}^j) = C((t, \tilde{x}^i), \mathbf{Z}_{ij})^{-1} \psi(t, \tilde{x}^i). \quad (156)$$

Observe that the change of physical reference frame is a unitary transformation of the relational quantum state ψ^u , as it is implemented by the $U(1)$ -valued 1-cocycle $C(\mathbf{Z})$.

5. Conclusion

It is interesting to interpret Equations (153)–(156), key results of the relational formulation of QM via DFM: A first reading would be to say that a reference subsystem s , here a particle $s_i = p_i$, “sees itself” as classical, and only the other subsystems (here the other $N - 1$ particles in the N particles configuration) are described quantum mechanically from its perspective. Yet, the $\mathcal{G}$ -transformations—the transformations of the 2nd kind in the DFM framework—implementing physical frame covariance ensure a “quantum democracy” in that any subsystem is as legitimate as any to witness (write) the quantum dynamics.

A more refined and accurate understanding would be to say that there is no meaningful way for a given reference subsystem $s_i = p_i$ to assign *itself* a relational quantum state: it can only do so for other subsystems, which are assigned a wave function with respect to $s_i = p_i$: the dressed/relational wave function ψ^{u_i} (138)/(148). Said otherwise still, for a given subsystem $s_i = p_i$, it only makes sense to assign a quantum state to the set of relations between itself and the rest of the world, which is what ψ^{u_i} represents. However, the physical reference frame covariance implemented by $\mathcal{G}$ -transformations ensures the equivalence (interchangeability) of any subsystem viewpoint to describe the collective relational quantum dynamics.

The reader will have certainly noticed that the dressed version of QM is very much alike the bare one: The dressed wave function ψ^{u_i} for $N - 1$ particles assigned by the i th particle from within the N particles configuration is not unlike the bare wave function ψ for the N particles configuration assigned by an idealized “external observer,” i.e., an “external” physical system s_{ext} . A benefit of the relational formulation via DFM is to clearly indicate that the initial bare ψ can indeed be understood as the dressed wave function assigned to the “internal” system $s_{\text{int}} = \{N \text{ particles configuration}\}$ by s_{ext} from within the total system $s_{\text{tot}} = s_{\text{int}} \cup s_{\text{ext}} (= \{(N + 1) \text{ particles configuration}\})$.

A further benefit of the DFM approach to relational QM is its built-in treatment of physical reference frame covariance by $\mathcal{G}$ -transformations: It dispels the possible a priori misconception that QM relies on “classical observers/systems” with a special status, showing instead how it naturally instantiate the above mentioned “quantum democracy.” This is broadly in line with heuristic ideas behind the literature on “quantum reference frames” (QRFs), as discussed, e.g., in ref. [21].¹³

The relational formulation of QM via DFM thus shows explicitly something that might be overlooked on first reading: that QM is a description of physics *from within*—without the need for an

¹³ We avoid the terminology “QRF” as in our estimation it is potentially misleading, suggesting a “quantum fuzziness” of the reference subsystem s_i in its own frame.

all-encompassing external viewpoint (“god’s eye view”), nor of a partition between classical measuring devices and quantum systems (“Heisenberg cut”). This is broadly in line with Rovelli’s “relational interpretation” of QM.^[18,37,45]

Remark also that the relational structure of QM we focus on here rests on that of classical mechanics, the DFM treating both in essentially the same way. From which follows that QM is plainly seen to be about quantization of relational d.o.f.: in the simple setup tackled here, the network of relative positions of classical particles.

There are a number of developments that we shall pursue. First, our framework may be naturally generalised to the mechanics of extended bodies, undergoing rotations, or of particles with spin. The latter case would be treated elegantly via bundle supergeometry, thereby going back to Berezin original motivation for the introduction of supergeometry in (established) physics.^[46] Another future natural development will be to extend our framework to write a bundle geometric formulation of relativistic QM—which may, e.g., reproduce and generalise^[47–49]—whose relational version would be obtained via the DFM. However, the immediate follow-up to the present work will be the companion,^[40] where we shall recast classical mechanics as a 1-dimensional “general-relativistic” gauge field theory, to which the DFM as developed in ref. [7] applies. We will thereby highlight another key facet of its relational structure and, in echo to the observation of the previous paragraph, we will establish the notion of *relational quantisation*. A notion we shall then develop for standard general-relativistic Gauge Field Theory.

Appendix A: The Case of Free Particles

We consider a configuration of N free particles represented by a point $p = (p_1, \dots, p_N) \in \mathcal{G}_t$ at time $t \in \mathcal{T}$, and a vector $\mathbf{x}|_p \in T_p\mathcal{G}$ which in a bundle chart reads $\mathbf{x} = \mathbf{x}^x \partial_x + \mathbf{x}^t \partial_t$, with $\mathbf{x}^x \partial_x \in T_p(\mathcal{G}_t) = V_p\mathcal{G} \simeq \mathbb{R}^{3N}$. At another configuration $p' = R_X p = p + X \in \mathcal{G}_t$, with $X \in H$, we have that $R_{X*}\mathbf{x}$ and $\mathbf{x}$ have the same t -component, i.e., the same projection as $\pi_* R_{X*}\mathbf{x} = \pi_* \mathbf{x}$. Furthermore, using the canonical parallel transport on $\mathcal{G}_t \simeq \mathbb{E}^{3N}$, allowing to identify $V_p\mathcal{G} \simeq V_{p'}\mathcal{G} \simeq \mathbb{R}^{3N}$, we may identify $R_{X*}(\mathbf{x}^x \partial_x) \simeq \mathbf{x}^x \partial_x$, so that $R_{X*}\mathbf{x} = \mathbf{x}$. Correspondingly, for $\Xi \in \text{Aut}_v(\mathcal{G})$ generated by $X \in H$ via $\Xi(p) = p + X(p)$, we have by (40) $\Xi_*\mathbf{x} = \mathbf{x} + \{dX(\mathbf{x})\}^v$. We have $dX(\mathbf{x}) = \mathbf{x}^x \partial_x X + \mathbf{x}^t \partial_t X = \mathbf{x}^t X'$, since $\partial_x X = 0$ for $X \in H$ are basic on $\mathcal{G}$ so x -independent, so that $\Xi_*\mathbf{x} \simeq (\mathbf{x}^x + \mathbf{x}^t X') \partial_x + \mathbf{x}^t \partial_t$.

The fiber metric is here the standard flat metric on $\mathbb{R}^{3N} \simeq V\mathcal{G}$, $\langle _, _ \rangle : \mathbb{R}^{3N} \times \mathbb{R}^{3N} \rightarrow \mathbb{R}$, $(w, w') \mapsto \langle w, w' \rangle$. The particles have masses $m = (m_1, \dots, m_N)$, and we define the free Lagrangian on $\mathcal{G}$ as $L|_p = \frac{m}{2} \langle v(\mathbf{x}), v(\mathbf{x}) \rangle|_p dt|_p = \sum_{k=1}^N \frac{m_k}{2} \langle v_k(\mathbf{x}), v_k(\mathbf{x}) \rangle|_p dt|_p$, where $dt \in \Omega^1_{\text{basic}}(\mathcal{G})$ and we defined $v := \frac{dx}{dt}$, so that $v(\mathbf{x}) = \frac{\mathbf{x}^x}{\mathbf{x}^t}$. It gauge transforms as

$$\begin{aligned} L^X &:= \Xi^* L = \frac{m}{2} \langle v(\Xi_* \mathbf{x}), v(\Xi_* \mathbf{x}) \rangle \Xi^* dt \\ &= \frac{m}{2} \langle v(\mathbf{x}) + X', v(\mathbf{x}) + X' \rangle dt \\ &= \frac{m}{2} \langle v(\mathbf{x}), v(\mathbf{x}) \rangle dt + m \left(\langle v(\mathbf{x}), X' \rangle + \frac{1}{2} \langle X', X' \rangle \right) dt \\ &= L + c(p, X). \end{aligned} \quad (\text{A1})$$

One checks that $c(p, X)$ is a 1-cocycle for the action of the structure group H of $\mathcal{G}$:

$$\begin{aligned} c(p, X + Y) &= m \left(\langle v(\mathbf{x}), X' + Y' \rangle + \frac{1}{2} \langle X' + Y', X' + Y' \rangle \right) dt \\ &= m \left(\langle v(\mathbf{x}), X' \rangle + \langle v(\mathbf{x}), Y' \rangle \right. \\ &\quad \left. + \frac{1}{2} \langle X', X' \rangle + \langle X', Y' \rangle + \frac{1}{2} \langle Y', Y' \rangle \right) dt \\ &= c(p, X) + \left(\langle v(\mathbf{x}) + X', Y' \rangle + \frac{1}{2} \langle Y', Y' \rangle \right) dt \\ &= c(p, X) + c(p + X, Y). \end{aligned} \quad (\text{A2})$$

The infinitesimal gauge transformation of L is thus, for $\bar{\chi} = \frac{d}{d\epsilon} X_\epsilon|_{\epsilon=0} \in \text{Lie}H$,

$$\mathfrak{L}_{\bar{\chi}} L = \frac{d}{d\epsilon} c(p, X_\epsilon)|_{\epsilon=0} = m \langle v(\mathbf{x}), \chi' \rangle dt =: a(\chi, p). \quad (\text{A3})$$

Given a kinematical history $\gamma : I \rightarrow \mathcal{G}$, $\tau \mapsto \gamma(\tau) \simeq (t(\tau), x(\tau))$, we have $\dot{\gamma} \simeq \dot{x} \partial_x + i \partial_t$. We have $v(\dot{\gamma}) = \frac{\dot{x}}{i} = x'$, the physical velocity (and $\dot{x} = x'$ for the choice $i = 1$). By (56) we have $\Xi_* \dot{\gamma} = \dot{\gamma} + \{\mathbf{x}\}^v \simeq (\dot{x} + i X') \partial_x + i \partial_t$, so $v(\Xi_* \dot{\gamma}) = x' + X'$. The restriction of L on γ is $\ell_\gamma := \gamma^* L \simeq \frac{m}{2} \langle x', x' \rangle \gamma^* dt = \frac{m}{2} \langle \dot{x}, \dot{x} \rangle i^{-1} d\tau$, which is the so-called “parametrized” Lagrangian—reducing for $i = 1$ to the standard unparametrized $\ell_\gamma = \frac{m}{2} \langle x', x' \rangle dt$. The H -transformation of ℓ_γ is, by (54) and above,

$$\begin{aligned} \ell_\gamma^X &= \ell_\gamma + c_\gamma(X) = \ell_\gamma + c(\gamma, X(\gamma)) \\ &\simeq \ell_\gamma + m \left(\langle \dot{x}, \dot{x} \rangle + \frac{1}{2} \langle \dot{x}, \dot{x} \rangle \right) i^{-1} d\tau. \end{aligned} \quad (\text{A4})$$

The linearisation of which gives

$$\begin{aligned} \delta_\chi \ell_\gamma &= a_\gamma(\chi) = a(\chi(\gamma), \gamma) = m \langle \dot{x}, \dot{x} \rangle i^{-1} d\tau \\ &= -m \langle x'', \chi \rangle i d\tau + \left(\frac{d}{d\tau} m \langle x', \chi \rangle \right) d\tau, \end{aligned} \quad (\text{A5})$$

where we identify the Euler–Lagrange 1-form $E(\chi, \gamma) = \chi(\gamma) \cdot EL[\gamma] d\tau = -\langle \chi, mx'' \rangle i d\tau$, and the presymplectic potential giving the conserved Noether charge $\langle mx', \chi \rangle$. The gauge transformation in the unparametrized version for $i = 1$ reads

$$\ell_\gamma^X = \ell_\gamma + m \left(\langle x', X' \rangle + \frac{1}{2} \langle X', X' \rangle \right) dt. \quad (\text{A6})$$

So, $\delta_\chi \ell_\gamma = a_\gamma(\chi) = -\langle \chi, mx'' \rangle dt + \frac{d}{dt} \langle \chi, mx' \rangle dt$, where we recognise the standard expression of the canonical momentum $\pi = mx'$ and of the Euler–Lagrange 1-form $E(\chi, \gamma) = \chi(\gamma) \cdot EL[\gamma] dt = -\langle \chi, \pi' \rangle dt$.

The Equations (A4)–(A6) give the transformation of the free Lagrangian under the so-called “extended Galilean transformations” (or boosts)^[22] as discussed in Section 3.1. Standard Galilean boosts are reproduced for $X(t) = vt$ with $v \in H$, in which

case the Lagrangian is quasi-invariant—i.e., invariant up to a boundary term—as

$$\begin{aligned}\ell_\gamma^X &= \ell_\gamma + m \left(\langle x', v \rangle + \frac{1}{2} \langle v, v \rangle \right) dt \\ &= \ell_\gamma + \frac{d}{dt} \left(m \left(\langle x, v \rangle + \frac{1}{2} \langle v, v \rangle t \right) \right) dt \\ &=: \ell_\gamma + \frac{d}{dt} \Delta((t, x), v) dt.\end{aligned}\quad (A7)$$

This reproduces Equation (3.1.5) in ref. [50], where $\Delta((t, x), v)$ is called a 1-cocycle for the Galilean group, satisfying a special case of the defining property (46)–(A2) of $c(p, X)$ and $c_\gamma(X)$. The rigid limit $X \rightarrow X$ of (A4)/(A6) gives the trivial H -equivariance of ℓ_γ , $R_X^* \ell_\gamma = \ell_\gamma$, i.e., the well-known invariance of the free Lagrangian under translations.

The action functional is $S[\gamma] = \int_\gamma L = \int_\gamma \ell_\gamma = \int_I \frac{m}{2} \langle \dot{x}, \dot{x} \rangle t^{-1} d\tau$, or, in the unparametrised form, $S[\gamma] = \int \frac{m}{2} \langle x', x' \rangle dt$. Its H -transformation is

$$\begin{aligned}S[\gamma]^X &= S[\gamma] + \int_I c_\gamma(X) \\ &\simeq S[\gamma] + \int_I m \left(\langle \dot{x}, \dot{X} \rangle + \frac{1}{2} \langle \dot{X}, \dot{X} \rangle \right) t^{-1} d\tau \\ &\simeq S[\gamma] + \int m \left(\langle x', X' \rangle + \frac{1}{2} \langle X', X' \rangle \right) dt, \quad \text{for } t = 1.\end{aligned}\quad (A8)$$

Which linearises as

$$\begin{aligned}\delta_\chi S[\gamma] &= \int_I a_\gamma(\chi) \\ &\simeq \int_I -m \langle x'', \chi \rangle t d\tau + \left(\frac{d}{d\tau} m \langle x', \chi \rangle \right) d\tau \\ &\simeq \int -\langle \chi, m x'' \rangle dt + \frac{d}{dt} \langle \chi, m x' \rangle dt, \quad \text{for } t = 1, \\ &= \int \chi(\gamma) \cdot EL[\gamma] dt + \langle \chi(\gamma), \pi \rangle \Big|_{t_0}^{t_1}, \quad \text{with } \chi(\gamma) = \delta_\chi \gamma.\end{aligned}\quad (A9)$$

From which we obtain the variational principle: $\delta_\chi S[\gamma_c] \equiv 0$ for any $\chi \in \text{Lie} \mathcal{H}_{0,1}$, giving the equation of motion $EL[\gamma_c] = m x'' = \pi' = 0$, and the conserved Noether charge π .

The wave function for N free particles is $\psi_\gamma(p) \simeq \psi(t, x) = \psi(t; x_1, \dots, x_N)$ and its H -transformation is

$$\begin{aligned}\psi_\gamma^X &= C_\gamma(X)^{-1} \psi_\gamma = \exp \left\{ \frac{i}{\hbar} \int_I c_\gamma(X) \right\} \psi_\gamma, \\ \hookrightarrow \psi(t, x + X) &= \exp \left\{ \frac{i}{\hbar} \int_I m \left(\langle \dot{x}, \dot{X} \rangle + \frac{1}{2} \langle \dot{X}, \dot{X} \rangle \right) t^{-1} d\tau \right\} \psi(t, x) \\ &= \exp \left\{ \frac{i}{\hbar} \int m \left(\langle x', X' \rangle + \frac{1}{2} \langle X', X' \rangle \right) dt \right\} \psi(t, x), \\ &\quad \text{for } t = 1.\end{aligned}\quad (A10)$$

For the special case $X(t) = vt$, by (A7) this reduces to

$$\begin{aligned}\psi(t, x + X) &= \exp \left\{ \frac{i}{\hbar} \int \frac{d}{dt} \Delta((t, x), v) dt \right\} \psi(t, x) \\ &= \exp \left\{ \frac{i}{\hbar} \Delta((t, x), v) \right\} \psi(t, x) \\ &= \exp \left\{ \frac{i}{\hbar} m \left(\langle x, v \rangle + \frac{1}{2} \langle v, v \rangle t \right) \right\} \psi(t, x),\end{aligned}\quad (A11)$$

reproducing Equation (3.1.9) in ref. [50], the well-known transformation of the wave function under Galilean boosts.

Relational Description via the DFM: Here we aim only to illustrate in a simple case how the relational, dressed, Lagrangian ℓ_γ^u and physical frame covariance may be computed using the cocycle. Taking, for concreteness, first the viewpoint of the i th particle, i.e., the dressing $u(p) = u_i(p) = -x_i$, we have,

$$c_\gamma(u_i) \simeq \sum_{k=1}^N m_k \left(\langle \dot{x}_k, \dot{u}_i \rangle + \frac{1}{2} \langle \dot{u}_i, \dot{u}_i \rangle \right) t^{-1} d\tau. \quad (A12)$$

Using (119), we thus have the dressed Lagrangian

$$\begin{aligned}\ell_\gamma^{u_i} &= \ell_\gamma + c_\gamma(u_i) \simeq \sum_{k=1}^N \frac{m_k}{2} \left(\langle \dot{x}_k, \dot{x}_k \rangle + 2 \langle \dot{x}_k, \dot{u}_i \rangle + \langle \dot{u}_i, \dot{u}_i \rangle \right) t^{-1} d\tau \\ &= \sum_{k=1}^N \frac{m_k}{2} \left(\langle \dot{x}_k, \dot{x}_k \rangle - 2 \langle \dot{x}_k, \dot{x}_i \rangle + \langle \dot{x}_i, \dot{x}_i \rangle \right) t^{-1} d\tau \\ &= \sum_{k=1}^N \frac{m_k}{2} \langle \dot{x}_k - \dot{x}_i, \dot{x}_k - \dot{x}_i \rangle t^{-1} d\tau \\ &= \sum_{k=1}^N \frac{m_k}{2} \langle \dot{\tilde{x}}_k^i, \dot{\tilde{x}}_k^i \rangle t^{-1} d\tau = \frac{m}{2} \langle \dot{\tilde{x}}^i, \dot{\tilde{x}}^i \rangle t^{-1} d\tau \\ &= \sum_{\substack{k=1 \\ k \neq i}}^N \frac{m_k}{2} \langle \dot{\tilde{x}}_k^i, \dot{\tilde{x}}_k^i \rangle t^{-1} d\tau = \ell_{\gamma^{u_i}}.\end{aligned}\quad (A13)$$

This Lagrangian describes the relational dynamics of $N - 1$ particles with respect to the i th particle. It is manifestly $\mathcal{H}_{\text{ext}}$ -invariant, and its residual transformation (1st kind) under $\mathcal{H}_{\text{int}}^{u_i} \simeq \mathcal{H}_{\text{int}}$ is, by (120), analogous to (A4):

$$\begin{aligned}(\ell_\gamma^{u_i})^{\tilde{Y}} &= \ell_\gamma^{u_i} + c_{\gamma^{u_i}}(\tilde{Y}) = \ell_\gamma^{u_i} + c(\gamma^{u_i}, \tilde{Y}(\gamma)) \\ &\simeq \ell_\gamma^{u_i} + m \left(\langle \dot{\tilde{x}}^i, \dot{\tilde{Y}}^i \rangle + \frac{1}{2} \langle \dot{\tilde{Y}}^i, \dot{\tilde{Y}}^i \rangle \right) t^{-1} d\tau,\end{aligned}\quad (A14)$$

with $\tilde{Y}^i = Y - Y_i = (Y_1 - Y_i, \dots, 0, \dots, Y_N - Y_i)$ by (86) and (96). It is easily checked directly by plugging $\tilde{x}^{\tilde{Y}} = \tilde{x} + \tilde{Y}$ in the expression (A13) for $\ell_\gamma^{u_i}$.

Similarly, the dressed wave function for free particles is, by (138)–(139),

$$\psi_{\gamma}^{u_i} = C_{\gamma}(\mathbf{u}_i)^{-1} \psi_{\gamma} \simeq \psi(t, \bar{x}^i) = \psi(t, \bar{x}_1^i, \dots, 0, \dots, \bar{x}_N^i), \quad (\text{A15})$$

and describes the relational quantum state between the i th particle and the $N - 1$ others. It is by construction $\mathcal{H}_{\text{ext}}$ -invariant, and its $\mathcal{H}_{\text{int}}^{u_i} \simeq \mathcal{H}_{\text{int}}$ -transformation is, by (141), similar to (A10):

$$\begin{aligned} (\psi_{\gamma}^{u_i})^{\bar{Y}} &= C_{\gamma}(\bar{Y})^{-1} \psi_{\gamma} = \exp \left\{ \frac{i}{\hbar} \int_I c_{\gamma}(\bar{Y}) \right\} \psi_{\gamma}, \\ \hookrightarrow \psi(t, \bar{x}^i + \bar{Y}^i) &= \exp \left\{ \frac{i}{\hbar} \int_I m \left(\langle \dot{\bar{x}}^i, \dot{\bar{Y}}^i \rangle + \frac{1}{2} \langle \dot{\bar{Y}}^i, \dot{\bar{Y}}^i \rangle \right) t^{-1} d\tau \right\} \\ &\psi(t, \bar{x}^i). \end{aligned} \quad (\text{A16})$$

We may now look at how physical frame covariance is implemented by $\mathcal{G}$ -transformations. By (97)–(98), the change from the reference frame of the i th particle to that of the j th particle is given by $\gamma^{u_j} = (\gamma^{u_i})^{Z_{ij}} = \gamma^{u_i} + Z_{ij}(\gamma)$ for $Z_{ij} = x_i - x_j \in \mathcal{G}$. Then, by (122), we have

$$\begin{aligned} \ell_{\gamma}^{u_j} &= \ell_{\gamma}^{u_i} + c(\gamma^{u_i}, Z_{ij}(\gamma)) \\ &= \ell_{\gamma}^{u_i} + m \left(\langle \dot{\bar{x}}^i, \dot{Z}_{ij} \rangle + \frac{1}{2} \langle \dot{Z}_{ij}, \dot{Z}_{ij} \rangle \right) t^{-1} d\tau \\ &= \sum_{k=1}^N \frac{m_k}{2} \left(\langle \dot{x}_k, \dot{x}_k \rangle - 2 \langle \dot{x}_k, \dot{x}_i \rangle + \langle \dot{x}_i, \dot{x}_i \rangle \right) t^{-1} d\tau \\ &\quad + m_k \left(\langle \dot{x}_k - \dot{x}_i, \dot{x}_i - \dot{x}_j \rangle + \frac{1}{2} \langle \dot{x}_i - \dot{x}_j, \dot{x}_i - \dot{x}_j \rangle \right) t^{-1} d\tau \\ &= \sum_{k=1}^N \frac{m_k}{2} \left(\langle \dot{x}_k, \dot{x}_k \rangle - \cancel{2 \langle \dot{x}_k, \dot{x}_i \rangle} + \langle \dot{x}_i, \dot{x}_i \rangle \right. \\ &\quad + \cancel{2 \langle \dot{x}_k, \dot{x}_i \rangle} - 2 \langle \dot{x}_k, \dot{x}_j \rangle - \cancel{2 \langle \dot{x}_i, \dot{x}_i \rangle} + \cancel{2 \langle \dot{x}_i, \dot{x}_j \rangle} \\ &\quad \left. + \langle \dot{x}_i, \dot{x}_i \rangle - 2 \langle \dot{x}_i, \dot{x}_j \rangle + \langle \dot{x}_j, \dot{x}_j \rangle \right) t^{-1} d\tau \\ &= \sum_{k=1}^N \frac{m_k}{2} \langle \dot{x}_k - \dot{x}_j, \dot{x}_k - \dot{x}_j \rangle t^{-1} d\tau \\ &= \sum_{k=1}^N \frac{m_k}{2} \langle \dot{\bar{x}}_k^j, \dot{\bar{x}}_k^j \rangle t^{-1} d\tau = \frac{m}{2} \langle \dot{\bar{x}}^j, \dot{\bar{x}}^j \rangle t^{-1} d\tau, \end{aligned} \quad (\text{A17})$$

showing explicitly that the cocyclic structure controls the physical frame covariance. The frame covariance of the wave function is, by (154)–(156),

$$\begin{aligned} \psi_{\gamma}^{u_j} &= (\psi_{\gamma}^{u_i})^{Z_{ij}} = C_{\gamma^{u_i}}(Z_{ij})^{-1} \psi_{\gamma}^{u_i} = \exp \left\{ \frac{i}{\hbar} \int_I c_{\gamma^{u_i}}(Z_{ij}) \right\} \psi_{\gamma}^{u_i}, \\ \hookrightarrow \psi(t, \bar{x}^j) &= \exp \left\{ \frac{i}{\hbar} \int_I m \left(\langle \dot{\bar{x}}^i, \dot{Z}_{ij} \rangle + \frac{1}{2} \langle \dot{Z}_{ij}, \dot{Z}_{ij} \rangle \right) t^{-1} d\tau \right\} \psi(t, \bar{x}^i). \end{aligned} \quad (\text{A18})$$

It describes the relational quantum state between the j th particle and the $N - 1$ others. It is by construction $\mathcal{H}_{\text{ext}}$ -invariant, and its $\mathcal{H}_{\text{int}}^{u_j} \simeq \mathcal{H}_{\text{int}}$ is, *mutatis mutandis*, the same as (A16). We thus see the “relational quantum democracy” at play, whereby any subsystem can be used to describe the collective quantum state, and change of subsystem is implemented by $\mathcal{G}$ -transformations via the cocyclic structure.

All the above applies identically for a configuration of N harmonic oscillators, whose dynamics is given by the Lagrangian $L = \frac{m}{2} \langle v, v \rangle dt - \frac{k}{2} \langle x, x \rangle dt = \sum_{l=1}^N \frac{m_l}{2} \langle v_l, v_l \rangle dt - \frac{k_l}{2} \langle x_l, x_l \rangle dt$, with spring constants $k = (k_1, \dots, k_N)$, whose $\mathcal{H}$ -transformation is given by the 1-cocycle $c(p, X) := m \left(\langle v, X' \rangle + \frac{1}{2} \langle X', X' \rangle \right) dt - k \left(\langle x, X \rangle + \frac{1}{2} \langle X, X \rangle \right) dt$.

Acknowledgements

J.F. is supported by the Austrian Science Fund (FWF), grant [P 36542], and by the Czech Science Foundation (GAČR), grant GA24-10887S. L.R. is supported by the research grant GrIFOS, funding scheme PNRR Young Researchers, MSCA Seal of Excellence (SoE), CUP E13C24003600006, ID SOE2024_0000103, of which this paper is part. She would also like to thank the University of Graz (Graz, Austria), the Department of Mathematics and Statistics and the Department of Physics at the Faculty of Science of MUNI (Brno, Czechia) for the hospitality during her visits in which part of this work was developed.

Open access publishing facilitated by Politecnico di Torino, as part of the Wiley - CRUI-CARE agreement.

Conflict of Interest

The authors declare that they have no conflicts of interest.

Data Availability Statement

Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

Keywords

Bundle geometry, Cocyclic connection, Physical frame covariance, Quantum mechanics, Relationality

Received: August 7, 2025

Revised: August 7, 2025

- [1] N. M. J. Woodhouse, *Geometric Quantization*, Oxford Mathematical Monographs, 2nd edn, Oxford University Press, Oxford, UK 1998.
- [2] T. W. B. Kibble, *Commun. Math. Phys.* **1979**, 65, 189.
- [3] T. A. Schilling, *PhD Thesis*, The Pennsylvania State University (United States – Pennsylvania) 1996, <https://www.proquest.com/dissertations-theses/geometry-quantum-mechanics/docview/304309562/se-2?accountid=28840>.
- [4] A. Ashtekar, T. Schilling, in *On Einstein's Path*, Springer, New York, NY 1999.
- [5] J. Janyška, M. Modugno, *An Introduction to Covariant Quantum Mechanics*, Fundamental Theories of Physics, 1st edn, Springer, Cham, Switzerland 2022.
- [6] J. François, L. Ravera, *Found. Phys.* **2025**, 55, 38.
- [7] J. T. François, L. Ravera, *Fortschr. Phys.* **2024**, 73, 2400149.
- [8] J. François, L. Ravera, *J. High Energy Phys.* **2024**, 2024, 41.
- [9] J. François, L. Ravera, *Phys. Rev. D* **2025**, 111, 12.
- [10] J. François, *Adv. Theor. Math. Phys.* **2021**, 25, 1389.
- [11] J. Attard, J. François, *Class. Quantum Grav.* **2017**, 34, 8.
- [12] J. Attard, J. François, *Adv. Theor. Math. Phys.* **2018**, 22, 1831.
- [13] P. A. M. Dirac, *Phys. Z. Sowjetunion* **1933**, 3, 64.
- [14] R. P. Feynman, *Rev. Mod. Phys.* **1948**, 20, 367.

- [15] R. P. Feynman, in *Nobel Lectures, Physics 1963-1970*, Elsevier Publishing Company, Amsterdam, <https://www.nobelprize.org/prizes/physics/1965/feynman/lecture/>, **1965**.
- [16] J. François, Thesis, Aix-Marseille Université (Marseille, iFrance), **2014**, <https://hal.archives-ouvertes.fr/tel-01217472>.
- [17] J. François, *J. Phys. A: Math. Theor.* **2024**, 57, 30.
- [18] C. Rovelli, *Int. J. Theor. Phys.* **1996**, 35, 1637.
- [19] Y. Aharonov, T. Kaufherr, *Phys. Rev. D* **1984**, 30, 368.
- [20] S. D. Bartlett, T. Rudolph, R. W. Spekkens, *Rev. Mod. Phys.* **2007**, 79, 555.
- [21] F. Giacomini, E. Castro-Ruiz, C. Brukner, *Nat. Commun.* **2019**, 10, 494.
- [22] D. M. Greenberger, *Am. J. Phys.* **1979**, 47, 35.
- [23] V. Guillemin, S. Sternberg, *Supersymmetry and Equivariant de Rham Theory*, Springer-Verlag, Berlin, Heidelberg **1999**.
- [24] C. Fournel, J. François, S. Lazzarini, T. Masson, *Int. J. Geom. Methods Mod. Phys.* **2014**, 11, 1450016.
- [25] J. François, *J. High Energy Phys.* **2021**, 2021, 225.
- [26] M. Zajac, *J. Geom. Mech.* **2023**, 15, 128.
- [27] A. Riello, M. Schiavina, *Annales Henri Poincaré* **2024**.
- [28] J. François, *Philos. Sci.* **2019**, 86, 472.
- [29] P. Berghofer, J. François, S. Friederich, H. Gomes, G. Hetzroni, A. Maas, R. Sondenheimer, *Gauge Symmetries, Symmetry Breaking, and Gauge-Invariant Approaches*, Elements in the Foundations of Contemporary Physics, Cambridge University Press, Cambridge, UK **2023**.
- [30] W. J. Wolf, J. Read, N. J. Teh, *Found. Phys.* **2022**, 53, 3.
- [31] A. Trautman, *Fiber Bundles, Gauge Field and Gravitation, in General Relativity and Gravitation*, Vol. 1, Plenum Press, New-York, NY **1979**.
- [32] C. V. Westenholz, *Acta Phys. Acad. Sci. Hungaricae* **1980**, 48, 213.
- [33] S. Sternberg, *Group Theory and Physics*, Cambridge University Press, Cambridge, UK **1994**.
- [34] P. Berghofer, J. François, *Found. Phys.* **2024**, 54, 72.
- [35] C. Rovelli, *Class. Quantum. Grav.* **1991**, 8, 297.
- [36] C. Rovelli, *Phys. Rev. D* **2002**, 65, 124013.
- [37] C. Rovelli, *Quantum gravity*, Cambridge Monographs on Mathematical Physics. Cambridge University Press, Cambridge, UK **2004**.
- [38] C. Rovelli, *Found. Phys.* **2014**, 44, 91.
- [39] J. Tamborino, *SIGMA* **2012**, 8, 017.
- [40] J. François, L. Ravera, Mechanics as a “general-relativistic” gauge field theory and Relational Quantization. *In preparation* **2025**.
- [41] J. Attard, J. François, S. Lazzarini, T. Masson, *Foundations of Mathematics and Physics one Century After Hilbert: New Perspectives*, chapter The dressing field method of gauge symmetry reduction, a review with examples, Springer, Berlin **2018**.
- [42] M. Göckeler, T. Schücker, *Differential Geometry, Gauge Theory and Gravity*, Cambridge Monographs on Mathematical Physics, Cambridge University Press, Cambridge, UK **1987**.
- [43] L. Bonora, *Fermions and Anomalies in Quantum Field Theories*, Theoretical and Mathematical Physics, 1st edition, Springer, Cham, Switzerland **2023**.
- [44] R. A. Bertlmann, *Anomalies In Quantum Field Theory*, International Series of Monographs on Physics, vol. 91, Oxford University Press, Oxford, UK **1996**.
- [45] C. Rovelli, F. Vidotto, *Covariant Loop Quantum Gravity: An Elementary Introduction to Quantum Gravity and Spinfoam Theory*, Cambridge University Press, Cambridge, UK **2014**.
- [46] F. Berezin, M. Marinov, *Ann. Phys.* **1977**, 104, 336.
- [47] D. W. Chiou, *Class. Quantum. Grav.* **2013**, 30, 125004.
- [48] E. Seidewitz, *J. Math. Phys.* **2006**, 47, 112302.
- [49] I. H. Redmount, W. M. Suen, *Int. J. Mod. Phys. A* **1993**, 08, 1629.
- [50] J. D. Azcarraga, J. M. Izquierdo, *Lie Groups, Lie Algebras, Cohomology and some Applications in Physics*, Cambridge Monographs on Mathematical Physics, Cambridge University Press, Cambridge, UK **1995**.