

Biomechanical Stabilization of the Keratoconic Cornea: a Review of Current Therapeutic Strategies

Original

Biomechanical Stabilization of the Keratoconic Cornea: a Review of Current Therapeutic Strategies / Ragonese, G.M., Grosso, E., Carbonaro, D., Torta, E., Lepore, E., Gallo, D.. - In: FRONTIERS IN MEDICAL TECHNOLOGY. - ISSN 2673-3129. - ELETTRONICO. - 8:(2026). [10.3389/fmedt.2026.1919880]

Availability:

This version is available at: 11583/3015783 since: 2026-09-22T09:59:40Z

Publisher:

Frontiers Media

Published

DOI:10.3389/fmedt.2026.1919880

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)



OPEN ACCESS

EDITED BY

Sridhar Poosapadi Arjunan,
RMIT University, Australia

REVIEWED BY

Matteo Frigelli,
University of Bern, Switzerland
Ana Mercedes Garcia Albusua,
Asociacion Para Evitar la Ceguera en
México Hospital Dr. Luis Sánchez Bulnes,
Mexico
Bora Deniz Argon,
University of Health Sciences, Türkiye

*CORRESPONDENCE

Diego Gallo

✉ diego.gallo@polito.it

RECEIVED 25 June 2026

REVISED 06 September 2026

ACCEPTED 16 September 2026

PUBLISHED 14 October 2026

CITATION

Ragonese GM, Grosso E, Carbonaro D,
Torta E, Lepore E and Gallo D (2026)
Biomechanical stabilization of the
keratoconic cornea: a review of current
therapeutic strategies.
Front. Med. Technol. 8:1919880.
doi: 10.3389/fmedt.2026.1919880

COPYRIGHT

© 2026 Ragonese, Grosso, Carbonaro,
Torta, Lepore and Gallo. This is an open-
access article distributed under the
terms of the [Creative Commons
Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use,
distribution or reproduction in other
forums is permitted, provided the
original author(s) and the copyright
owner(s) are credited and that the
original publication in this journal is
cited, in accordance with accepted
academic practice. No use, distribution
or reproduction is permitted which does
not comply with these terms.

Biomechanical stabilization of the keratoconic cornea: a review of current therapeutic strategies

Graziana M. Ragonese^{1,2}, Edoardo Grosso², Dario Carbonaro³,
Elena Torta¹, Emiliano Lepore² and Diego Gallo^{1*}

¹Polito^{BIO}Med Lab, Department of Mechanical and Aerospace Engineering, Politecnico di Torino, Torino, Italy, ²Recornea Srl, Trieste, Italy, ³Department of Mechanical, Energetic and Management Engineering, Università della Calabria, Arcavacata di Rende, Italy

Keratoconus is a progressive corneal ectatic disorder characterized by stromal thinning and biomechanical weakening of the cornea, leading to progressive corneal protrusion and visual impairment. Therapeutic strategies have evolved from visual correction toward approaches specifically targeting the biomechanical instability underlying disease progression. This review provides a biomechanical perspective on current and emerging strategies for the biomechanical stabilization of the keratoconic cornea. Conservative treatments based on contact lenses remain essential for visual rehabilitation but do not modify the underlying ectatic process. Corneal collagen crosslinking represents the reference treatment for progressive keratoconus, as it increases stromal stiffness through the formation of additional covalent bonds. However, its efficacy depends on treatment protocol, epithelial status, corneal thickness, and disease stage. Corneal implants, i.e., synthetic intracorneal ring segments and corneal allogenic intrastromal ring segments, provide stretch-induced geometric improvement of the corneal shape and, consequently, of visual acuity, although predictability remains limited. Combined approaches, such as ring implantation followed by cross-linking, aim to integrate corneal reshaping with intrinsic stromal reinforcement. Emerging approaches, including cell-based therapies, and implantation of stromal lenticule or tissue-engineered scaffolds, suggest a future shift toward a multimodal therapeutic strategy integrating biomechanical stabilization, restoration of stromal structure, and biological regeneration. Advancing keratoconus management will require objective biomechanical assessment and predictive modeling tools to guide individualized treatment planning and improve outcome predictability. Ultimately, future management should increasingly incorporate a biomechanical perspective, with interventions designed to modify the mechanical behavior of the ectatic cornea, arrest progression, and restore functional corneal integrity.

KEYWORDS

collagen crosslinking, corneal biomechanics, ectatic cornea, intracorneal ring segments, keratoconus, visual rehabilitation

1 Introduction

Keratoconus (KCN) is a progressive corneal ectasia characterized by stromal thinning (1) and biomechanical weakening, which reduces the ability of the cornea to maintain its physiological curvature under intraocular pressure (IOP). This leads to conical protrusion and consequent vision distortion (2). Corneal shape is maintained by the highly ordered

organization of stromal collagen fibrils. In KCN, this organization is disrupted and has been associated with reduced natural collagen crosslinking, further impairing structural integrity (3). A recent global systematic review and meta-analysis estimated a prevalence of approximately 289 cases per 100,000 individuals (0.24%) and an incidence of 4.0 per 100,000 person-years, with markedly higher prevalence in Africa and South Asia (4). Disease onset typically occurs during the second decade of life with mild corneal thinning and slight refractive changes (5), and it may progress until the fourth decade, when corneal morphology generally stabilizes (6). As the ectasia advances, increasing corneal irregularity leads to visual disturbances such as glare, halos, photophobia, monocular diplopia, and progressive visual blurring (7). End-stage disease may range from mild refractive errors to severe astigmatism, myopia, or visual loss (8). As KCN worsens, surgical intervention becomes necessary in 15% to 20% of affected individuals (9). Corneal transplantation (i.e., keratoplasty) has been the mainstay of surgical treatment for patients with advanced KCN. There is, however, a considerable shortage of corneal graft tissue, with only 1 human donor cornea available for 70 needed (10). Moreover, in about 10% of cases keratoplasty is associated with a long-term risk of immune-mediated endothelial rejection, endothelial cell loss, and complications such as expulsive hemorrhage and endophthalmitis (11).

Several procedures and implants have been developed to biomechanically stabilize the keratoconic cornea, thereby delaying or potentially avoiding the need for keratoplasty. These approaches range from external mechanical support, through intrinsic reinforcement of the corneal stroma, to structural implants that reshape and stabilize the cornea (Table 1). In the early stages of the disease, KCN management relies on conservative approaches, mainly spectacles and rigid contact lenses (CLs), which can provide satisfactory refractive and visual correction (12) but do not address the underlying biomechanical instability of the cornea. Corneal collagen crosslinking (CXL) was subsequently introduced as an early-stage therapeutic option to halt keratoconus progression by inducing additional cross-links within the collagen matrix and thus increasing corneal biomechanical stiffness (13). In recent years, minimally invasive surgical options such as intracorneal ring segments (ICRSs) made of polymethyl methacrylate (PMMA) or derived from allogenic tissue (corneal allogenic intrastromal ring segments, CAIRS) have also been proposed (14). These approaches act on the corneal periphery to induce central corneal flattening, thereby reducing corneal steepening in keratoconic eyes.

While recent reviews have thoroughly investigated keratoconus treatment strategies and their clinical outcomes, less focus has been placed on comparing these approaches according to their biomechanical effects. This mini-review summarizes current strategies aimed at stabilizing the keratoconic cornea from a biomechanical perspective, distinguishing external optical/mechanical support, intrinsic stromal reinforcement, and stromal implants that redistribute corneal stress and reshape the keratoconic cornea. Throughout this review, corneal stiffening, geometric flattening, visual improvement, and disease stabilization are treated as distinct outcomes. Stiffening reflects increased resistance to deformation, flattening reflects changes in corneal shape, visual improvement reflects functional benefit, and

stabilization requires longitudinal evidence of slowed or arrested progression. Therefore, geometric or visual improvement should not be considered equivalent to true stabilization.

This narrative mini-review was based on a targeted literature search conducted primarily through PubMed. Search terms included combinations of “keratoconus”, “corneal biomechanics,” and keywords related to the principal therapeutic approaches for keratoconus. Additional treatment-specific terms were combined with biomechanical keywords to identify studies addressing the effects of each intervention on corneal mechanical behavior.

2 Optical rehabilitation and external biomechanical support with conservative treatments

In the early stages of KCN, visual acuity can often be corrected with spectacles; however, as the disease progresses, their effectiveness declines and CLs become the primary option for visual rehabilitation (15). Fitting CLs in ectatic corneas is challenging and depends on disease severity, cone location, and patient tolerance (16). The mainstay of visual correction is represented by rigid gas permeable (RGP) CLs, typically manufactured from combinations of silicone and acrylate-based materials (17). Their rigid surface enables the formation of a tear lens between the contact lens and the irregular cornea, which neutralizes a large portion of the corneal irregular astigmatism (18). However, the high rigidity of RGP CLs (Young's modulus in the range 1 - 1.9 MPa) (19–21), has been associated with reduced contact area with the cornea, high contact pressure, and poor centration and conformity in highly irregular corneas (22). Excessive contact pressure remains a critical factor that may lead to discomfort, epithelial damage, corneal deformation (22), disruption of the tear film and high frictional shear stress on the corneal surface (21, 23). To mitigate these issues, soft contact lenses based on hydrogel materials have been considered (24). Nevertheless, their tendency to drape over the irregular corneal surface often limits visual improvement (25). Furthermore, their lower stiffness (Young's modulus 0.4 to 0.6 MPa) (20) can make handling more difficult and may reduce adequate lens movement on the eye, thereby limiting effective tear exchange beneath the lens (26). Alternative lens designs have been developed for patients with poor tolerance or inadequate fitting of conventional RGP CLs. Hybrid lenses combine a rigid central zone with a soft peripheral skirt to preserve the optical benefits of RGP CLs while improving centration and comfort (27), whereas piggyback systems place a rigid CL over a soft silicone-hydrogel lens (28, 29). In more advanced or irregular KCN, scleral lenses may be considered, as they vault the cornea and limbus, creating a fluid reservoir that masks irregular astigmatism and reduces direct corneal contact (30–34). However, limited tear exchange may lead to complications such as midday fogging, epithelial bogging, and inflammatory responses (35).

Overall, CLs primarily improve visual function by providing optical correction and, depending on lens design and fitting, may offer external structural support (34). However, they do not intrinsically reinforce the ectatic stroma or modify the underlying ectatic process. If improperly fitted, CLs may induce

TABLE 1 Comparative overview of established and emerging approaches for keratoconus management.

	Approach	Mechanism of Action	Biomechanical Impact	Clinical Outcomes	Challenges/Limitations	Biomechanical Reinforcement	Geometrical Modification
Established approaches	Contact Lenses (CLs)	Formation of a tear lens between the CL and the cornea that masks corneal irregularity and improves the optical surface	Possible external mechanical support, depending on CL design and fitting; no intrinsic stromal reinforcement	Improved visual quality and reduction of irregular astigmatism in mild disease	No modification of the underlying ectatic process; intolerance or poor fitting in advanced disease; risk of epithelial damage or corneal thinning if improperly fitted	 Very low: external support only	 Very low: no direct reshaping
	Corneal Collagen Cross-Linking (CXL)	Riboflavin/UV-A-induced formation of additional covalent bonds within the stromal collagen matrix	Direct increase in stromal stiffness; increased resistance to ectatic deformation	Proven efficacy in slowing or halting disease progression, with long-term stabilization reported up to 10 years	Limited corneal reshaping; protocol-dependent efficacy; possible complications include haze, infection, and delayed epithelial healing	 Very high: direct stromal stiffening	 Very low: no direct reshaping
	Intracorneal Ring Segment (ICRS)/ Corneal Allogenic Intrastromal Ring Segment (CAIRS)	Stromal lamellar separation with arc shortening effect leading to central cornea flattening	Redistribution of internal stresses, with corneal flattening mainly driven by geometric remodeling rather than intrinsic tissue stiffening	Improvement in corneal shape and visual acuity	Variable outcomes; empirical nomograms; limited predictability; minimal intrinsic stromal stiffening except possibly with continuous designs; risk of migration or extrusion	 Moderate: structural support	 High: corneal flattening
	Combined ICRS + CXL	Combination of implant-induced geometric remodeling with CXL-induced stromal stiffening	Potentially synergistic effect of structural reshaping by ICRS and increased stromal rigidity by CXL	Enhanced biomechanical stability through combined corneal reshaping and reduction of ectatic progression	Optimal timing and sequencing not standardized; variable patient-specific response	 Very high: combined stiffening and support	 High: corneal flattening
Emerging approaches	Gene Therapy/Cell-Based Approaches	Targeted modulation of disease-related pathways and/or restoration of the keratocyte function	Potential restoration of stromal homeostasis and ECM organization	Potential modulation of ECM turnover and tissue repair; visual improvement remains unproven clinically	Experimental stage; lack of robust clinical evidence	 High: potential stromal stiffening through tissue remodeling	 Moderate: potential geometrical reshaping
	Stromal Lenticule Implantation	Intrastromal implantation of a lenticule to increase stromal thickness and modify corneal curvature	Direct structural reinforcement through tissue addition; improved stress distribution	Increased corneal thickness with variable or moderate corneal flattening	Limited long-term data; surgical complexity; donor tissue variability.	 High: potential tissue reinforcement	 High: potential corneal flattening
	Tissue-Engineered Scaffolds	Replacement or reinforcement of stromal tissue using biosynthetic scaffolds that mimic the corneal ECM	Structural support with potential biological integration and ECM-like stromal remodeling	Corneal re-epithelialization; stromal thickness restoration	Early clinical stage; biomaterial optimization required; variability in integration and long-term stability	 High: potential stromal stiffening through tissue remodeling	 High: potential corneal flattening

CXL, corneal crosslinking; ECM, extracellular matrix; ICRS, intracorneal ring segments. The red-to-green scale qualitatively indicates the relative contribution of each strategy to biomechanical reinforcement or geometric modification. Biomechanical reinforcement, geometric modification, visual improvement, and disease stabilization are distinct outcomes. Biomechanical reinforcement refers to intrinsic or structural resistance to deformation; geometrical modification refers to changes in corneal shape; clinical outcomes refer mainly to visual or refractive improvement; and disease stabilization requires longitudinal evidence of slowed or arrested progression. The ratings represent a synthesis based on the dominant mechanism of action, the available experimental or biomechanical data, and the available clinical evidence. Low indicates absent, limited, or temporary contribution; moderate indicates partial or indirect contribution; high indicates a direct or more established contribution. For emerging approaches, the ratings reflect proposed or potential effects, as clinical evidence remains limited.

epithelial injury, discomfort, disruption of the tear film, or other corneal surface complications, which may ultimately contribute to worsening of the condition (28, 35). Thus, CLs should be regarded primarily for the management of KCN symptoms, underscoring the need for additional strategies aimed at biomechanical reinforcement of the cornea.

3 Intrinsic stromal biomechanical reinforcement: corneal crosslinking (CXL)

Corneal biomechanical stability is primarily governed by the molecular structure of collagen and its spatial organization within the stromal matrix (36). On this principle, CXL aims to stabilize the ectatic cornea by inducing the formation of additional covalent bonds between collagen fibrils via riboflavin-mediated stromal photoactivation with ultraviolet-A (UV-A) irradiation, enhancing corneal stiffness and biomechanical integrity (37) and thereby slowing or halting the progression of KCN. CXL can be performed with epithelial removal (epithelium-off, “epi-off”) to maximize stromal riboflavin penetration, or with the epithelium left intact (transepithelial, “epi-on”) to reduce postoperative pain and complications. These two approaches are discussed in the followings.

The standard “epi-off” approach, known as the Dresden protocol, involves removal of the corneal epithelium, followed by riboflavin saturation and UV-A irradiation at 370 nm with an irradiance of 3 mW/cm² for 30 min, delivering a total fluence of 5.4 J/cm² (38). This process generates reactive oxygen species, which induce covalent bond formation between collagen fibrils, resulting in increased corneal rigidity and resistance to ectatic deformation (13). The biomechanical effect of standard epi-off CXL has been demonstrated in both experimental and clinical settings. *Ex vivo* stress-strain testing showed an approximately 4.5-fold increase in corneal rigidity (Young’s modulus increase from 1.3 MPa to 5.9 MPa) (39). *In vivo* Corvis ST measurements suggest a possible increase in biomechanical resistance after CXL, with stiffness parameter at first applanation (SP-A1), an indicator of the resistance of the cornea to deformation, increasing from 75.7 mmHg/mm to 77.5 mmHg/mm (40). However, SP-A1 is influenced by intraocular pressure and corneal thickness and should therefore be interpreted cautiously. The efficacy of the standard epi-off technique in stopping the progression of KCN is well established by both short- and long-term studies, with sustained stabilization of KCN and maintained improvements in clinical outcomes reported for up to 10 years of follow-up (41–43).

Several modifications of standard CXL have been introduced, including accelerated protocols that increase UV-A irradiance while reducing exposure time without altering the total energy delivered (44), or protocols with pulsed UV-A delivery to enhance stromal oxygen availability during irradiation, thereby potentially augmenting the biomechanical stiffening effect (7). Direct comparisons among CXL protocols should be interpreted cautiously, as available studies differ in experimental model, measurement method, oxygen availability during irradiation, and follow-up duration that may influence the reported treatment effects.

Transepithelial CXL (T-CXL) is performed without epithelial removal (“epi-on”) to accelerate recovery and reduce pain and complications associated with debridement, including infectious keratitis (45). However, tight epithelial junctions limit stromal riboflavin diffusion during the procedure, potentially limiting its effectiveness in the biomechanical reinforcement of the cornea with respect to the standard epi-off technique (46). In this regard, *in vivo* studies in rabbit corneas have shown that Young’s modulus increased by approximately 21.3% following epi-on CXL (from 9.81 MPa to 11.9 MPa), compared with an increase of approximately 102.5% after standard epi-off CXL (from 9.91 MPa to 19.86 MPa) (47). This observation may also explain the inferior impact on limiting corneal ectasia progression of epi-on vs. epi-off CXL (48, 49).

Iontophoresis-assisted CXL (I-CXL) has been proposed as an alternative approach to enhance riboflavin penetration into intact corneal tissue by means of a low-intensity electric current applied to a negatively charged, low-molecular weight riboflavin solution (46, 50). However, stress-strain analysis performed through uniaxial tensile testing of corneal strips demonstrated that I-CXL significantly increases corneal stiffness in terms of Young’s modulus relative to untreated controls, although to a lesser extent than standard CXL (I-CXL: 17.2 ± 8.2 MPa; standard CXL: 18.8 ± 7.2 MPa; untreated control: 8.6 ± 3.5 MPa) (51). As a consequence, its corneal flattening effect is less pronounced than that observed with standard CXL, with a 2-year mean keratometry (Kmean) reduction of 2.1% for I-CXL compared to 4.3% for standard CXL (52). Although the primary therapeutic goal of CXL is to arrest ectatic progression through stromal stiffening, corneal flattening may occur as a secondary outcome (53–55). This effect is less predictable than stabilization and depends on disease stage, cone location, corneal thickness, epithelial remodeling, and treatment protocol. Customized CXL protocols have been investigated to modulate this biomechanical response by delivering spatially selective UV-A irradiation based on individual corneal topography (53, 54). In a prospective study, a flattening of ≥2.0 D in 37% of eyes treated with personalized CXL compared to 11% after conventional CXL ($P = 0.03$) was reported, suggesting a more pronounced flattening effect with personalized irradiation patterns (53), while patient-specific computational models are emerging to predict CXL-induced corneal shape changes and guide individualized treatment planning (55). Overall, standard (epi-off) CXL remains the reference solution for progressive KCN, whereas T-CXL and I-CXL protocols offer improved tolerability but generally show weaker or less consistent biomechanical effects (47, 51). In thin corneas, standard CXL is limited by endothelial safety considerations. Modified approaches, including hypo-osmolar riboflavin swelling (56), contact lens-assisted CXL (57), individualized fluence protocols such as Sub400 (13), and stromal expansion with refractive lenticules (58) have been proposed to extend biomechanical reinforcement to thin ectatic corneas. In particular, clinical evidence for contact lens-assisted approaches is expanding. A recent retrospective study evaluating accelerated contact lens-assisted CXL using a UV-filter-free Nelfilcon A daily hydrogel lens in thin corneas reported improved best-corrected visual acuity, modest keratometric flattening, stable endothelial cell counts, and only transient clinically insignificant haze during 6 months

of follow-up (59). However, the efficacy of these modified protocols in providing biomechanical reinforcement and their long-term comparability with standard epi-off CXL remain areas of active investigation.

4 Biomechanical reinforcement with implants: synthetic and allogenic ring segments

Implants can be broadly divided into synthetic PMMA ring segments (ICRS) and biologic allogenic corneal tissue segments (CAIRS). They are inserted with a minimally invasive procedure into the deep corneal stroma, within stromal tunnels or a corneal pocket. Unlike CXL, which primarily increases stromal material stiffness, these implants act as spacer elements between stromal collagen lamellae, inducing an arc-shortening effect that flattens the central area of the cornea (60–64). Computational biomechanical models based on finite element analysis have shown that the flattening effect observed after ICRS implantation is directly proportional to the thickness of the segment and inversely proportional to the corneal diameter (65). Additionally, ICRS implantation induces stretch in the strain-stiffening collagen lamellae at the corneal periphery that increases the effective structural stiffness, contributing to limit corneal protrusion (66). However, the main measurable effect of ICRS/CAIRS is geometric remodeling, whereas their contribution to long-term disease stabilization remains supported by limited evidence and depends on whether such increase in effective structural stiffness is sufficient to reduce further ectatic deformation. Clinical decision-making is guided by implantation nomograms, which define parameters such as number of segments, arc length, thickness, and implantation site. However, these nomograms are largely empirical, often based on unpublished clinical data, and lack correspondence with predictions of corneal biomechanical response (62). A variety of ICRS designs with different geometries, diameters, and thicknesses are currently available, including Intacs, KeraRing, Ferrara Ring, MyoRing, and more recently the allogenic option CAIRS (67).

The main characteristics of ICRSs are summarized in Table 2. In detail, Intacs (Addition Technology Inc, Des Plaines, IL, USA), KeraRing (Mediphacos Belo Horizonte, Brazil), and Ferrara segments (Mediphacos Belo Horizonte, Brazil) are crescent-shaped PMMA implants inserted into stromal tunnels to induce corneal flattening through peripheral tissue addition and arc-shortening (68, 69). Differences in cross-sectional geometry, arc length, thickness, and implantation diameter modulate the flattening effect and visual outcome (Table 2) (65). Clinical studies generally report reductions in keratometry and improvements in corrected visual acuity (Table 2), although the magnitude of response varies according to cone morphology, disease severity, tunnel depth, segment design, and nomogram selection (70–72). Importantly, available biomechanical data suggest that synthetic ICRS mainly provide structural support rather than directly increasing intrinsic stromal stiffness.

The continuous 360° configuration of MyoRing (DIOPTEx GmbH, Linz, Austria) provides a circumferential support effect that differs from segmental crescent-shaped ICRSs. Unlike segmental

ICRS, which are implanted in a stromal tunnel, MyoRing is inserted into a corneal pocket (73). Biomechanically, the implant acts as an artificial limbus, redistributing corneal load around the entire corneal periphery and inducing more symmetric central flattening than segmental ICRSs. This effect has been associated with an increase in effective corneal stiffness; however, the magnitude of this increase may depend on the measurement approach used (73, 74). Comparative studies suggest a greater reduction in maximum keratometry (Kmax) with MyoRing (from 51.46 ± 4.38 D to 43.71 ± 1.57 D) compared to KeraRing (from 49.66 ± 3.73 D to 45.04 ± 2.0 D) (75). However, the evidence base remains smaller than for conventional ICRS, and additional comparative studies are needed.



CAIRS may represent a shift from inert mechanical spacing toward biologically integrated stromal reinforcement, but its long-term stabilizing effect and comparative performance against synthetic ICRS require further validation. CAIRS are derived from donor corneal stromal tissue and implanted into stromal channels (76). Their mechanism relies on stromal integration and corneal reshaping through tissue addition, making it compatible also for thinner corneas. Furthermore, CAIRS can be customized in size and shape to better match individual corneal anatomy and ectasia patterns. However, this technique remains relatively recent, with limited long-term clinical data available (77). Additionally, unlike synthetic ICRS, there is no universally accepted nomogram, and the possibility of adjusting parameters such as tunnel depth and width increases procedural complexity (78).

As intrastromal implants and CXL act through different biomechanical mechanisms, combination strategies have been increasingly explored. ICRS or CAIRS can regularize corneal geometry and redistribute stromal stress, whereas CXL can increase stromal material stiffness and reduce the risk of further ectatic progression (79, 80). The theoretical advantage is therefore complementary: reshaping plus biomechanical reinforcement. Available comparative clinical evidence suggests a possible benefit of combined treatment. In a retrospective nonrandomized comparative case series, the addition of CXL to inferior-segment ICRSs implantation resulted in greater functional and topographic improvements compared with ICRSs alone. Specifically, the combined treatment group demonstrated a larger reduction in Kmean (-1.34 ± 1.27 D) versus the ICRS-only group (-0.64 ± 2.40 D). The higher best-corrected visual acuity gain (6.5 lines vs. 0.93 lines) suggests a functional advantage associated with the combined approach (81). Future studies should clarify whether combined procedures produce additive biomechanical stabilization, beyond topographic flattening and refractive benefit.

5 Current challenges and future perspectives

Although current therapeutic strategies can provide external support, increase stromal stiffness, or remodel corneal geometry, important challenges remain, particularly in advanced keratoconus. These include tolerability, limited predictability of refractive outcomes, and uncertain long-term biomechanical stability. Accordingly, current research is increasingly directed

TABLE 2 Comparative characteristics of currently available synthetic and allogenic intrastromal ring implants for keratoconus (intracorneal ring segments, ICRS, and corneal allogenic intrastromal ring segments, CAIRS, respectively).

	Cross-sectional Shape	Arc Length	Available Thicknesses (μm)	Indications	Reported Flattening Effect	Complications
Intacs	 Hexagonal (68)	150° (68)	250 to 450 (68)	Mild-to-moderate keratoconus ($K_{\text{max}} < 58$ D), minimum corneal thickness ≥ 400 μm , central or paracentral cones (100)	K_{max} reduction of approximately 3.8 D in 3 years (70)	Extrusion in 0.4% of cases, associated with trauma (70)
KeraRing	 Triangular (101)	90°, 160°, 210° (101)	150 to 300 (101)	Moderate-to-advanced keratoconus ($46 \leq K_{\text{max}} < 65$ D) with minimum corneal thickness ≥ 400 μm (102)	K_{mean} reduction of 2.24 D at 6 months (71); K_{mean} decreased from 50.63 ± 3.97 D to 47.56 ± 4.46 D at 24 months (73)	Early segment migration in ~6% of eyes; no extrusion reported (71, 103)
Ferrara	 Triangular (104)	90°, 120°, 140°, 160°, 210° (104)	150 to 300 (104)	Progressive keratoconus, increasing corneal steepening, and contact lens intolerance. Contraindicated in advanced disease with apical scarring and keratometry > 60 D (104)	K_{max} decreased from 54.07 D to 48.09 D at 5 years (72); from 56.8 D to 54.4 D in another 5-year study (105)	Extrusion (4.8%), repositioning/exchange (12.9%), visual disturbances (~10%), disease progression (26%) (100, 105)
MyoRing	 Convex to concave (106)	360° (106)	~200 to 400 (106)	Keratoconus with contact lens intolerance, K_{max} 42–65 D, and central pachymetry ≤ 400 μm (75)	K_{mean} decreased from 49.85 D to 47.51 D (-2.34 D) at 3 years (107); greater K_{max} reduction than KeraRing (-7.75 D vs -4.62 D) (75)	No major complications reported; occasional need for additional interventions due to patient dissatisfaction (107)
CAIRS	Not standardized: allogenic corneal stromal tissue; customized uniform, tapered, or asymmetric segments.	Not standardized; variable/customized; reported values of approximately 160° (108, 109).	Not standardized; variable/customized.	Moderate-to-advanced keratoconus (78) with minimum corneal thickness 350 μm (110)–400 μm (109).	K_{max} reduction from 55.85 ± 7.53 D to 50.69 ± 6.38 D at 1 year (78).	No major complications reported in early series; possible segment displacement/repositioning; long-term safety data remain limited (76, 77).

KCN, keratoconus; K_{max} , maximum keratometry; K_{mean} , mean keratometry.

toward strategies that aim not only to stabilize the ectatic cornea, but also to restore stromal thickness, extracellular matrix organization, and corneal function (82, 83).

Among these approaches, cell-based therapies aim at re-establishing the anatomical integrity and physiological function of the keratoconic cornea through restoration of the keratocyte population, i.e., the principal stromal cells (84–86). Corneal stromal stem cells and adipose-derived adult stem cells (ADASCs) have been investigated as potential sources. Early clinical studies using autologous ADASCs (85), decellularized stromal lamellas, or ADASC-recellularized lamellas (87) reported favorable safety profiles but moderate improvements in visual and pachymetric outcomes (88). However, these findings remain preliminary, with limited direct assessment of corneal

biomechanical properties. Complementary strategies such as gene- and growth factor-based approaches aim to modulate molecular pathways involved in stromal cell activity and extracellular matrix turnover. In principle, these strategies could promote controlled tissue repair and matrix reorganization, thereby restoring stromal homeostasis and biomechanical stability of the keratoconic cornea (85). However, their application to keratoconus remains mainly preclinical, and clinical evidence demonstrating biomechanical stabilization is currently lacking.

Additional strategies seek to replace or reinforce diseased stromal tissues. Stromal lenticule addition techniques aim to restore stromal volume by implanting donor or refractive lenticules within a stromal pocket. From a biomechanical

perspective, these procedures may reinforce the ectatic cornea by increasing tissue thickness, modifying corneal curvature, and improving stress distribution (89). When combined with CXL, lenticule implantation combines tissue addition with stromal stiffening. Early clinical studies have reported increased corneal thickness, corneal flattening, and favorable short-term safety profiles (90), but the evidence remains limited by small cohorts and short follow-up, with reports of persistent interface haze beyond 12 months postoperatively (91). Alternatively, tissue-engineering strategies use synthetic, biological, or hybrid scaffolds designed to mimic the corneal extracellular matrix. Synthetic polymers such as polylactic acid, polyglycolic acid or poly(lactic-co-glycolic) acid offer tunable mechanical and degradation properties, whereas biologically derived matrices composed of extracellular components (e.g., collagen, laminin, or chitosan) may provide greater bioactivity and cellular compatibility (85). Collagen-based scaffolds are particularly relevant because collagen is the principal structural component of the corneal stroma, constituting approximately 70% of the corneal extracellular matrix (89), and a major determinant of transparency and biomechanical integrity (89, 92). Early clinical studies of biosynthetic corneal substitutes have shown stromal integration, re-epithelialization, and preservation of corneal transparency in selected patients (93). Nevertheless, their application to keratoconus remains at an early stage, with challenges related to optical transparency, biomechanical strength, degradation behavior, host integration, epithelial healing, and long-term stability.

Emerging therapies for keratoconus converge on two complementary biomechanical goals: restoration of stromal tissue and modulation of biological remodeling. Tissue-addition strategies, such as lenticule implantation and tissue-engineered scaffolds, primarily aim to modify the corneal geometry by increasing stromal volume. Alternatively, cell-based and molecular approaches seek to restore stromal homeostasis and extracellular matrix organization. These approaches suggest a shift toward combined structural, biological, and biomechanical restoration. However, they remain investigational, and their role in biomechanical stabilization will require controlled studies with longer follow-up and standardized biomechanical endpoints capable of quantifying the structural reinforcement of the ectatic cornea. In this sense, future studies investigating novel therapeutic strategies for ectatic corneas should complement conventional clinical outcomes, such as corneal thickness, curvature, and visual acuity, with objective and standardized biomechanical measurements. Quantifications based on, for example, stromal stiffness, corneal hysteresis (94), Brillouin microscopy (95), optical coherence elastography (96), or patient-specific computational modeling (97), would help determine whether a novel therapeutic approach truly reinforces the ectatic cornea, rather than merely improving its thickness or shape.

In conclusion, the therapeutic landscape for keratoconus has progressively evolved from purely refractive compensation toward strategies explicitly targeting corneal biomechanics. Future developments delineate a transition toward a comprehensive, multimodal therapeutic strategy in which biomechanical stabilization, corneal remodeling, and biological regeneration are integrated according to disease stage and

individual corneal behavior. In this context, the digital twin paradigm may represent a promising framework, allowing patient-specific corneal models to integrate tomographic, biomechanical, and clinical data to simulate disease progression, predict treatment outcomes, and support individualized planning of the therapeutic strategy (98, 99). Ultimately, future keratoconus management should increasingly incorporate a biomechanical perspective, with interventions designed to modify the mechanical behavior of the ectatic cornea, arrest progression, and restore functional corneal integrity.

Author contributions

GR: Investigation, Writing – original draft, Writing – review & editing, Formal analysis. EG: Investigation, Writing – review & editing. DC: Conceptualization, Writing – review & editing. ET: Methodology, Writing – review & editing. EL: Conceptualization, Project administration, Supervision, Writing – review & editing. DG: Conceptualization, Funding acquisition, Project administration, Resources, Supervision, Writing – original draft, Writing – review & editing.

Funding

The author(s) declared that financial support was received for this work and/or its publication. This work is part of the project PNRR-NGEU, which has received funding from the MUR – DM 352/2022.

Conflict of interest

The author(s) declared that this work was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

The author DG declared that they were an editorial board member of *Frontiers* at the time of submission. This had no impact on the peer review process and the final decision.

Generative AI statement

The author(s) declared that generative AI was used in the creation of this manuscript solely to support language editing, grammar checking, and improvement of the clarity of the manuscript. The authors critically reviewed and edited all AI-assisted suggestions and take full responsibility for the final content.

Any alternative text (alt text) provided alongside figures in this article has been generated by *Frontiers* with the support of artificial intelligence and reasonable efforts have been made to ensure accuracy, including review by the authors wherever possible. If you identify any issues, please contact us.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated

organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

- Rabinowitz YS. Keratoconus. *Surv Ophthalmol.* (1998) 42(4):297–319. doi: 10.1016/s0039-6257(97)00119-7
- Ambekar R, Toussaint KC, Wagoner Johnson A. The effect of keratoconus on the structural, mechanical, and optical properties of the cornea. *J Mech Behav Biomed Mater.* (2011) 4(3):223–36. doi: 10.1016/j.jmbm.2010.09.014
- Vought R, Greenstein SA, Gelles J, Hersh PS. The pathophysiology of keratoconus. *Cornea.* (2025) 44(2):137–43. doi: 10.1097/ICO.0000000000003585
- Sriranganathan A, Chan CC, Dhillon J, Felfeli T. Global incidence and prevalence of keratoconus: a systematic review and meta-analysis. *Cornea.* (2026) 45:1080–92. doi: 10.1097/ICO.0000000000003973
- Olivares Jiménez JL, Guerrero Jurado JC, Bermudez Rodriguez FJ, Serrano Laborda D. Keratoconus: age of onset and natural history. *Optom Vis Sci.* (1997) 74(3):147–51. doi: 10.1097/00006324-199703000-00025
- Vazirani J, Basu S. Keratoconus: current perspectives. *Clin Ophthalmol.* (2013) 7:2019–30. doi: 10.2147/OPTH.S50119
- Singh RB, Koh S, Sharma N, Woreta FA, Hafezi F, Dua HS, et al. Keratoconus. *Nat Rev Dis Primers.* (2024) 10(1):81. doi: 10.1038/s41572-024-00565-3
- Choi JA, Kim M-S. Progression of keratoconus by longitudinal assessment with corneal topography. *Invest Ophthalmol Vis Sci.* (2012) 53(2):927–35. doi: 10.1167/iops.11-8118
- Liu H, Chen Y, Wang P, Li B, Wang W, Su Y, et al. Efficacy and safety of deep anterior lamellar keratoplasty vs. Penetrating keratoplasty for keratoconus: a meta-analysis. *PLoS One.* (2015) 10(1):e0113332. doi: 10.1371/journal.pone.0113332
- Gain P, Jullienne R, He Z, Aldossary M, Acquart S, Cognasse F, et al. Global survey of corneal transplantation and eye banking. *JAMA Ophthalmol.* (2016) 134(2):167–73. doi: 10.1001/jamaophthalmol.2015.4776
- Thompson RW, Price MO, Bowers PJ, Price FW. Long-term graft survival after penetrating keratoplasty. *Ophthalmology.* (2003) 110(7):1396–402. doi: 10.1016/S0161-6420(03)00463-9
- Deshmukh R, Ong ZZ, Rampat R, Alió del Barrio JL, Barua A, Ang M, et al. Management of keratoconus: an updated review. *Front Med.* (2023) 10:1212314. doi: 10.3389/fmed.2023.1212314
- Hafezi F, Kling S, Hafezi NL, Aydemir ME, Lu N-J, Hillen M, et al. Corneal cross-linking. *Prog Retin Eye Res.* (2025) 104:101322. doi: 10.1016/j.preteyeres.2024.101322
- D'Oria F, Bagaglia SA, Alio Del Barrio JL, Alessio G, Alio JL, Mazzotta C. Refractive surgical correction and treatment of keratoconus. *Surv Ophthalmol.* (2024) 69(1):122–39. doi: 10.1016/j.survophthal.2023.09.005
- Lim L, Lim EWL. Current perspectives in the management of keratoconus with contact lenses. *Eye (Lond).* (2020) 34(12):2175–96. doi: 10.1038/s41433-020-1065-z
- Saraç Ö, Kars ME, Temel B, Çağır N. Clinical evaluation of different types of contact lenses in keratoconus management. *Cont Lens Anterior Eye.* (2019) 42(5):482–6. doi: 10.1016/j.clae.2019.02.013
- Efron N. Rigid lens materials. In: *Contact Lens Practice*. Amsterdam: Elsevier (2024). p. 122–130. doi: 10.1016/B978-0-7020-8427-0.00011-8
- Downie LE, Lindsay RG. Contact lens management of keratoconus. *Clin Exp Optom.* (2015) 98(4):299–311. doi: 10.1111/cxo.12300
- Iqbal A, Thomas R, Mahadevan R. Impact of modulus of elasticity of silicone hydrogel contact lenses on meibomian glands morphology and function. *Clin Exp Optom.* (2021) 104(7):760–6. doi: 10.1080/08164622.2021.1887703
- Young G, Garofalo R, Peters S, Harmer O. The effect of temperature on soft contact lens modulus and diameter. *Eye Contact Lens.* (2011) 37(6):337–41. doi: 10.1097/ICL.0b013e31822e8c3b
- Horst CR, Brodland B, Jones LW, Brodland GW. Measuring the modulus of silicone hydrogel contact lenses. *Optom Vis Sci.* (2012) 89(10):1468–76. doi: 10.1097/OPX.0b013e3182691454
- Ramasubramanian D, Hernández-Verdejo JL, López-Alonso JM. Influence of contact lens parameters on cornea: biomechanical analysis. *Bioengineering (Basel).* (2024) 11(10):966. doi: 10.3390/bioengineering11100966
- Hart SM, McGhee EO, Uruña JM, Levings PP, Eikenberry SS, Schaller MA, et al. Surface gel layers reduce shear stress and damage of corneal epithelial cells. *Tribol Lett.* (2020) 68(4):106. doi: 10.1007/s11249-020-01344-3
- Musgrave CSA, Fang F. Contact lens materials: a materials science perspective. *Materials (Basel).* (2019) 12(2):261. doi: 10.3390/ma12020261
- Rico-Del-Viejo L, García-Montero M, Hernández-Verdejo JL, García-Lázaro S, Gómez-Sanz FJ, Lorente-Velázquez A. Nonsurgical procedures for keratoconus management. *J Ophthalmol.* (2017) 2017:9707650–17. doi: 10.1155/2017/9707650
- Kim E, Saha M, Ehrmann K. Mechanical properties of contact lens materials. *Eye Contact Lens.* (2018) 44(Suppl 2):S148–56. doi: 10.1097/ICL.0000000000000442
- Uçakhan ÖÖ, Yeşiltaş YS. Correction of irregular astigmatism with new-generation hybrid contact lenses. *Eye Contact Lens.* (2020) 46(2):91–8. doi: 10.1097/ICL.0000000000000618
- Şengör T, Aydın Kurna S. Update on contact lens treatment of keratoconus. *Turk J Ophthalmol.* (2020) 50(4):234–44. doi: 10.4274/tjo.galenos.2020.70481
- Kurna A, Sengor T, Ozkurt K, Kurna A, Akı FJ. High dk piggyback contact lens system for contact lens-intolerant keratoconus patients. *Clin Ophthalmol.* (2011) 5:331–5. doi: 10.2147/OPTH.S16727
- Segal O, Barkana Y, Hourovitz D, Behrman S, Kamun Y, Avni I, et al. Scleral contact lenses may help where other modalities fail. *Cornea.* (2003) 22(4):308–10. doi: 10.1097/00003226-200305000-00006
- Michaud L, Lipson M, Kramer E, Walker M. The official guide to scleral lens terminology. *Contact Lens Anterior Eye.* (2020) 43(6):529–34. doi: 10.1016/j.clae.2019.09.006
- Rathi VM, Mandathara PS, Taneja M, Dumpati S, Sangwan VS. Scleral lens for keratoconus: technology update. *Clin Ophthalmol.* (2015) 9:2013–8. doi: 10.2147/OPTH.S52483
- Bergmanson JPG, Walker MK, Johnson LA. Assessing scleral contact lens satisfaction in a keratoconus population. *Optom Vis Sci.* (2016) 93(8):855–60. doi: 10.1097/OPX.0000000000000882
- Visser E-S, Van der Linden BJJJ, Otten HM, Van der Lelij A, Visser R. Medical applications and outcomes of bitangential scleral lenses. *Optom Vis Sci.* (2013) 90(10):1078–85. doi: 10.1097/OPX.0000000000000018
- Walker MK, Bergmanson JP, Miller WL, Marsack JD, Johnson LA. Complications and fitting challenges associated with scleral contact lenses: a review. *Cont Lens Anterior Eye.* (2016) 39(2):88–96. doi: 10.1016/j.clae.2015.08.003
- Raiskup F, Herber R, Lenk J, Pillunat LE, Spoerl E. Crosslinking with UV-A and riboflavin in progressive keratoconus: from laboratory to clinical practice—developments over 25 years. *Prog Retin Eye Res.* (2024) 102:101276. doi: 10.1016/j.preteyeres.2024.101276
- Bui AD, Truong A, Pasricha ND, Indaram M. Keratoconus diagnosis and treatment: recent advances and future directions. *Clin Ophthalmol.* (2023) 17:2705–18. doi: 10.2147/OPTH.S392665
- Santhiago MR, Randleman JB. The biology of corneal cross-linking derived from ultraviolet light and riboflavin. *Exp Eye Res.* (2021) 202:108355. doi: 10.1016/j.exer.2020.108355
- Wollensak G, Spoerl E, Seiler T. Stress-strain measurements of human and porcine corneas after riboflavin-ultraviolet-A-induced cross-linking. *J Cataract Refract Surg.* (2003) 29(9):1780–5. doi: 10.1016/s0886-3350(03)00407-3
- Jabbarvand M, Moravvej Z, Shahraki K, Hashemian H, Ghasemi H, Berijani S, et al. Corneal biomechanical outcome of collagen cross-linking in keratoconic patients evaluated by corvis ST. *Eur J Ophthalmol.* (2021) 31(4):1577–83. doi: 10.1177/1120672120944798
- Bordais J, Cassagne M, Touboul D, Saunier V, Butterworth J, Malecaze F, et al. Conventional epithelial-off corneal crosslinking in patients with progressive keratoconus: 10-year outcomes. *Cornea.* (2024) 43(7):874–82. doi: 10.1097/ICO.0000000000003532
- Nicola C, Pop R, Rednik A, Nicola D. 10-Year Results of standard cross-linking in patients with progressive keratoconus in Romania. *J Ophthalmol.* (2019) 2019:8285649–5. doi: 10.1155/2019/8285649
- Raiskup F, Theuring A, Pillunat LE, Spoerl E. Corneal collagen crosslinking with riboflavin and ultraviolet-A light in progressive keratoconus: ten-year results. *J Cataract Refract Surg.* (2015) 41(1):41–6. doi: 10.1016/j.jcrs.2014.09.033
- Shetty R, Pahuja NK, Nuijts RMMA, Ajani A, Jayadev C, Sharma C, et al. Current protocols of corneal collagen cross-linking: visual, refractive, and tomographic outcomes. *Am J Ophthalmol.* (2015) 160(2):243–9. doi: 10.1016/j.ajo.2015.05.019

45. D'Oria F, Palazón A, Alió JL. Corneal collagen cross-linking epithelium-on vs. Epithelium-off: a systematic review and meta-analysis. *Eye Vis (Lond)*. (2021) 8(1):34. doi: 10.1186/s40662-021-00256-0
46. Subasinghe SK, Ogbuehi KC, Dias GJ. Current perspectives on corneal collagen crosslinking (CXL). *Graefes Arch Clin Exp Ophthalmol*. (2018) 256(8):1363–84. doi: 10.1007/s00417-018-3966-0
47. Wollensak G, Iomdina E. Biomechanical and histological changes after corneal crosslinking with and without epithelial debridement. *J Cataract Refract Surg*. (2009) 35(3):540–6. doi: 10.1016/j.jcrs.2008.11.036
48. Kobashi H, Rong SS, Ciolino JB. Transepithelial versus epithelium-off corneal crosslinking for corneal ectasia. *J Cataract Refract Surg*. (2018) 44(12):1507–16. doi: 10.1016/j.jcrs.2018.08.021
49. Caporossi A, Mazzotta C, Paradiso AL, Baiocchi S, Marigliani D, Caporossi T. Transepithelial corneal collagen crosslinking for progressive keratoconus: 24-month clinical results. *J Cataract Refract Surg*. (2013) 39(8):1157–63. doi: 10.1016/j.jcrs.2013.03.026
50. Vinciguerra P, Montericchio A, Catania F, Fossati G, Raimondi R, Legrottaglie EF, et al. New perspectives in keratoconus treatment: an update on iontophoresis-assisted corneal collagen crosslinking. *Int Ophthalmol*. (2021) 41(5):1909–16. doi: 10.1007/s10792-021-01713-4
51. Cassagne M, Laurent C, Rodrigues M, Galinier A, Spoerl E, Galiacy SD, et al. Iontophoresis transcorneal delivery technique for transepithelial corneal collagen crosslinking with riboflavin in a rabbit model. *Invest Ophthalmol Vis Sci*. (2016) 57(2):594–603. doi: 10.1167/iov.13-12595
52. Bikbova G, Bikbov M. Standard corneal collagen crosslinking versus transepithelial iontophoresis-assisted corneal crosslinking, 24 months follow-up: randomized control trial. *Acta Ophthalmol*. (2016) 94(7):e600–6. doi: 10.1111/aos.13032
53. Seiler TG, Fischinger I, Koller T, Zapp D, Frueh BE, Seiler T. Customized corneal cross-linking: one-year results. *Am J Ophthalmol*. (2016) 166:14–21. doi: 10.1016/j.ajo.2016.02.029
54. Nordström M, Schiller M, Fredriksson A, Behndig A. Refractive improvements and safety with topography-guided corneal crosslinking for keratoconus: 1-year results. *Br J Ophthalmol*. (2017) 101(7):920–5. doi: 10.1136/bjophthalmol-2016-309210
55. Frigelli M, Ariza Gracia MA, Aydemir ME, Torres-Netto EA, Hafezi F, Rozema J, et al. Predicting the effects of customized corneal cross-linking on corneal geometry. *Invest Ophthalmol Vis Sci*. (2025) 66(12):51. doi: 10.1167/iov.66.12.51
56. Raiskup F, Spoerl E. Corneal cross-linking with hypo-osmolar riboflavin solution in thin keratoconic corneas. *Am J Ophthalmol*. (2011) 152(1):28–32.e1. doi: 10.1016/j.ajo.2011.01.016
57. Jacob S, Kumar DA, Agarwal A, Basu S, Sinha P, Agarwal A. Contact lens-assisted collagen cross-linking (CACXL): a new technique for cross-linking thin corneas. *J Refract Surg*. (2014) 30(6):366–72. doi: 10.3928/1081597X-20140523-01
58. Sachdev MS, Gupta D, Sachdev G, Sachdev R. Tailored stromal expansion with a refractive lenticule for crosslinking the ultrathin cornea. *J Cataract Refract Surg*. (2015) 41(5):918–23. doi: 10.1016/j.jcrs.2015.04.007
59. Argon BD, Avcı Ö. Avcı, safety and efficacy of contact lens-assisted accelerated corneal cross-linking using nelfilcon A in thin corneas. *Int Ophthalmol*. (2026) 46(1):73. doi: 10.1007/s10792-026-03943-w
60. Roberts CJ, Dupps WJ. Biomechanics of corneal ectasia and biomechanical treatments. *J Cataract Refract Surg*. (2014) 40(6):991–8. doi: 10.1016/j.jcrs.2014.04.013
61. Hamon L, Schlötzer-Schrehardt U, Flockerzi FA, Seitz B, Daas L. Morphological characterization and clinical effects of stromal alterations after intracorneal ring segment implantation in keratoconus. *Graefes Arch Clin Exp Ophthalmol*. (2022) 260(7):2299–308. doi: 10.1007/s00417-022-05572-2
62. Sakellaris D, Balidis M, Gorou O, Szentmary N, Alexoudis A, Grieshaber MC, et al. Intracorneal ring segment implantation in the management of keratoconus: an evidence-based approach. *Ophthalmol Ther*. (2019) 8(Suppl 1):5–14. doi: 10.1007/s40123-019-00211-2
63. Morales P, Durán JA. Advances in intracorneal ring segment (ICRS) implantation for keratoconus: a comprehensive literature review, clinical insights, and future prospects. *J Clin Med*. (2025) 14(13):4454. doi: 10.3390/jcm14134454
64. Burris TE, Baker PC, Ayer CT, Loomas BE, Mathis ML, Silvestrini TA. Flattening of central corneal curvature with intrastromal corneal rings of increasing thickness: an eye-bank eye study. *J Cataract Refract Surg*. (1993) 19:182–7. doi: 10.1016/s0886-3350(13)80404-x
65. Vega-Estrada A, Alió JL. The use of intracorneal ring segments in keratoconus. *Eye Vis (Lond)*. (2016) 3:8. doi: 10.1186/s40662-016-0040-z
66. Bagheri NM, Mosaddegh P, Kakhkhaei M. Effects of intracorneal ring segments on the biomechanical response of the ectatic cornea to air-puff: a patient-specific numerical analysis. *Math Mech Solids*. (2022) 27(3):390–409. doi: 10.1177/10812865211025581
67. Hayat UK, Shiwani HA, Memon D, Walkden A. Intracorneal ring segments in keratoconus: a narrative literature review. *Clin Ophthalmol*. (2025) 19:2249–57. doi: 10.2147/OPTH.S520174
68. Ertan A, Kamburoğlu G. Intacs implantation using a femtosecond laser for management of keratoconus: comparison of 306 cases in different stages. *J Cataract Refract Surg*. (2008) 34(9):1521–6. doi: 10.1016/j.jcrs.2008.05.028
69. Hamdi IM. Optical and topographic changes in keratoconus after implantation of ferrara intracorneal ring segments. *J Refract Surg*. (2010) 26(11):871–80. doi: 10.3928/1081597X-20100114-05
70. Warrak EL, Serhan HA, Ayash JG, Wahab CH, Baban TA, Daoud RC, et al. Long-term follow up of intracorneal ring segment implantation in 932 keratoconus eyes. *J Fr Ophthalmol*. (2020) 43(10):1020–4. doi: 10.1016/j.jfo.2020.03.019
71. Shabayek MH, Alió JL. Intrastromal corneal ring segment implantation by femtosecond laser for keratoconus correction. *Ophthalmology*. (2007) 114(9):1643–52. doi: 10.1016/j.ophtha.2006.11.033
72. Torquetti L, Berbel RF, Ferrara P. Long-term follow-up of intrastromal corneal ring segments in keratoconus. *J Cataract Refract Surg*. (2009) 35(10):1768–73. doi: 10.1016/j.jcrs.2009.05.036
73. Daxer A. Biomechanics of corneal ring implants. *Cornea*. (2015) 34(11):1493–8. doi: 10.1097/ICO.0000000000000591
74. Daxer A, Ettl A, Hörantner R. Long-term results of MyoRing treatment of keratoconus. *J Optom*. (2017) 10(2):123–9. doi: 10.1016/j.optom.2016.01.002
75. Al-Tuwairqi WS, Osuagwu UL, Razzouk H, AlHarbi A, Ogbuehi KC. Clinical evaluation of two types of intracorneal ring segments (ICRS) for keratoconus. *Int Ophthalmol*. (2017) 37(5):1185–98. doi: 10.1007/s10792-016-0385-2
76. Jacob S, Patel SR, Agarwal A, Ramalingam A, Saijimal AI, Raj JM. Corneal allogenic intrastromal ring segments (CAIRS) combined with corneal cross-linking for keratoconus. *J Refract Surg*. (2018) 34(5):296–303. doi: 10.3928/1081597X-20180223-01
77. Levy I, Mukhija R, Nanavaty MA. Corneal allogenic intrastromal ring segments: a literature review. *J Clin Med*. (2025) 14(4):1340. doi: 10.3390/jcm14041340
78. Asfar KE, Bteich Y, Abou Mrad A, Assaf JF, Jacob S, Hafezi F, et al. Corneal allogenic intrastromal ring segments (CAIRS) versus synthetic segments: a single segment comparative analysis using propensity score matching. *J Refract Surg*. (2024) 40(11):e863–76. doi: 10.3928/1081597X-20241002-02
79. Kankariya VP, Dube AB, Grentzelos MA, Kontadakis GA, Diakonis VF, Petrelli M, et al. Corneal cross-linking (CXL) combined with refractive surgery for the comprehensive management of keratoconus: cXL plus. *Indian J Ophthalmol*. (2020) 68(12):2757–72. doi: 10.4103/ijo.IJO_1841_20
80. Kymionis GD, Grentzelos MA, Portaliou DM, Kankariya VP, Randleman JB. Corneal collagen cross-linking (CXL) combined with refractive procedures for the treatment of corneal ectatic disorders: cXL plus. *J Refract Surg*. (2014) 30(8):566–76. doi: 10.3928/1081597X-20140711-10
81. Chan CCK, Sharma M, Wachler BSB. Effect of inferior-segment intacs with and without C3-R on keratoconus. *J Cataract Refract Surg*. (2007) 33(1):75–80. doi: 10.1016/j.jcrs.2006.09.012
82. Giannaccare G, Lixi F, Slidsborg C, Ozkan G, Gheorghe AG, Arghirescu A-M, et al. Current landscape and future prospects of corneal regenerative medicine. *Ophthalmol Ther*. (2026) 15(1):85–128. doi: 10.1007/s40123-025-01286-w
83. Nubile M, Alió del Barrio JL, Cerino L, Salgari N, El Zarif M, Totta M, et al. Ex Vivo lenticule customization for stromal lenticule advanced keratoplasty. *Invest Ophthalmol Vis Sci*. (2024) 65(10):9. doi: 10.1167/iov.65.10.9
84. Ahadi M, Ramin S, Abbasi A, Tahmouri H, Hosseini SB. Mini review: human clinical studies of stem cell therapy in keratoconus. *BMC Ophthalmol*. (2024) 24:35. doi: 10.1186/s12886-024-03297-w
85. Javed Z, Daigavane S. Harnessing corneal stromal regeneration for vision restoration: a comprehensive review of the emerging treatment techniques for keratoconus. *Cureus*. (2024) 16(9):e69835. doi: 10.7759/cureus.69835
86. Atalay E, Özalp O, Yıldırım N. Advances in the diagnosis and treatment of keratoconus. *Ther Adv Ophthalmol*. (2021) 13:25158414211012796. doi: 10.1177/25158414211012796
87. El Zarif M, Alió JL, Alió del Barrio JL, De Miguel MP, Abdul Jawad K, Makdissy N. Corneal stromal regeneration: a review of human clinical studies in keratoconus treatment. *Front Med*. (2021) 8:650724. doi: 10.3389/fmed.2021.650724
88. El Zarif M, Alió JL, Alió del Barrio JL, Abdul Jawad K, Palazón-Bru A, Abdul Jawad Z, et al. Corneal stromal regeneration therapy for advanced keratoconus: long-term outcomes at 3 years. *Cornea*. (2021) 40(6):741–54. doi: 10.1097/ICO.00000000000002646
89. Fasolo A, Galzignato A, Pedrotti E, Chiarego C, Cozzini T, Bonacci E, et al. Femtosecond laser-assisted implantation of corneal stroma lenticule for keratoconus. *Int Ophthalmol*. (2021) 41(5):1949–57. doi: 10.1007/s10792-021-01739-8
90. Ganesh S, Brar S. Femtosecond intrastromal lenticular implantation combined with accelerated collagen cross-linking for the treatment of keratoconus—initial clinical result in 6 eyes. *Cornea*. (2015) 34(10):1331–9. doi: 10.1097/ICO.0000000000000539
91. Hayashi T, Igarashi A, Fujiwara K, Kobashi H, Son H-S, Kling S, et al. Stromal transplantation and corneal-sparing techniques in ectatic diseases. *Prog Retinal Eye Res*. (2026) 113:101487. doi: 10.1016/j.preteyeres.2026.101487
92. Huang X, Islam MM, Watson SL, Patra HK. Seeing through collagen: integrative pro-regenerative corneal implants for clearer future. *NPJ Regen Med*. (2026) 11:21. doi: 10.1038/s41536-026-00471-0

93. Fagerholm P, Lagali NS, Merrett K, Jackson WB, Munger R, Liu Y, et al. A biosynthetic alternative to human donor tissue for inducing corneal regeneration: 24-month follow-up of a phase 1 clinical study. *Sci Transl Med.* (2010) 2(46):46ra61. doi: 10.1126/scitranslmed.3001022
94. Touboul D, Roberts C, Kérautret J, Garra C, Maurice-Tison S, Saubusse E, et al. Correlations between corneal hysteresis, intraocular pressure, and corneal central pachymetry. *J Cataract Refract Surg.* (2008) 34(4):616–22. doi: 10.1016/j.jcrs.2007.11.051
95. Hammoud B, Schumacher JS, Zhang H, Dutra BAL, Susanna BN, Scarcelli G, et al. Brillouin microscopy for focal biomechanical measurements in normal and keratoconic corneas: a narrative review. *Surv Ophthalmol.* (2026) 71(2):662–73. doi: 10.1016/j.survophthal.2025.06.004
96. De Stefano VS, Ford MR, Seven I, Dupps WJ Jr. Depth-dependent corneal biomechanical properties in normal and keratoconic subjects by optical coherence elastography. *Trans Vis Sci Tech.* (2020) 9(7):4. doi: 10.1167/tvst.9.7.4
97. Sinha Roy A, Dupps WJ Jr. Patient-specific computational modeling of keratoconus progression and differential responses to collagen cross-linking. *Invest Ophthalmol Vis Sc.* (2011) 52(12):9174–87. doi: 10.1167/iops.11-7395
98. Fantaci B, Calvo B, Rodríguez JF. Modeling biological growth of human keratoconus: on the effect of tissue degradation, location and size. *Comput Biol Med.* (2024) 180:108976. doi: 10.1016/j.compbiomed.2024.108976
99. Bagheri NM, Kadkhodaei M, Pirhadi S, Mosaddegh P. Effects of intracorneal ring segments implementation technique and design on corneal biomechanics and keratometry in a personalized computational analysis. *Sci Rep.* (2021) 11(1):14433. doi: 10.1038/s41598-021-93821-5
100. Tlaiss Y, Warrak J, Warrak E. Intrastromal corneal ring segments for keratoconus: a comprehensive review of different types. *Ther Adv Ophthalmol.* (2025) 17:25158414251371521. doi: 10.1177/25158414251371521
101. Coskunseven E, Kymionis GD, Tsiklis NS, Atun S, Arslan E, Siganos CS, et al. Complications of intrastromal corneal ring segment implantation using a femtosecond laser for channel creation: a survey of 850 eyes with keratoconus. *Acta Ophthalmol.* (2011) 89(1):54–7. doi: 10.1111/j.1755-3768.2009.01605.x
102. Dal A, Erdağ M, Canleblebici M. A comparative study of two intrastromal corneal ring segment models in keratoconus: A retrospective study. *Indian J Ophthalmol.* (2025) 73(9):1307–13. doi: 10.4103/IJO.IJO_14_25
103. Coskunseven E, Kymionis GD, Tsiklis NS, Atun S, Arslan E, Jankov MR, et al. One-year results of intrastromal corneal ring segment implantation (KeraRing) using femtosecond laser in patients with keratoconus. *Am J Ophthalmol.* (2008) 145(5):775–9. doi: 10.1016/j.ajo.2007.12.022
104. Torquetti L, Sandes J, Ferrara Intrastromal Corneal Ring Segments. *International Journal of Keratoconus and Ectatic Corneal Diseases.* (2016) 5(3):114–27. doi: 10.5005/jp-journals-10025-1132
105. de Araujo BS, Kubo L, Marinho DR, Kwitko S. Keratoconus progression after intrastromal corneal ring segment implantation according to age: 5-year follow-up cohort study. *Int Ophthalmol.* (2020) 40(11):2847–54. doi: 10.1007/s10792-020-01468-4
106. Hosny M, El-Mayah E, Sidky MK, Anis M. Femtosecond laser-assisted implantation of complete versus incomplete rings for keratoconus treatment. *Clin Ophthalmol.* (2015) 9:121–7. doi: 10.2147/OPTH.S73855
107. Janani L, Jadidi K, Mosavi SA, Nejat F, Naderi M, Nourijelyani K. MyoRing Implantation in Keratoconic Patients: 3 years Follow-up Data. *J Ophthalmic Vis Res.* (2016) 11(1):26–31. doi: 10.4103/2008-322X.180713
108. Nacaroglu SA, Yesilkaya EC, Perk FFNK, Tanriverdi C, Taneri S, Kilic A. Efficacy and safety of intracorneal allogenic ring segment implantation in keratoconus: 1-year results. *Eye (Lond).* (2023) 37(18):3807–12. doi: 10.1038/s41433-023-02618-5
109. Yucekul B, Tanriverdi C, Taneri S, Keskin Perk FFN, Karaca Y, Kilic A. Effect of Corneal Allogenic Intrastromal Ring Segment (CAIRS) Implantation Surgery in Patients With Keratoconus According to Prior Corneal Cross-linking Status. *Journal of Refractive Surgery.* (2024) 40(6):e392–7. doi: 10.3928/1081597X-20240501-01
110. Kirgiz A, Kemer Atik B, Emul M, Taskapili M. Clinical outcomes of femtosecond laser-assisted corneal allogenic intrastromal ring segment (CAIRS) in the treatment of keratoconus. *Clinical & Experimental Ophthalmology.* (2024) 52(7):713–23. doi: 10.1111/ceo.14411